

DELSYM COUGH RELIEF PLUS SOOTHING ACTION- dextromethorphan hydrobromide and menthol lozenge
Reckitt Benckiser LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Delsym®
Cough Relief Plus Soothing Action

Drug Facts

Active ingredients (in each lozenge)	Purpose
Dextromethorphan HBr 5 mg	Cough suppressant
Menthol 5 mg	Oral pain reliever

Uses

temporarily relieves

- sore throat
- sore mouth
- minor mouth irritation
- cough due to minor throat and bronchial irritation as may occur with the common cold

Warnings

Sore throat warning

- If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting consult a doctor promptly.

Do not use in a child under 6 years of age.

Do not use if you are now taking prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that is accompanied by excessive phlegm (mucus)

Stop use and ask a doctor or dentist if

- sore mouth symptoms do not improve in 7 days
- irritation, pain or redness persists or worsens
- swelling develops
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness

If pregnant or breast-feeding, ask a health professional before use.

Keep this and all drugs out of the reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 12 years and older: take 2 lozenges (one immediately after the other) and allow each lozenge to dissolve slowly in the mouth; may be repeated every 4 hours, not to exceed 12 lozenges in any 24-hour period, or as directed by a doctor
- children 6 to under 12 years of age: take 1 lozenge and allow to dissolve slowly in the mouth; may be repeated every 4 hours, not to exceed 6 lozenges in 24 hours, or as directed by a doctor
- children under 6 years of age: do not use

Other information

- tamper evident: product is sealed in blister package for your protection. Do not use if carton is damaged or if printed seal on blister is broken or missing.
- store at room temperature 20–25°C (68–77°F)

Inactive ingredients

D&C yellow #10, FD&C yellow #6, flavors, liquid glucose, propylene glycol, sucralose, sucrose, water

Questions?

Call 1-888-963-3382

You may also report side effects to this phone number.

Dist. by: Reckitt Benckiser
Parsippany, NJ 07054-0224

PRINCIPAL DISPLAY PANEL - 16 Lozenge Carton

NDC 63824-160-16

NEW!

Delsym®

COUGH RELIEF PLUS
SOOTHING ACTION

Dextromethorphan HBr 5 mg (Cough Suppressant)

Menthol 5 mg (Oral Pain Reliever)

- **Quiets cough**
- **Relieves sore throat**
- **Soothing action**

16 LOZENGES

Honey

Lemon



Delsym

COUGH RELIEF PLUS
SOOTHING ACTION

Dextromethorphan HBr 5 mg (Cough Suppressant)
Menthol 5 mg (Oral Pain Reliever)

NDC 63824-160-16

NEW!

Delsym

COUGH RELIEF PLUS
SOOTHING ACTION

Dextromethorphan HBr 5 mg (Cough Suppressant)
Menthol 5 mg (Oral Pain Reliever)

- Quiets cough
- Relieves sore throat
- Soothing action



16 LOZENGES

**Honey
Lemon**



Dextromethorphan HBr 5 mg (Cough Suppressant)
Menthol 5 mg (Oral Pain Reliever)

Delsym

COUGH RELIEF PLUS
SOOTHING ACTION

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Drug Facts

Drug Facts (continued)

Please visit
our web site:
www.delsym.com

COUGH RELIEF PLUS
SOOTHING ACTION

Delsym
Dextromethorphan HBr 5 mg (Cough Suppressant)
Menthol 5 mg (Oral Pain Reliever)

Drug Facts

Active ingredients (in each lozenge)
Dextromethorphan HBr 5 mg...Cough suppressant
Menthol 5 mg.....Oral pain reliever

Purpose

Uses temporarily relieves ■ sore throat
■ sore mouth ■ minor mouth irritation
■ cough due to minor throat and bronchial
irritation as may occur with the common cold

Warnings

Sore throat warning
■ If sore throat is severe, persists for more
than 2 days, is accompanied or followed by
fever, headache, rash, nausea or vomiting
consult a doctor promptly.

Do not use in a child under 6 years of age.
Do not use if you are now taking a prescription
monoamine oxidase inhibitor (MAOI) (certain
drugs for depression, psychiatric, or emotional
conditions, or Parkinson's disease), or for
2 weeks after stopping the MAOI drug. If you
do not know if your prescription drug contains
an MAOI, ask a doctor or pharmacist before
taking this product.

Ask a doctor before use if you have
■ persistent or chronic cough such as occurs
with smoking, asthma, or emphysema
■ cough that is accompanied by excessive
phlegm (mucus)

Stop use and ask a doctor or dentist if
■ sore mouth symptoms do not improve in
7 days
■ irritation, pain or redness persists or worsens
■ swelling develops
■ cough lasts more than 7 days, comes back, or
occurs with fever, rash, or persistent headache.
These could be signs of a serious illness

**If pregnant or breast-feeding, ask a health
professional before use. Keep this and all
drugs out of the reach of children.**

In case of overdose, get medical help or contact
a Poison Control Center right away.

Directions ■ adults and children 12 years
and older: take 2 lozenges (one immediately
after the other) and allow each lozenge to
dissolve slowly in the mouth; may be repeated
every 4 hours, not to exceed 12 lozenges in
any 24-hour period, or as directed by a doctor
■ children 6 to under 12 years of age: take
1 lozenge and allow to dissolve slowly in
the mouth; may be repeated every 4 hours,
not to exceed 6 lozenges in 24 hours, or
as directed by a doctor
■ children under 6 years of age: do not use

Other information

■ tamper evident: product is sealed in blister
package for your protection. Do not use
if carton is damaged or if printed seal on
blister is broken or missing.
■ store at room temperature 20-25°C (68-77°F)

Inactive ingredients D&C yellow #10,
FD&C yellow #6, flavors, liquid glucose,
propylene glycol, sucralose, sucrose, water

Questions? Call 1-888-963-3382

You may also report side effects to this
phone number.



Dist. by: Reckitt Benckiser
Parsippany, NJ 07054-0224 010412
Made in England ©2012 RB 0386858

Lot No.:	Lane 1	Lane 1
	Lane 2	Lane 2
	Lane 3	Lane 3

0386858

DELSYM COUGH RELIEF PLUS SOOTHING ACTION

dextromethorphan hydrobromide and menthol lozenge

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63824-160
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Dextromethorphan Hydrobromide (UNII: 9D2RTI9KYH) (Dextromethorphan - UNII:7355X3ROTS)	Dextromethorphan Hydrobromide	5 mg
Menthol (UNII: L7T10EIP3A) (Menthol - UNII:L7T10EIP3A)	Menthol	5 mg

Inactive Ingredients

Ingredient Name	Strength
D&C yellow No. 10 (UNII: 35SW5USQ3G)	
FD&C yellow No. 6 (UNII: H77VEI93A8)	
propylene glycol (UNII: 6DC9Q167V3)	
sucralose (UNII: 96K6UQ3ZD4)	
sucrose (UNII: C151H8M554)	
water (UNII: 059QF0K00R)	

Product Characteristics

Color	YELLOW	Score	no score
Shape	ROUND	Size	27mm
Flavor	HONEY, LEMON	Imprint Code	D
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63824-160-16	16 in 1 BLISTER PACK		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	07/01/2012	

Labeler - Reckitt Benckiser LLC (094405024)

Revised: 1/2012

Reckitt Benckiser LLC