

**DG HEALTH ANTI DIARRHEAL- loperamide hydrochloride tablet, film coated
Dolgenercorp, LLC**

Dolgenercorp, LLC Anti-Diarrheal Drug Facts

Active ingredient (in each caplet)

Loperamide HCl 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- store at 20° -25° C (68° -77° F)
- see end panel for lot number and expiration date

Inactive ingredients

anhydrous lactose, carnauba wax, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch

Questions or comments?

1-888-309-9030

Principal Display Panel

SEE NEW WARNINGS

Compare to the active ingredient of Imodium® A-D

Loperamide Hydrochloride Tablets, 2 mg

Anti-Diarrheal

Controls the symptoms of diarrhea

Anti-Diarrheal

12 Caplets*

*Capsule-Shaped Tablets

Actual Caplet Size

2 mg

OPEN OTHER END
CODE AREA

22453 VT G1

Drug Facts (continued)
 Inactive ingredients: anhydrous lactose, croscarmellose, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch.
 100% Satisfaction Guarantee (BBB) 309-9030
 Questions or comments? 1-888-309-9030
Drug Facts (continued)

Drug Facts (continued)
Warnings
 Do not use if you have ever had a severe allergic reaction to loperamide HCl. Heart attack: Taking more than directed can cause serious heart problems or death. Do not use if you have bloody or black stools. Ask a doctor before use if you have:
 ■ a history of abnormal heart rhythm
 ■ a history of liver disease
 ■ fever
 ■ muscle aches
 Ask a doctor before use if you have:
 ■ a history of abnormal heart rhythm
 ■ a history of liver disease
 ■ fever
 ■ muscle aches
 If you are taking a prescription drug, Loperamide may interact with certain prescription drugs.
 When using this product, the drowsiness or dizziness may occur. Be careful when driving or operating machinery.
 Stop use and ask a doctor if:
 ■ symptoms get worse
 ■ diarrhea lasts for more than 2 days
 ■ you get abdominal swelling or bloating.
 These may be signs of a serious condition. If pregnant or breastfeeding, ask a health professional before use.

Directions
 Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222, Control Center right away).
Drug Facts (continued)

Active ingredient
 Loperamide HCl 2 mg. Amt. diarr. eal.
Purpose
 Loperamide HCl 2 mg. Amt. diarr. eal.

Use
 Controls symptoms of diarrhea, including Traveler's Diarrhea.

Other information
 ■ Store at 20°-25°C (68°-77°F); excursions permitted to 15°-30°C (59°-86°F) as shown on the label.
 ■ See expiration date for lot number and expiration date.

adults and children 12 years and over	2 caplets after the first stool, 1 caplet thereafter each subsequent dose; do not use more than 4 caplets in 24 hours
children 9-11 years	1 caplet after the first stool, 1/2 caplet thereafter each subsequent dose; do not use more than 3 caplets in 24 hours
children 6-8 years	1 caplet after the first stool, 1/2 caplet thereafter each subsequent dose; do not use more than 2 caplets in 24 hours
children 2-5 years	ask a doctor
children under 2 years (up to 30 lbs)	do not use

3 7003016946 7
 16946
 700301
 3
 A1378

DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN
 DISTRIBUTED BY ODD EAST MAIN CO., 100 MISSION RIDGE, GOODLETTSVILLE, TN 37072.
 *This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Imodium® A-D.

Loperamide Hydrochloride Tablets, 2 mg

Anti-Diarrheal

DG™ health

SEE NEW WARNINGS
 Compare to the active ingredient of Imodium® A-D**

Loperamide Hydrochloride Tablets, 2 mg

Anti-Diarrheal

- Controls the symptoms of diarrhea
- Anti-Diarrheal



12 Caplets*

2 mg

Actual Caplet Size

*Capsule-Shaped Tablets



DG HEALTH ANTI DIARRHEAL

loperamide hydrochloride tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55910-224
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	

Product Characteristics

Color	GREEN	Score	2 pieces
Shape	OVAL	Size	10mm
Flavor		Imprint Code	L2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-224-67	1 in 1 CARTON	12/28/2009	01/31/2022
1		48 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:55910-224-23	1 in 1 CARTON	09/11/2012	12/31/2021
2		72 in 1 BOTTLE; Type 0: Not a Combination Product		

3	NDC:55910-224-72	1 in 1 CARTON	05/02/2017	07/31/2021
3		60 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:55910-224-62	24 in 1 CARTON	02/14/2020	
4		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
5	NDC:55910-224-53	12 in 1 CARTON	07/06/2020	
5		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
6	NDC:55910-224-91	6 in 1 CARTON	07/08/2020	08/31/2023
6		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075232	12/28/2009	

Labeler - Dolgencorp, LLC (068331990)

Revised: 11/2024

Dolgencorp, LLC