

**UMBRELLA- octinoxate titanium dioxide lotion**  
**CA-BOTANA INTERNATIONAL**

-----  
**Umbrella Sunscreen**

**Warning and Precautions Section**

For external use only. Do not use on damaged or broken skin. When using this product keep out of eyes. Rinse with water to remove. Stop use and ask a doctor if rash occurs. Keep out of reach of children. If product is swallowed, get medical help or contact a poison control center right away.

**DOSAGE & ADMINISTRATION SECTION**

Helps prevent sunburn. If used as directed with other sun protection measured decreases the risk of skin cancer and early skin aging caused by the sun. Apply liberally 15 minutes before sun exposure. Use a water resistant sunscreen if swimming or sweating. Reapply: at least 2 hours. Children under 6 months: Ask a doctor. Sun protection measurements. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including: limit time in the sun, especially from 10 a.m - 2 p.m. Wear long-sleeve shirts, pants, hats, and sunglasses. Protect this product from excessive heat and direct sun

**INACTIVE INGREDIENT SECTION**

ALOE VERA LEAF  
ALUMINUM STARCH OCTENYLSUCCINATE  
ASCORBIC ACID  
ASCORBYL PALMITATE  
BORON NITRIDE  
CAMELLIA SINENSIS FLOWER  
CAPRYLIC/CAPRIC MONO/DIGLYCERIDES  
STEARYL ALCOHOL  
CETYL ALCOHOL  
CETEARYL ESNONANOATE  
CITRIC ACID MONOHYDRATE  
CITRUS AURANTIUM FLOWER OIL  
COCO-CAPRYLATE  
ETHYLHEXYLGLYCERIN

GLYCERIN  
SUNFLOWER OIL  
NEACEN  
PANTHENOL  
PHENOXYETHANOL  
EGG PHOSPHOLIPIDS  
POTASSIUM SORBATE  
PROPANEDIOL  
ROSMARINUS OFFICINALIS FLOWERING TOP OIL  
PEG-9 DIGLYCIDYL ETHER/SODIUM HYALURONATE CROSSPOLYMER  
SORBITAN STEARATE  
TOCOPHEROL  
GRAPE SEED OIL  
XANTHAN GUM  
WATER

**OTC - ACTIVE INGREDIENT SECTION**

OCTINOXATE  
TITANIUM DIOXIDE

**OTC - PURPOSE SECTION**

**keep out of reach of children section**

For external use only.

Avoid contact with eyes.

Keep out of reach of children.

Do not apply to open wounds.

STOP USE AND ask a doctor if condition worsens or symptoms persist for more than seven days, discontinue use of the product.

For external use only. Do not use on damaged or broken skin. When using this product keep out of eyes. Rinse with water to remove. Stop use and ask a doctor if rash occurs. Keep out of reach of children. If product is swallowed, get medical help or contact a poison control center right away.

Helps prevent sunburn. If used as directed with other sun protection measured decreases the risk of skin cancer and early skin aging caused by the sun. Apply liberally 15 minutes before sun exposure. Use a water resistant sunscreen if swimming or sweating. Reapply: at least 2 hours. Children under 6 months: Ask a doctor. Sun protection measurements. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including: limit time in the sun, especially from 10 a.m - 2 p.m. Wear long-sleeve shirts, pants, hats, and sunglasses. Protect this product from excessive heat and direct sun

DOCTOR D. SCHWAB®  
**UMBRELLA SUNSCREEN**  
BROAD SPECTRUM  
SPF 30  
ENVIRONMENTAL PROTECTION  
CRÈME SOLAIRE HYDRATANTE  
HUMECTANTE PROTECCIÓN SOLAR

SUN

2 oz / 56.7 g

www.DoctorSchwabCA.com

DOCTOR D. SCHWAB®  
Manufactured by CA BOTANICA  
San Diego, CA 92101, USA.  
Made in USA

REFRESH, NDC # 35192-007-01

**Drug Facts**

Active Ingredients	Purpose
Titanium Dioxide 4%	Sunscreen
Ocinoxate 4%	Sunscreen

**Uses** • helps prevent sunburn • **Warnings** • **For external use only** • Do not use on damaged or broken skin • **When using this product** keep out of eyes. Rinse with water to remove • **Stop use and ask a doctor** if rash occurs • **Keep out of reach of children.** If product is swallowed, get medical help or contact a Poison Control Center right away.

**Directions** • apply liberally 15 minutes before sun exposure • use a water resistant sunscreen if swimming or sweating • reapply at least every 2 hours • children under 6 months: Ask a doctor • **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including: • limit time in the sun, especially from 10 a.m. – 2 p.m. • wear long-sleeve shirts, pants, hats, and sunglasses

**Inactive ingredients:** Aloe barbadensis leaf extract, Aluminum starch octenylsuccinate, Ascorbic acid, Ascorbyl palmitate, Boron nitride, Camellia sinensis (White Tea) extract, Caprylic/Capric triglyceride, Stearyl alcohol (and) Ceteareth-20, Cetearyl isonanoate, Cetyl alcohol, Citric acid, Citrus aurantium dulcis (Orange) oil, Coco caprylate, Ethylhexylglycerin, Glycerine, Helianthus annuus (Sunflower) seed oil, Niacinamide, Panthenol, Phenylethanol, Phospholipids, Potassium sorbate, Propanediol (Zemea), Rosmarinus officinalis leaf extract CO2, Sodium hyaluronate, Sorbitan monostearate, Tetrasodium EDTA, Tocopherol, Vitis vinifera (Grape) seed extract, Xanthan gum, Water (Aqua).

**Other Information** • protect this product from excessive heat and direct sun.

octinoxate titanium dioxide lotion

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:35192-007
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>OCTINOXATE</b> (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.268 g in 56.7 g
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2.268 g in 56.7 g

### Inactive Ingredients

Ingredient Name	Strength
<b>ALOE VERA LEAF</b> (UNII: ZY81Z83H0X)	
<b>ALUMINUM STARCH OCTENYLSUCCINATE</b> (UNII: I9PJ006294)	
<b>ASCORBIC ACID</b> (UNII: PQ6CK8PD0R)	
<b>ASCORBYL PALMITATE</b> (UNII: QN83US2B0N)	
<b>BORON NITRIDE</b> (UNII: 2U4T60A6YD)	
<b>CAMELLIA SINENSIS FLOWER</b> (UNII: 9I2BJY2J17)	
<b>CAPRYLIC/CAPRIC MONO/DIGLYCERIDES</b> (UNII: U72Q2I8C85)	
<b>STEARYL ALCOHOL</b> (UNII: 2KR89I4H1Y)	
<b>CETYL ALCOHOL</b> (UNII: 936JST6JCN)	
<b>CETEARYL ISONONANOATE</b> (UNII: P5O01U99NI)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>CITRUS AURANTIUM FLOWER OIL</b> (UNII: D4BGE91OXH)	
<b>COCO-CAPRYLATE</b> (UNII: 4828G836N6)	
<b>ETHYLHEXYLGLYCERIN</b> (UNII: 147D247K3P)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>SUNFLOWER OIL</b> (UNII: 3W1JG795YI)	
<b>NIACIN</b> (UNII: 2679MF687A)	
<b>PANTHENOL</b> (UNII: WW9CM0067Z)	
<b>PHENOXYETHANOL</b> (UNII: HIE492ZZ3T)	
<b>EGG PHOSPHOLIPIDS</b> (UNII: 1Z74184RGV)	
<b>POTASSIUM SORBATE</b> (UNII: 1VPU26JZZ4)	
<b>PROPANEDIOL</b> (UNII: 5965N8W85T)	
<b>ROSMARINUS OFFICINALIS FLOWERING TOP OIL</b> (UNII: OXN0D3N28L)	
<b>PEG-9 DIGLYCIDYL ETHER/SODIUM HYALURONATE CROSSPOLYMER</b> (UNII: 788QAG3W8A)	
<b>SORBITAN MONOSTEARATE</b> (UNII: NVZ4I0H58X)	
<b>TOCOPHEROL</b> (UNII: R0ZB2556P8)	
<b>GRAPE SEED OIL</b> (UNII: 930MLC8XGG)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>WATER</b> (UNII: 059QF0KO0R)	

### Packaging

		<b>Marketing Start</b>	<b>Marketing End</b>
--	--	------------------------	----------------------

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:35192-007-01	56.7 g in 1 TUBE; Type 0: Not a Combination Product	08/01/2013	

  

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	08/01/2013	

**Labeler** - CA-BOTANA INTERNATIONAL (106276728)

**Registrant** - RODOLFO UGELSTAD (106276728)

### Establishment

Name	Address	ID/FEI	Business Operations
CA-BOTANA INTERNATIONAL		106276728	manufacture(35192-007)

Revised: 12/2024

CA-BOTANA INTERNATIONAL