

SEIZALAM - midazolam hydrochloride injection, solution

Meridian Medical Technologies LLC

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use SEIZALAM[®] safely and effectively. See full prescribing information for SEIZALAM[®].

SEIZALAM[®] (midazolam injection), for intramuscular use, CIV
Initial U.S. Approval: 1985

WARNING: RISKS FROM CONCOMITANT USE WITH OPIOIDS; ABUSE, MISUSE, AND ADDICTION; and DEPENDENCE AND WITHDRAWAL REACTIONS

See full prescribing information for complete boxed warning.

- Concomitant use of benzodiazepines and opioids may result in profound sedation, respiratory depression, coma, and death. Monitor patients for respiratory depression and sedation. (5.1, 7.1)
- The use of benzodiazepines, including SEIZALAM, exposes users to risks of abuse, misuse, and addiction, which can lead to overdose or death. Before prescribing SEIZALAM and throughout treatment, assess each patient's risk for abuse, misuse, and addiction. (5.2)
- Although SEIZALAM is indicated only for intermittent use (1, 2), if used more frequently than recommended, abrupt discontinuation or rapid dosage reduction of SEIZALAM may precipitate acute withdrawal reactions, which can be life-threatening. For patients using SEIZALAM more frequently than recommended, to reduce the risk of withdrawal reactions, use a gradual taper to discontinue SEIZALAM. (5.3)

----- RECENT MAJOR CHANGES -----

Warnings and Precautions (5.9)

1/2023

----- INDICATIONS AND USAGE -----

SEIZALAM is a benzodiazepine indicated for the treatment of status epilepticus in adults. (1)

----- DOSAGE AND ADMINISTRATION -----

- The recommended dose is 10 mg, administered by intramuscular injection. (2.1)
- Inject in the mid-outer thigh (vastus lateralis muscle). (2.2)
- Continuous monitoring of respiratory and cardiac function is recommended. (2.3)

----- DOSAGE FORMS AND STRENGTHS -----

Injection: 50 mg/10 mL (5 mg/mL) in a multiple-dose vial. (3)

----- CONTRAINDICATIONS -----

Hypersensitivity to midazolam. (4)

----- WARNINGS AND PRECAUTIONS -----

- Risks of Cardiorespiratory Adverse Reactions: Serious cardiorespiratory adverse reactions have occurred, sometimes resulting in death or permanent neurologic injury, after administration of midazolam. (5.4)
- Other Adverse Reactions: Agitation can occur. (5.5)
- Risks from Concomitant Use of Central Nervous System (CNS) Depressants: May increase risk of hypoventilation, airway obstruction, desaturation, or apnea, and may contribute to profound and/or prolonged drug effect. Practitioners administering SEIZALAM must have the skills necessary to manage serious cardiorespiratory adverse reactions, including skills in airway management. (5.6)
- Impaired Cognitive Function: Because of partial or complete impairment of recall, patients should not operate hazardous machinery or a motor vehicle until drug effects have subsided. (5.7)
- Glaucoma: Patients with open-angle glaucoma may need to have their ophthalmologic status evaluated following treatment with SEIZALAM. SEIZALAM is not recommended in patients with narrow-angle glaucoma. (5.8)

- Neonatal Sedation and Withdrawal Syndrome: SEIZALAM use during pregnancy can result in neonatal sedation and/or neonatal withdrawal (5.9, 8.1)

-----ADVERSE REACTIONS-----

The most common adverse reactions (incidence >3%) in clinical trials in patients with status epilepticus were upper airway obstruction, agitation, and pyrexia. (6.1)

The most common adverse reaction (incidence >5%) observed in clinical trials for uses other than that for which SEIZALAM is indicated was decreased tidal volume and/or respiratory rate decrease (11%). (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Meridian Medical Technologies[®], LLC at 1-833-739-0945 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

-----DRUG INTERACTIONS-----

- Benzodiazepines and Opioids: Risk of respiratory depression is increased (7.1)
- Other CNS Depressants and Alcohol: Sedative effect of midazolam is increased. (7.2)
- Cytochrome P450-3A4 Inhibitors: May result in prolonged sedation due to decreased plasma clearance of midazolam. (7.3)

-----USE IN SPECIFIC POPULATIONS-----

Pregnancy: Based on animal data, may cause fetal harm. (8.1)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 1/2023

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: RISKS FROM CONCOMITANT USE WITH OPIOIDS

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dose

2.2 Important Administration Instructions

2.3 Monitoring

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

5.1 Risks from Concomitant Use with Opioids

5.2 Abuse, Misuse, and Addiction

5.3 Dependence and Withdrawal Reactions After Use of SEIZALAM More Frequently Than Recommended

5.4 Risks of Cardiorespiratory Adverse Reactions

5.5 Other Adverse Reactions

5.6 Risks from Concomitant Use of Central Nervous System Depressants

5.7 Impaired Cognitive Function

5.8 Glaucoma

5.9 Neonatal Sedation and Withdrawal Syndrome

5.10 Risk of Serious Adverse Reactions in Infants due to Benzyl Alcohol Preservative

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

7 DRUG INTERACTIONS

7.1 Effect of Concomitant Use of Benzodiazepines and Opioids

7.2 Other CNS Depressants and Alcohol

7.3 Cytochrome P450-3A4 Inhibitors

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment
- 8.7 Congestive Heart Failure

9 DRUG ABUSE AND DEPENDENCE

- 9.1 Controlled Substance
- 9.2 Abuse
- 9.3 Dependence

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

- 16.1 How Supplied
- 16.2 Storage and Handling

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: RISKS FROM CONCOMITANT USE WITH OPIOIDS

- **Concomitant use of benzodiazepines and opioids may result in profound sedation, respiratory depression, coma, and death. Monitor patients for respiratory depression and sedation [see Warnings and Precautions (5.1), Drug Interactions (7.1)].**
- **The use of benzodiazepines, including SEIZALAM, exposes users to risks of abuse, misuse, and addiction, which can lead to overdose or death. Abuse and misuse of benzodiazepines commonly involve concomitant use of other medications, alcohol, and/or illicit substances, which is associated with an increased frequency of serious adverse outcomes. Before prescribing SEIZALAM and throughout treatment, assess each patient's risk for abuse, misuse, and addiction [see Warnings and Precautions (5.2)].**
- **The continued use of benzodiazepines may lead to clinically significant physical dependence. The risks of dependence and withdrawal increase with longer treatment duration and higher daily dose. Although SEIZALAM is indicated only for intermittent use [see Indications and Usage (1) and Dosage and Administration (2)], if used more frequently than recommended, abrupt discontinuation or rapid dosage reduction of SEIZALAM may precipitate acute withdrawal reactions, which can be life-threatening. For patients using SEIZALAM more frequently than recommended, to reduce the risk of withdrawal reactions, use a gradual taper to discontinue SEIZALAM [see Warnings and Precautions (5.3)].**

1 INDICATIONS AND USAGE

SEIZALAM is indicated for the treatment of status epilepticus in adults.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dose

The recommended dose of SEIZALAM is 10 mg, administered by intramuscular injection.

2.2 Important Administration Instructions

SEIZALAM should be administered by a healthcare professional who has had adequate training in the recognition and treatment of status epilepticus.

SEIZALAM is for intramuscular use only. Inject in the mid-outer thigh (vastus lateralis muscle).

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit [see *Dosage Forms and Strengths (3)*].

2.3 Monitoring

After administration of SEIZALAM, continuous monitoring of respiratory and cardiac function is recommended until the patient is stabilized. Serious and life-threatening cardiorespiratory adverse reactions, such as hypoventilation, airway obstruction, apnea, and hypotension have been reported with the use of midazolam. Patients should be monitored in a setting that allows for immediate access to resuscitative drugs. Appropriate resuscitation equipment and personnel trained in their use and skilled in airway management should be available [*see Warnings and Precautions (5.4), Adverse Reactions (6.1)*].

Observation for signs of cardiorespiratory depression is particularly important in patients with chronic obstructive pulmonary disease (COPD), patients 60 or more years of age, and patients who have received concomitant narcotics or other central nervous system (CNS) depressants.

3 DOSAGE FORMS AND STRENGTHS

Injection: 50 mg/10 mL (5 mg/mL) of a sterile, clear, colorless to light yellow liquid solution in a multiple-dose vial.

4 CONTRAINDICATIONS

SEIZALAM is contraindicated in patients with a known hypersensitivity to midazolam.

5 WARNINGS AND PRECAUTIONS

5.1 Risks from Concomitant Use with Opioids

Concomitant use of benzodiazepines, including SEIZALAM, and opioids may result in profound sedation, respiratory depression, coma, and death. If a decision is made to use midazolam concomitantly with opioids, monitor patients closely for respiratory depression and sedation [*see Drug Interactions (7.1)*].

Practitioners administering SEIZALAM must have the skills necessary to manage serious cardiorespiratory adverse reactions, including skills in airway management.

5.2 Abuse, Misuse, and Addiction

The use of benzodiazepines, including SEIZALAM, exposes users to the risks of abuse, misuse, and addiction, which can lead to overdose or death. Abuse and misuse of benzodiazepines often (but not always) involve the use of doses greater than the maximum recommended dosage and commonly involve concomitant use of other medications, alcohol, and/or illicit substances, which is associated with an increased frequency of serious adverse outcomes, including respiratory depression, overdose, or death [*see Drug Abuse and Dependence (9.2)*].

Before prescribing SEIZALAM and throughout treatment, assess each patient's risk for abuse, misuse, and addiction. Use of SEIZALAM, particularly in patients at elevated risk, necessitates counseling about the risks and proper use of SEIZALAM along with monitoring for signs and symptoms of abuse, misuse, and addiction. Do not exceed the recommended dosing frequency; avoid or minimize concomitant use of CNS depressants and other substances associated with abuse, misuse, and addiction (e.g.,

opioid analgesics, stimulants); and advise patients on the proper disposal of unused drug. If a substance use disorder is suspected, evaluate the patient and institute (or refer them for) early treatment, as appropriate.

5.3 Dependence and Withdrawal Reactions After Use of SEIZALAM More Frequently Than Recommended

For patients using SEIZALAM more frequently than recommended, to reduce the risk of withdrawal reactions, use a gradual taper to discontinue SEIZALAM (a patient-specific plan should be used to taper the dose).

Patients at an increased risk of withdrawal adverse reactions after benzodiazepine discontinuation or rapid dosage reduction include those who take higher dosages, and those who have had longer durations of use.

Acute Withdrawal Reactions

The continued use of benzodiazepines may lead to clinically significant physical dependence. Although SEIZALAM is indicated only for intermittent use [*see Indications and Usage (1) and Dosage and Administration (2)*], if used more frequently than recommended, abrupt discontinuation or rapid dosage reduction of SEIZALAM, or administration of flumazenil (a benzodiazepine antagonist) may precipitate acute withdrawal reactions, which can be life-threatening (e.g., seizures) [*see Drug Abuse and Dependence (9.3)*].

Protracted Withdrawal Syndrome

In some cases, benzodiazepine users have developed a protracted withdrawal syndrome with withdrawal symptoms lasting weeks to more than 12 months [*see Drug Abuse and Dependence (9.3)*].

5.4 Risks of Cardiorespiratory Adverse Reactions

Serious cardiorespiratory adverse reactions have occurred after administration of midazolam. These have included respiratory depression, airway obstruction, oxygen desaturation, apnea, respiratory arrest and/or cardiac arrest, sometimes resulting in death or permanent neurologic injury. There have also been rare reports of hypotensive episodes requiring treatment during or after diagnostic or surgical manipulations, particularly in patients with hemodynamic instability. Hypotension occurs more frequently in patients premedicated with a narcotic. The danger of hypoventilation, airway obstruction, or apnea is greater in elderly patients and those with chronic disease states or decreased pulmonary reserve [*see Use in Specific Populations (8.5, 8.7)*]; patients with COPD are highly sensitive to the respiratory depressant effect of midazolam. SEIZALAM should be administered with caution to patients in shock or coma with depression of vital signs.

Practitioners administering SEIZALAM must have the skills necessary to manage serious cardiorespiratory adverse reactions, including skills in airway management.

5.5 Other Adverse Reactions

Reactions such as agitation, involuntary movements (including tonic/clonic movements and muscle tremor), hyperactivity, and combativeness have been reported with midazolam when used for sedation. These reactions may be caused by inadequate or

excessive dosing or improper administration of midazolam; however, consideration should be given to the possibility of cerebral hypoxia or true paradoxical reactions. Agitation also occurred in the randomized controlled clinical study of SEIZALAM in patients with status epilepticus [see *Adverse Reactions (6.1)*].

5.6 Risks from Concomitant Use of Central Nervous System Depressants

Concomitant use of barbiturates, alcohol or other central nervous system depressants may increase the risk of hypoventilation, airway obstruction, desaturation, or apnea and may contribute to profound and/or prolonged drug effect. SEIZALAM should be administered with caution to patients in acute alcohol intoxication with depression of vital signs. Narcotic premedication also depresses the ventilatory response to carbon dioxide stimulation.

The efficacy and safety of midazolam in clinical use are functions of the dose administered, the clinical status of the individual patient, and the use of concomitant medications capable of depressing the central nervous system (CNS). Anticipated effects range from mild sedation to deep levels of sedation virtually equivalent to a state of general anesthesia where the patient may require external support of vital functions. Practitioners administering SEIZALAM must have the skills necessary to manage serious cardiorespiratory adverse reactions, including skills in airway management. For information regarding withdrawal, see *Drug Abuse and Dependence (9.3)*.

5.7 Impaired Cognitive Function

Midazolam is associated with a high incidence of partial or complete impairment of recall for several hours following an administered dose. Gross tests of recovery from the effects of midazolam cannot be relied upon to predict reaction time under stress. It is recommended that no patient operate hazardous machinery or a motor vehicle until the effects of the drug, such as drowsiness, have subsided, and as their medical condition permits.

5.8 Glaucoma

Benzodiazepines, including SEIZALAM, can increase intraocular pressure in patients with glaucoma. Measurements of intraocular pressure in patients without eye disease show a moderate lowering following induction with midazolam; patients with glaucoma have not been studied. Patients with open-angle glaucoma may need to have their ophthalmologic status evaluated following treatment with SEIZALAM. SEIZALAM is not recommended in patients with narrow-angle glaucoma.

5.9 Neonatal Sedation and Withdrawal Syndrome

Use of SEIZALAM late in pregnancy can result in sedation (respiratory depression, lethargy, hypotonia) and/or withdrawal symptoms (hyperreflexia, irritability, restlessness, tremors, inconsolable crying, and feeding difficulties) in the neonate [see *Use in Specific Populations (8.1)*]. Monitor neonates exposed to SEIZALAM during pregnancy or labor for signs of sedation and monitor neonates exposed to SEIZALAM during pregnancy for signs of withdrawal; manage these neonates accordingly.

5.10 Risk of Serious Adverse Reactions in Infants due to Benzyl Alcohol Preservative

SEIZALAM is not approved for use in neonates or infants. Serious and fatal adverse reactions including “gaspings syndrome” can occur in neonates and low birth weight infants treated with benzyl alcohol-preserved drugs, including SEIZALAM. The “gaspings syndrome” is characterized by CNS depression, metabolic acidosis, and gasping respirations. The minimum amount of benzyl alcohol at which serious adverse reactions may occur is not known (SEIZALAM contains 10 mg of benzyl alcohol per mL) [see Use in Specific Populations (8.4)].

6 ADVERSE REACTIONS

The following serious adverse reactions are discussed in greater detail in other sections:

- Risks from Concomitant Use with Opioids [see Warnings and Precautions (5.1)]
- Abuse, Misuse, and Addiction [see Warnings and Precautions (5.2)]
- Dependence and Withdrawal Reactions After Use of SEIZALAM More Frequently Than Recommended [see Warnings and Precautions (5.3)]
- Risks of Cardiorespiratory Adverse Reactions [see Warnings and Precautions (5.4)]
- Other Adverse Reactions [see Warnings and Precautions (5.5)]
- Risks from Concomitant Use of Central Nervous System Depressants [see Warnings and Precautions (5.6)]
- Impaired Cognitive Function [see Warnings and Precautions (5.7)]
- Glaucoma [see Warnings and Precautions (5.8)]
- Neonatal Sedation and Withdrawal Syndrome [see Warnings and Precautions (5.9)]
- Risk of Serious Adverse Reactions in Infants due to Benzyl Alcohol Preservative [see Warnings and Precautions (5.10)]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Adverse Reactions in the Controlled Study of Intramuscular Midazolam in Patients with Status Epilepticus

In a double-blind, randomized, active-controlled clinical study, 448 patients were assigned to receive intramuscular (IM) midazolam via an autoinjector, and 445 were assigned to receive intravenous (IV) lorazepam. Approximately 45% of patients were female, and the mean age was 43 years. Patients were administered treatment by a healthcare professional (e.g., paramedic) prior to arrival at a hospital.

Table 1 lists the adverse reactions occurring in 2% or more of the IM midazolam-treated patients and at a rate greater than the IV lorazepam-treated patients.

Table 1 Adverse Reactions in 2% or More of IM Midazolam-Treated Patients and More Frequent than in IV Lorazepam-Treated Patients in Out of Hospital Treatment of Status Epilepticus

| Adverse Reaction | IM Midazolam N=448 (%) | IV Lorazepam N=445 (%) |
|--------------------------|---------------------------------------|---------------------------------------|
| Upper airway obstruction | 5 | 3 |

| | | |
|-----------------------|---|---|
| Agitation | 4 | 3 |
| Pyrexia | 4 | 3 |
| Mental status changes | 3 | 2 |
| Postictal state | 3 | 2 |
| Acute renal failure | 2 | 1 |

Adverse Reactions in Other Midazolam Studies

Fluctuations in vital signs were the most frequently seen findings following parenteral administration of midazolam in adults for uses other than that for which SEIZALAM is indicated, and included decreased tidal volume and/or respiratory rate decrease [11% of patients following intramuscular administration], as well as variations in blood pressure and pulse rate. The majority of serious adverse effects, particularly those associated with oxygenation and ventilation, have been reported when midazolam was administered with other medications capable of depressing the CNS. The incidence of such events was higher in patients undergoing procedures involving the airway without the protective effect of an endotracheal tube (e.g., upper endoscopy and dental procedures).

The following additional adverse reactions were reported after intramuscular administration in adults: Headache (1.3%), and local effects at the IM injection site including pain (3.7%), induration (0.5%), redness (0.5%), and muscle stiffness (0.3%).

7 DRUG INTERACTIONS

7.1 Effect of Concomitant Use of Benzodiazepines and Opioids

The concomitant use of benzodiazepines and opioids increases the risk of respiratory depression because of actions at different receptor sites in the CNS that control respiration. Benzodiazepines interact at GABA_A sites and opioids interact primarily at mu receptors. When benzodiazepines and opioids are combined, the potential for benzodiazepines to significantly worsen opioid-related respiratory depression exists. Limit dosage and duration of concomitant use of benzodiazepines and opioids. Monitor patients closely for respiratory depression and sedation.

7.2 Other CNS Depressants and Alcohol

The sedative effect of SEIZALAM is accentuated by concomitantly administered medication that depresses the central nervous system, particularly opioids (e.g., morphine, meperidine, and fentanyl), secobarbital, and droperidol, and also by alcohol [*see Warnings and Precautions (5.1, 5.4, 5.6)*].

7.3 Cytochrome P450-3A4 Inhibitors

Caution is advised when SEIZALAM is administered concomitantly with drugs that are known to inhibit the P450-3A4 enzyme system (e.g., cimetidine, erythromycin, diltiazem, verapamil, ketoconazole, and itraconazole). These drug interactions may result in prolonged sedation caused by a decrease in plasma clearance of midazolam [*see Clinical Pharmacology (12.3)*].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Exposure Registry

There is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to antiepileptic drugs (AEDs), such as SEIZALAM, during pregnancy. Healthcare providers are encouraged to recommend that pregnant women who are taking SEIZALAM during pregnancy enroll in the North American Antiepileptic Drug (NAAED) pregnancy registry by calling 1-888-233-2334 or visiting <http://www.aedpregnancyregistry.org/>.

Risk Summary

Neonates born to mothers using benzodiazepines late in pregnancy have been reported to experience symptoms of sedation and/or neonatal withdrawal [*see Warnings and Precautions (5.9) and Clinical Considerations*]. Available data from published observational studies of pregnant women exposed to benzodiazepines do not report a clear association with benzodiazepines and major birth defects (*see Data*).

Administration of midazolam to rats and rabbits during the period of organogenesis or to rats during late pregnancy and throughout lactation at doses greater than those used clinically did not result in adverse effects on development (*see Animal Data*). Data for other benzodiazepines suggest the possibility of increased neuronal cell death and long-term effects on neurobehavioral and immunological function based on findings in animals following prenatal or early postnatal exposure at clinically relevant doses.

The background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defects, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

Clinical Considerations

Fetal/Neonatal Adverse Reactions

Benzodiazepines cross the placenta and may produce respiratory depression, hypotonia, and sedation in neonates. Monitor neonates exposed to SEIZALAM during pregnancy or labor for signs of sedation, respiratory depression, hypotonia, and feeding problems. Monitor neonates exposed to SEIZALAM during pregnancy for signs of withdrawal. Manage these neonates accordingly [*see Warnings and Precautions (5.9)*].

Data

Human Data

Published data from observational studies on the use of benzodiazepines during pregnancy do not report a clear association with benzodiazepines and major birth defects.

Although early studies reported an increased risk of congenital malformations with diazepam and chlordiazepoxide, there was no consistent pattern noted. In addition, the majority of more recent case-control and cohort studies of benzodiazepine use during pregnancy, which were adjusted for confounding exposures to alcohol, tobacco and

other medications, have not confirmed these findings.

Animal Data

When midazolam (0, 0.2, 1, or 4 mg/kg/day) was administered intravenously to pregnant rats during the period of organogenesis, no adverse effects on embryofetal development were observed. The highest dose tested, which was associated with minimal evidence of maternal toxicity, is approximately 4 times the recommended human dose (RHD) of 10 mg based on body surface area (mg/m²).

When midazolam (0, 0.2, 0.6, and 2 mg/kg/day) was administered intravenously to rabbits during the period of organogenesis, no adverse effects on embryofetal development were reported. The high dose, which was not associated with evidence of maternal toxicity, is approximately 4 times the RHD on a mg/m² basis.

When midazolam (0, 0.2, 1, or 4 mg/kg/day) was administered intravenously to female rats during late gestation and throughout lactation, no clear adverse effects were noted in the offspring. The high dose, which was not associated with evidence of maternal toxicity, is approximately 4 times the RHD on a mg/m² basis.

In published animal studies, administration of benzodiazepines or other drugs that enhance GABAergic inhibition to neonatal rats has been reported to result in widespread apoptotic neurodegeneration in the developing brain at plasma concentrations relevant for seizure control in humans. The window of vulnerability to these changes in rats (postnatal days 0-14) includes a period of brain development that takes place during the third trimester of pregnancy in humans.

8.2 Lactation

Risk Summary

Midazolam is excreted in human milk. There are reports of sedation, poor feeding and poor weight gain in infants exposed to benzodiazepines through breast milk. There are no data on the effects of midazolam on milk production.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for SEIZALAM and any potential adverse effects on the breastfed child from SEIZALAM or from the underlying maternal condition.

Clinical Considerations

Infants exposed to SEIZALAM through breast milk should be monitored for sedation, poor feeding and poor weight gain.

8.4 Pediatric Use

Safety and effectiveness in pediatric patients have not been established. Benzodiazepines are not recognized as a treatment for status epilepticus in neonates and should not be used in this population.

SEIZALAM is not approved for use in neonates or infants. Serious adverse reactions including fatal reactions and the "gasping syndrome" occurred in premature neonates and low-birth-weight infants in the neonatal intensive care unit who received drugs containing benzyl alcohol as a preservative. In these cases, benzyl alcohol dosages of 99 to 234 mg/kg/day produced high levels of benzyl alcohol and its metabolites in the blood and urine (blood levels of benzyl alcohol were 0.61 to 1.378 mmol/L). Additional adverse

reactions included gradual neurological deterioration, seizures, intracranial hemorrhage, hematologic abnormalities, skin breakdown, hepatic and renal failure, hypotension, bradycardia, and cardiovascular collapse. Preterm, low-birth-weight infants may be more likely to develop these reactions because they may be less able to metabolize benzyl alcohol. The minimum amount of benzyl alcohol at which serious adverse reactions may occur is not known (SEIZALAM contains 10 mg of benzyl alcohol per mL) [see *Warnings and Precautions* (5.10)].

8.5 Geriatric Use

Of the total number of patients from the intent-to-treat (ITT) population in the clinical trial of SEIZALAM, 14.9 percent were 65 years of age and over, while 8.3 percent were 75 years of age and over.

Geriatric patients may have altered drug distribution; diminished hepatic and/or renal function; longer elimination half-lives for midazolam and its metabolites, and subjects over 70 years of age may be particularly sensitive [see *Clinical Pharmacology* (12.3)]. Administration of IM midazolam to elderly patients has been associated with rare reports of death under circumstances compatible with cardiorespiratory depression [see *Warnings and Precautions* (5.4)]. In most of these cases, the patients also received other CNS depressants capable of depressing respiration, especially narcotics [see *Warnings and Precautions* (5.1, 5.6)]. Close monitoring of geriatric patients is recommended.

8.6 Renal Impairment

Patients with renal impairment may have a slower elimination of midazolam and its metabolites, which may result in prolonged drug exposure [see *Clinical Pharmacology* (12.3)].

8.7 Congestive Heart Failure

Patients with congestive heart failure eliminate midazolam more slowly, which may result in prolonged drug exposure [see *Clinical Pharmacology* (12.3)].

9 DRUG ABUSE AND DEPENDENCE

9.1 Controlled Substance

SEIZALAM contains midazolam hydrochloride, a Schedule IV controlled substance.

9.2 Abuse

SEIZALAM is a benzodiazepine and a CNS depressant with a potential for abuse and addiction. Abuse is the intentional, non-therapeutic use of a drug, even once, for its desirable psychological or physiological effects. Misuse is the intentional use, for therapeutic purposes, of a drug by an individual in a way other than prescribed by a health care provider or for whom it was not prescribed. Drug addiction is a cluster of behavioral, cognitive, and physiological phenomena that may include a strong desire to take the drug, difficulties in controlling drug use (e.g., continuing drug use despite harmful consequences, giving a higher priority to drug use than other activities and obligations), and possible tolerance or physical dependence. Even taking

benzodiazepines as prescribed may put patients at risk for abuse and misuse of their medication. Abuse and misuse of benzodiazepines may lead to addiction.

Abuse and misuse of benzodiazepines often (but not always) involve the use of doses greater than the maximum recommended dosage and commonly involve concomitant use of other medications, alcohol, and/or illicit substances, which is associated with an increased frequency of serious adverse outcomes, including respiratory depression, overdose, or death. Benzodiazepines are often sought by individuals who abuse drugs and other substances, and by individuals with addictive disorders [*see Warnings and Precautions (5.2)*].

The following adverse reactions have occurred with benzodiazepine abuse and/or misuse: abdominal pain, amnesia, anorexia, anxiety, aggression, ataxia, blurred vision, confusion, depression, disinhibition, disorientation, dizziness, euphoria, impaired concentration and memory, indigestion, irritability, muscle pain, slurred speech, tremors, and vertigo.

The following severe adverse reactions have occurred with benzodiazepine abuse and/or misuse: delirium, paranoia, suicidal ideation and behavior, seizures, coma, breathing difficulty, and death. Death is more often associated with polysubstance use (especially benzodiazepines with other CNS depressants such as opioids and alcohol).

Midazolam was actively self-administered in primate models used to assess the positive reinforcing effects of psychoactive drugs. Midazolam produced physical dependence of a mild to moderate intensity in cynomolgus monkeys after 5 to 10 weeks of administration.

Available data concerning the drug abuse and dependence potential of midazolam suggest that its abuse potential is at least equivalent to that of diazepam.

9.3 Dependence

Physical Dependence After Use of SEIZALAM More Frequently Than Recommended

SEIZALAM may produce physical dependence if used more frequently than recommended. Physical dependence is a state that develops as a result of physiological adaptation in response to repeated drug use, manifested by withdrawal signs and symptoms after abrupt discontinuation or a significant dose reduction of a drug. Although SEIZALAM is indicated only for intermittent use [*see Indications and Usage (1) and Dosage and Administration (2)*], if used more frequently than recommended, abrupt discontinuation or rapid dosage reduction or administration of flumazenil, a benzodiazepine antagonist, may precipitate acute withdrawal reactions, including seizures, which can be life-threatening. Patients at an increased risk of withdrawal adverse reactions after benzodiazepine discontinuation or rapid dosage reduction include those who take higher dosages (i.e., higher and/or more frequent doses) and those who have had longer durations of use [*see Warnings and Precautions (5.3)*].

For patients using SEIZALAM more frequently than recommended, to reduce the risk of withdrawal reactions, use a gradual taper to discontinue SEIZALAM [*see Warnings and Precautions (5.3)*].

Acute Withdrawal Signs and Symptoms

Acute withdrawal signs and symptoms associated with benzodiazepines have included abnormal involuntary movements, anxiety, blurred vision, depersonalization, depression,

derealization, dizziness, fatigue, gastrointestinal adverse reactions (e.g., nausea, vomiting, diarrhea, weight loss, decreased appetite), headache, hyperacusis, hypertension, irritability, insomnia, memory impairment, muscle pain and stiffness, panic attacks, photophobia, restlessness, tachycardia, and tremor. More severe acute withdrawal signs and symptoms, including life-threatening reactions, have included catatonia, convulsions, delirium tremens, depression, hallucinations, mania, psychosis, seizures and suicidality.

Protracted Withdrawal Syndrome

Protracted withdrawal syndrome associated with benzodiazepines is characterized by anxiety, cognitive impairment, depression, insomnia, formication, motor symptoms (e.g., weakness, tremor, muscle twitches), paresthesia, and tinnitus that persists beyond 4 to 6 weeks after initial benzodiazepine withdrawal. Protracted withdrawal symptoms may last weeks to more than 12 months. As a result, there may be difficulty in differentiating withdrawal symptoms from potential re-emergence or continuation of symptoms for which the benzodiazepine was being used.

Tolerance

Tolerance to SEIZALAM may develop after use more frequently than recommended. Tolerance is a physiological state characterized by a reduced response to a drug after repeated administration (i.e., a higher dose of a drug is required to produce the same effect that was once obtained at a lower dose). Tolerance to the therapeutic effect of benzodiazepines may develop; however, little tolerance develops to the amnestic reactions and other cognitive impairments caused by benzodiazepines.

10 OVERDOSAGE

Overdosage of benzodiazepines is characterized by central nervous system depression ranging from drowsiness to coma. In mild to moderate cases, symptoms can include drowsiness, confusion, dysarthria, lethargy, hypnotic state, diminished reflexes, ataxia, and hypotonia. Rarely, paradoxical or disinhibitory reactions (including agitation, irritability, impulsivity, violent behavior, confusion, restlessness, excitement, and talkativeness) may occur. In severe overdosage cases, patients may develop respiratory depression and coma. Overdosage of benzodiazepines in combination with other CNS depressants (including alcohol and opioids) may be fatal [see *Warnings and Precautions (5.2)*]. Markedly abnormal (lowered or elevated) blood pressure, heart rate, or respiratory rate raise the concern that additional drugs and/or alcohol are involved in the overdosage.


In managing benzodiazepine overdosage, employ general supportive measures, including intravenous fluids and airway maintenance. Flumazenil, a specific benzodiazepine receptor antagonist indicated for the complete or partial reversal of the sedative effects of benzodiazepines in the management of benzodiazepine overdosage, can lead to withdrawal and adverse reactions, including seizures, particularly in the context of mixed overdosage with drugs that increase seizure risk (e.g., tricyclic and tetracyclic antidepressants) and in patients with long-term benzodiazepine use and physical dependency. The risk of withdrawal seizures with flumazenil use may be increased in patients with epilepsy. Flumazenil is contraindicated in patients who have received a benzodiazepine for control of a potentially life-threatening condition (e.g., status epilepticus). If the decision is made to use flumazenil, it should be used as an

adjunct to, not as a substitute for, supportive management of benzodiazepine overdose. See the flumazenil injection Prescribing Information.

Consider contacting the Poison Help line (1-800-222-1222) or a medical toxicologist for additional overdose management recommendations.

11 DESCRIPTION

Midazolam is a white to light yellow crystalline compound, insoluble in water. The hydrochloride salt of midazolam, which is formed *in situ*, is soluble in aqueous solutions. Chemically, midazolam HCl is 8-chloro-6-(2-fluorophenyl)-1-methyl-4*H*-imidazo[1,5-*a*][1,4]benzodiazepine hydrochloride. Midazolam hydrochloride has the empirical formula $C_{18}H_{13}ClFN_3 \cdot HCl$, a calculated molecular weight of 362.24 and the following structural formula:



SEIZALAM is a sterile, nonpyrogenic solution for intramuscular injection. Each mL contains 5 mg midazolam (equivalent to 5.6 mg midazolam hydrochloride) compounded with 1% benzyl alcohol as preservative, 0.01% edetate disodium, and 0.8% sodium chloride. The pH is adjusted to approximately 3 with hydrochloric acid and, if necessary, sodium hydroxide.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The exact mechanism of action for midazolam in the treatment of status epilepticus is not fully understood, but is thought to involve potentiation of GABAergic neurotransmission resulting from binding at the benzodiazepine site of the GABA_A receptor.

12.2 Pharmacodynamics

The effects of midazolam on the CNS are dependent on the dose administered, the route of administration, and the presence or absence of other medications.

12.3 Pharmacokinetics

The pharmacokinetics of midazolam were evaluated in a single dose-escalation trial in

healthy subjects; following IM injection of midazolam at doses ranging from fixed total amounts of 5 mg to 30 mg (one half to three times the recommended dose) or body weight based dose of 0.10 mg/kg to 0.49 mg/kg, the overall median time to maximum plasma concentration (T_{max}) was observed at approximately 0.5 hours after administration. The rate and extent of systemic exposure, as assessed by maximum plasma concentration (C_{max}) and area under the plasma drug concentration-time curve from time 0 to infinity ($AUC_{0-\infty}$), tended to increase proportionally with increasing dose from 5 mg to 30 mg.

Absorption

Following IM administration of a single 10 mg midazolam dose to healthy subjects, midazolam was absorbed with median T_{max} (range) of 0.5 (0.25 to 0.5) hours; midazolam mean (\pm SD) C_{max} and $AUC_{0-\infty}$ were 113.9 (\pm 30.9) ng/mL and 402.7 (\pm 97.0) ng•hr/mL, respectively.

Distribution

The mean (\pm SD) apparent volume of distribution (V_z/F) of midazolam following a single IM dose of 10 mg midazolam was 2117 (\pm 845.1) mL/kg in healthy subjects.

In humans, midazolam has been shown to cross the placenta and enter into fetal circulation, and has been detected in human milk and CSF [see *Use in Specific Populations* (8.1, 8.2)].

In adults, midazolam is approximately 97% bound to plasma protein, principally albumin. The 1-hydroxy metabolite is approximately 89% bound to plasma protein.

Elimination

Elimination of the parent drug takes place via hepatic metabolism of midazolam to hydroxylated metabolites that are conjugated and excreted in the urine.

Metabolism

In vitro studies with human liver microsomes indicate that the biotransformation of midazolam is mediated by the cytochrome P450-3A4 (CYP3A4). This enzyme is present in gastrointestinal tract mucosa, as well as in the liver. The 1-hydroxy-midazolam (also termed alpha-hydroxymidazolam) metabolite comprises 60% to 70% of the biotransformation products of midazolam, while 4-hydroxy-midazolam constitutes 5% or less. Small amounts of a dihydroxy derivative have also been detected, but not quantified. The principal urinary excretion products are glucuronide conjugates of the hydroxylated derivatives.

Studies of the intravenous administration of 1-hydroxy-midazolam in humans suggest that 1-hydroxy-midazolam is at least as potent as the parent compound, and may contribute to the net pharmacologic activity of midazolam. *In vitro* studies have demonstrated that the affinities of 1- and 4-hydroxy-midazolam for the benzodiazepine receptor are approximately 20% and 7%, respectively, relative to midazolam.

Excretion

Following IM administration of 10 mg midazolam, mean (\pm SD) elimination half-life and apparent total body clearance (CL/F) of midazolam were 4.2 (\pm 1.87) hours and 367.3 (\pm 73.5) mL/hr/kg, respectively.

The principal urinary excretion product is 1-hydroxy-midazolam in the form of a

glucuronide conjugate; smaller amounts of the glucuronide conjugates of 4-hydroxy- and dihydroxy-midazolam are detected as well. The amount of midazolam excreted unchanged in the urine after a single IV dose is less than 0.5%. Following a single IV infusion in 5 healthy volunteers, 45% to 57% of the dose was excreted in the urine as the 1-hydroxymethyl midazolam conjugate.

Specific Populations

Changes in the pharmacokinetic profile of midazolam due to drug interactions, physiological variables, etc., may result in changes in the plasma concentration-time profile and pharmacological response to midazolam in these patients. For example, patients with acute renal failure (ARF) appear to have a longer elimination half-life for midazolam [see *Use in Specific Populations (8.6)*]. In other groups, the relationship between prolonged half-life and duration of effect has not been established.

Obesity

In a study comparing normal (n=20) and obese patients (n=20), the mean half-life was greater in the obese group (5.9 versus 2.3 hours). This was due to an increase of approximately 50% in the volume of distribution (V_d) corrected for total body weight. The clearance was not significantly different between groups.

Geriatric Patients

In three parallel-group studies, the pharmacokinetics of midazolam administered IV or IM were compared in young (mean age 29 years, n=52) and healthy elderly subjects (mean age 73 years, n=53). Plasma half-life was approximately 2-fold higher in the elderly. The mean V_d based on total body weight increased consistently between 15% and 100% in the elderly. The mean CL (total clearance) decreased approximately 25% in the elderly in two studies, and was similar to that of the younger patients in the other [see *Use in Specific Populations (8.5)*].

Male and Female Patients

No meaningful differences in midazolam exposures (C_{max} and AUC) were observed between male and female adults after IM administration.

Patients with Congestive Heart Failure

In patients suffering from congestive heart failure, a 2-fold increase in the elimination half-life, a 25% decrease in the plasma clearance, and a 40% increase in the volume of distribution of midazolam were observed.

Patients with Renal Impairment

Patients with renal impairment may have longer elimination half-lives for midazolam and its metabolites [see *Use in Specific Populations (8.6)*].

Midazolam and 1-hydroxy-midazolam pharmacokinetics were compared between 6 intensive care unit (ICU) patients who developed acute renal failure (ARF) and a control group of subjects with normal renal function. Midazolam was administered as an IV infusion (5 to 15 mg/hour). Midazolam clearance was reduced (1.9 versus 2.8 mL/min/kg), and the half-life was prolonged (7.6 hours versus 13 hours) in the ARF patients. The renal clearance of the 1-hydroxy-midazolam glucuronide was prolonged in the ARF group (4 versus 136 mL/min), and the half-life was prolonged (12 hours versus >25 hours). Plasma levels accumulated in all ARF patients to about ten times that of the

parent drug. The relationship between accumulating metabolite levels and prolonged sedation is unclear.

In a study of chronic renal failure patients (n=15) receiving a single IV dose of midazolam, there was a 2-fold increase in the clearance and volume of distribution, but the half-life remained unchanged. Metabolite levels were not studied.

Patients with Hepatic Impairment

Midazolam pharmacokinetics were studied after a single IV dose (0.075 mg/kg) was administered to 7 patients with biopsy-proven alcoholic cirrhosis and 8 control patients. The mean half-life of midazolam increased 2.5-fold in the patients with cirrhosis. Clearance was reduced by 50% and V_d increased by 20%. In another study in 21 male patients with cirrhosis, without ascites and with normal kidney function as determined by creatinine clearance, no changes in the pharmacokinetics of midazolam or 1-hydroxy-midazolam were observed when compared to healthy individuals. The clinical significance of these findings is unknown.

Drug Interaction Studies

CYP3A4 Inhibitors

Drugs that inhibit the activity of CYP3A4 may inhibit midazolam clearance and elevate midazolam concentrations [see *Drug Interactions (7.3)*].

- The effect of single oral doses of 800 mg cimetidine and 300 mg ranitidine on steady-state concentrations of oral midazolam was examined in a randomized crossover study (n=8). Cimetidine increased the mean midazolam steady-state concentration from 57 to 71 ng/mL. Ranitidine increased the mean steady-state concentration to 62 ng/mL. No change in choice reaction time or sedation index was detected after dosing with the H₂ receptor antagonists.
- In a placebo-controlled study, erythromycin administered as a 500 mg dose, three times a day, for 1 week (n=6), reduced the clearance of midazolam following a single 0.5 mg/kg IV dose. The half-life was approximately doubled.
- The effects of diltiazem (60 mg three times a day) and verapamil (80 mg three times a day) on the pharmacokinetics and pharmacodynamics of midazolam were investigated in a three-way crossover study (n=9). The half-life of midazolam increased from 5 to 7 hours when midazolam was taken in conjunction with verapamil or diltiazem. No interaction was observed in healthy subjects between midazolam and nifedipine.
- In a placebo-controlled study, where saquinavir or placebo was administered orally as a 1200 mg dose three times a day for 5 days (n=12), a 56% reduction in the clearance of midazolam following a single 0.05 mg/kg IV dose was observed. The half-life was approximately doubled.

CYP3A4 Inducers

Drugs that induce the activity of CYP3A4 may increase midazolam clearance and decrease midazolam concentrations.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Midazolam maleate was administered in the diet to mice and rats for 2 years at doses of 0, 1, 9, or 80 mg/kg/day. In female mice in the highest dose group there was a marked increase in the incidence of hepatic tumors. In high-dose male rats there was a small but statistically significant increase in benign thyroid follicular cell tumors. The highest dose not associated with increased tumor incidences in mice and rats (9 mg/kg/day) is approximately 4 and 9 times, respectively, the recommended human dose (RHD) of 10 mg based on body surface area (mg/m²). The pathogenesis of induction of these tumors is not known. These tumors were found after chronic administration, whereas human use will ordinarily be of single or several doses.

Mutagenesis

Midazolam was negative for genotoxicity in *in vitro* (Ames, mammalian cell clastogenicity) and *in vivo* (mouse bone marrow micronucleus) assays.

Impairment of Fertility

When midazolam (0, 1, 4, or 16 mg/kg) was orally administered to male and female rats prior to and during mating and continuing in females throughout gestation and lactation, no adverse effects on male or female fertility were noted. Midazolam plasma exposures (AUC) at the highest dose tested were approximately 6 times that in humans at the RHD.

14 CLINICAL STUDIES

The safety and effectiveness of SEIZALAM for the treatment of status epilepticus was established in a multi-center, randomized, double-blind (double-dummy), active-control trial comparing midazolam administered intramuscularly (IM) via an auto-injector to lorazepam administered intravenously (IV). Patients meeting the diagnosis of status epilepticus, with continuing convulsive seizure activity after the arrival of paramedics, were eligible for enrollment. The ITT population consisted of 893 patients who were randomized to receive either IM midazolam (n=448) or IV lorazepam (n=445). Following randomization, each patient received study treatments administered by a healthcare professional (e.g., paramedic) prior to arrival at a hospital. According to the double-dummy design, adult patients received 10 mg IM midazolam followed by IV placebo or received IM placebo followed by 4 mg IV lorazepam. The primary efficacy endpoint was the termination of convulsive seizure activity (without the need for rescue medication) prior to arrival at the emergency department (ED) as determined by the ED attending physician. A statistically significantly higher percentage of midazolam-treated patients met the primary efficacy endpoint, as shown in Table 2.

Table 2: Primary Efficacy Analysis Results: Seizure Termination (without rescue medication)

| | IM Midazolam (n=448) | IV Lorazepam (n= 445) |
|-----------------------|---------------------------------|----------------------------------|
| Treatment success (%) | 73.4 | 63.4 |
| p-value ^a | 0.002 | |

^a Fischer's exact test

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

SEIZALAM (midazolam injection) is a clear, colorless to light yellow sterile solution available in multiple-dose flip-top vials containing 50 mg/10 mL (5 mg/mL).

SEIZALAM is supplied in the following packaging configurations:

- One vial: NDC 11704-650-01
- Carton of 10 vials: NDC 11704-650-10

16.2 Storage and Handling

Store at 20 °C to 25 °C (68 °F to 77 °F); excursions permitted between 15 °C and 30 °C (59 °F and 86 °F) [See USP Controlled Room Temperature].

17 PATIENT COUNSELING INFORMATION

Patients having seizures will likely be unresponsive or may have difficulty in comprehending counseling information.

Risks from Concomitant Use with Opioids

Inform patients and caregivers that potentially fatal additive effects may occur if SEIZALAM is used with opioids and not to use such drugs concomitantly unless supervised by a healthcare provider [see *Warnings and Precautions (5.1), Drug Interactions (7.1)*].

Abuse, Misuse, and Addiction

Inform patients that the use of SEIZALAM more frequently than recommended, even at recommended dosages, exposes users to risks of abuse, misuse, and addiction, which can lead to overdose and death, especially when used in combination with other medications (e.g., opioid analgesics), alcohol, and/or illicit substances. Inform patients about the signs and symptoms of benzodiazepine abuse, misuse, and addiction; to seek medical help if they develop these signs and/or symptoms; and on the proper disposal of unused drug [see *Warnings and Precautions (5.2) and Drug Abuse and Dependence (9.2)*].

Withdrawal Reactions

Inform patients that use of SEIZALAM more frequently than recommended may lead to clinically significant physical dependence and that abrupt discontinuation or rapid dosage reduction of SEIZALAM may precipitate acute withdrawal reactions, which can be life-threatening. Inform patients that in some cases, patients taking benzodiazepines have developed a protracted withdrawal syndrome with withdrawal symptoms lasting weeks to more than 12 months [see *Warnings and Precautions (5.3) and Drug Abuse and Dependence (9.3)*].

Concomitant Medications

Advise patients to inform their physician about any alcohol consumption and medicine they are now taking, especially blood pressure medication and antibiotics, including drugs they buy without a prescription. Alcohol has an increased effect when consumed

with benzodiazepines; therefore, caution should be exercised regarding simultaneous ingestion of alcohol during benzodiazepine treatment [*see Warnings and Precautions (5.6), Drug Interactions (7)*].

Impaired Cognitive Function

Advise patients not to operate hazardous machinery or a motor vehicle until the effects of the drug, such as drowsiness, have subsided [*see Warnings and Precautions (5.7)*].

Pregnancy

Advise pregnant females that the use of SEIZALAM late in pregnancy can result in sedation (respiratory depression, lethargy, hypotonia) and/or withdrawal symptoms (hyperreflexia, irritability, restlessness, tremors, inconsolable crying, and feeding difficulties) in newborns [*see Warnings and Precautions (5.9) and Use in Specific Populations (8.1)*]. Instruct patients to inform their healthcare provider if they are pregnant. Encourage patients to enroll in the North American Antiepileptic Drug Pregnancy Registry if they become pregnant. This registry is collecting information about the safety of antiepileptic drugs during pregnancy [*see Use in Specific Populations (8.1)*].

Lactation

Counsel patients that midazolam, the active ingredient in SEIZALAM, is excreted in breast milk. Instruct patients to inform their healthcare provider if they are breastfeeding or intend to breastfeed. Instruct breastfeeding patients who have been administered SEIZALAM to monitor infants for excessive sedation, poor feeding and poor weight gain, and to seek medical attention if they notice these signs [*see Use in Specific Populations (8.2)*].

Manufactured by:

Hospira, Inc.
Lake Forest, IL 60045
A Pfizer Company

Distributed by:

Meridian Medical Technologies[®], LLC
St. Louis, MO 63146
0002218

Principal Display Panel - Seizalam 5mg/mL Carton Label

10 x 10 mL Multiple-Dose Vials

NDC 11704-650-10

Rx only

Seizalam[®] (midazolam injection)
for Intramuscular Use Only

50 mg / 10 mL (5 mg/mL)

For Adult Use Only

Sterile.


CIV

MERIDIAN MEDICAL TECHNOLOGIES®

10 x 10 mL Multiple-Dose Vials

NDC 11704-650-10
Rx only

Each mL contains 5 mg midazolam (equivalent to 5.6 mg midazolam hydrochloride) compounded with 1% benzyl alcohol as preservative, 0.01% edetate disodium, 0.8% sodium chloride. The pH is adjusted to approximately 3 with hydrochloric acid and, if necessary, sodium hydroxide.

Seizalam® (midazolam injection) 
for Intramuscular Use Only

50 mg / 10 mL (5 mg/mL)

For Adult Use Only

Sterile.




Sterile.

For Adult Use Only

50 mg / 10 mL
(5 mg/mL)

for Intramuscular Use Only

Seizalam® (midazolam injection) 

Rx only

NDC 11704-650-10

10 x 10 mL Multiple-Dose Vials

For Intramuscular use only.

Usual dosage: See insert.

Store at 20°C to 25°C (68°F to 77°F);
excursions permitted between 15°C and
30°C [See USP Controlled Room
Temperature.]

PAA196623

MADE IN ITALY
Manufactured by:
Hospira, Inc.
Lake Forest, IL 60045, a Pfizer company
Distributed by:
Meridian Medical Technologies®, LLC
St. Louis, MO 63146



GTIN 10311704650109
LOT AA####
SN #####
EXP DMMYYYY



(01)10311704650109

FPD - GS1 128 - 10 MII
(01)10311704650109

10 x 10 mL Multiple-Dose Vials

NDC 11704-650-10
Rx only

Each mL contains 5 mg midazolam (equivalent to 5.6 mg midazolam hydrochloride) compounded with 1% benzyl alcohol as preservative, 0.01% edetate disodium, 0.8% sodium chloride. The pH is adjusted to approximately 3 with hydrochloric acid and, if necessary, sodium hydroxide.

Seizalam® (midazolam injection)

for Intramuscular Use Only

50 mg / 10 mL (5 mg/mL)

For Adult Use Only

Sterile.



Sterile.

For Adult Use Only

50 mg / 10 mL (5 mg/mL)

for Intramuscular Use Only

Seizalam® (midazolam injection)

Rx only

NDC 11704-650-10

10 x 10 mL Multiple-Dose Vials

For Intramuscular use only.

Usual dosage: See insert.

Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C [See USP Controlled Room Temperature.]

MADE IN ISRAEL

Manufactured by:
Hospira, Inc.
Lake Forest, IL 60045, a Pfizer company
Distributed by:
Meridian Medical Technologies®, LLC
St. Louis, MO 63146

PAA196624



GTIN 10311704650109
LOT AA####
SN #####
EXP DMMYYYY



(01)10311704650109

FPO - GST 128 - 10 MII
(01)10311704650109

Principal Display Panel - Seizalam 5mg/mL Vial Label

Rx only

NDC 11704-650-01

Seizalam® CIV

(midazolam injection)

Multiple-Dose Vial for Intramuscular Use Only

50 mg / 10 mL (5 mg/mL)

For Adult Use Only

Sterile.

Rx only NDC 11704-650-01

Seizalam[®] **Ⓢ**
(midazolam injection)


Multiple-Dose Vial for Intramuscular Use Only

50 mg / 10 mL (5 mg/mL)

For Adult Use Only
Sterile.

Each mL contains 5 mg midazolam (equivalent to 5.6 mg midazolam hydrochloride) compounded with 1% benzyl alcohol as preservative, 0.01% edetate disodium, 0.8% sodium chloride. The pH is adjusted to approximately 3 with hydrochloric acid and, if necessary, sodium hydroxide. Store at 20°C to 25°C (68°F to 77°F).

Manufactured by: Hospira, Inc., Lake Forest, IL 60045
Distributed by: Meridian Medical Technologies[®], LLC
St. Louis, MO 63146



PAA196625
LOT AA###
EXP DMMYYY

SEIZALAM

midazolam hydrochloride injection, solution

Product Information

| | | | |
|-------------------------|-------------------------|--------------------|---------------|
| Product Type | HUMAN PRESCRIPTION DRUG | Item Code (Source) | NDC:11704-650 |
| Route of Administration | INTRAMUSCULAR | DEA Schedule | CIV |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|--------------|
| midazolam hydrochloride (UNII: W7TTW573JJ) (midazolam - UNII:R60L0SM5BC) | midazolam | 5 mg in 1 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|--------------------------------------|----------|
| sodium chloride (UNII: 451W47IQ8X) | |
| edetate disodium (UNII: 7FLD91C86K) | |
| benzyl alcohol (UNII: LKG8494WBH) | |
| hydrochloric acid (UNII: QTT17582CB) | |
| sodium hydroxide (UNII: 55X04QC32I) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------|----------------------|--------------------|
| 1 | NDC:11704-650-10 | 10 in 1 CARTON | 09/14/2018 | |

| | | | | |
|------------------------------|---|---|-----------------------------|---------------------------|
| 1 | NDC:11704-650-01 | 10 mL in 1 VIAL, GLASS; Type 0: Not a Combination Product | | |
| Marketing Information | | | | |
| Marketing Category | Application Number or Monograph Citation | | Marketing Start Date | Marketing End Date |
| NDA | NDA209566 | | 09/14/2018 | |

Labeler - Meridian Medical Technologies LLC (049504624)

Establishment

| Name | Address | ID/FEI | Business Operations |
|---------------|---------|-----------|---------------------|
| Hospira, Inc. | | 030606222 | ANALYSIS(11704-650) |

Establishment

| Name | Address | ID/FEI | Business Operations |
|---------------|---------|-----------|---|
| Hospira, Inc. | | 093132819 | ANALYSIS(11704-650) , LABEL(11704-650) , MANUFACTURE(11704-650) , PACK(11704-650) |

Establishment

| Name | Address | ID/FEI | Business Operations |
|---------------|---------|-----------|---------------------|
| Hospira, Inc. | | 827731089 | ANALYSIS(11704-650) |

Revised: 1/2023

Meridian Medical Technologies LLC