

ASPIRIN REGULAR STRENGTH- aspirin tablet, coated
Cardinal Health (Leader) 70000

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*

***nonsteroidal anti-inflammatory drug**

Purpose

Pain reliever

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscle pain
 - toothache
 - menstrual pain
 - colds
 - minor arthritis pain
- or as recommended by a doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of **Reye's syndrome**, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

if you are allergic to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- new symptoms occur
- ringing in the ears or loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.(1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years of age and over: take 1 to 2 tablets every 4 hours, while symptoms persist
- do not take more than 12 tablets in 24 hours unless directed by a doctor
- children under 12 years of age: do not use

Other information

- store between 20-25°C (68-77°F)

Inactive ingredients

corn starch, croscarmellose sodium, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, methacrylic acid copolymer, microcrystalline cellulose, mineral oil, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, talc, titanium dioxide, triethyl citrate

Questions or comments?

Call 1-877-753-3935 Monday-friday 9AM-5PM EST

Principal Display Panel

COMPARE TO ECOTRIN® REGULAR STRENGTH

Regular Strength

Aspirin

Tablets, 325 mg | Pain Reliever (NSAID)

Aspirin Regimen[‡]

Safety coated**

ENTERIC COATED TABLETS

†This product is not manufactured or distributed by Prestige Brands, Inc., distributor of Ecotrin®.

**Coating helps protect against stomach upset

‡Talk to your doctor or other healthcare provider before using this product for your heart.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BORKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DISTRIBUTED BY CARDINAL HEALTH

DUBLIN, OHIO 43017

www.myleader.com 1-800-200-6313

Package label

PLD-A13SD EC004916
 Lot No.:
 Exp. Date:

CIN 5425749 REV. 5/2018

0 96295 13403 2

Drug Facts (continued)
 medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years of age and over: take 1 to 2 tablets every 4 hours, while symptoms persist
- do not take more than 12 tablets in 24 hours unless directed by a doctor
- children under 12 years of age: do not use

Other information

- store between 20°-25°C (68°-77°F)

Inactive ingredients corn starch, croscarmellose sodium, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, methacrylic acid copolymer, microcrystalline cellulose, mineral oil, poly sorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, talc, titanium dioxide, triethyl citrate

Questions or comments?
 Call 1-877-753-3835 Monday-Friday 9AM-5PM EST

This product is not manufactured or distributed by Prestige Brands, Inc., distributor of Ecotrin®.

*Talk to your doctor or other healthcare provider before using this product for your heart.
 **Getting help protect against stomach upset

100% Money Back Guarantee

Return to place of purchase.

DISTRIBUTED BY CARDINAL HEALTH
 DUBLIN, OHIO 43017
 www.leadertm.com 1-800-200-655
 Essential to Care™ Since 1879

© 2018 Cardinal Health. All Rights Reserved. CARDINAL HEALTH, the Cardinal Health LOGO, LEADER, and the Leader LOGO are trademarks or registered trademarks of Cardinal Health.

Drug Facts (continued)

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - tarry, bloody or black stools
 - feel faint
 - vomit blood
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.
 KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Drug Facts

Active ingredient (in each tablet)
 Aspirin 325 mg (NSAID)* Pain reliever
 *nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscle pain
 - toothache
 - menstrual pain
 - colds
 - minor arthritis pain
- or as recommended by a doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

NDC 70000-0359-1

Regular Strength Aspirin
 Tablets, 325 mg | Pain Reliever (NSAID)

Aspirin Regimen†
 Safety Coated™

125 ENTERIC COATED TABLETS

Actual Size

COMPARE TO ECOTRIN® REGULAR STRENGTH active ingredient†

100% Money Back Guarantee

LEADER™

LEADER Aspirin 325 mg

ASPIRIN REGULAR STRENGTH

aspirin tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70000-0359
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
MINERAL OIL (UNII: T5L8T28FGP)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
ALUMINUM OXIDE (UNII: LM26O6933)	

Product Characteristics

Color	ORANGE	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	T
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70000-0359-1	1 in 1 BOX	03/31/2018	12/31/2022
1		125 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part343	03/31/2018	12/31/2022

Labeler - Cardinal Health (Leader) 70000 (097537435)