

**CICLOPIROX- ciclopirox shampoo**  
**Encube Ethicals, Inc.**

---

**HIGHLIGHTS OF PRESCRIBING INFORMATION**

These highlights do not include all the information needed to use Ciclopirox Shampoo safely and effectively. See full prescribing information for **CICLOPIROX SHAMPOO**.

**Ciclopirox Shampoo 1% For topical use**

**Initial U.S. Approval: 1982**

**INDICATIONS AND USAGE**

- Ciclopirox shampoo is an antifungal indicated for the topical treatment of seborrheic dermatitis of the scalp in adults. (1)

**DOSAGE AND ADMINISTRATION**

- Apply approximately 1 teaspoon of ciclopirox shampoo to the scalp twice per week for 4 weeks. (2)
- For topical use only. Not for ophthalmic, oral, or intravaginal use. (2)

**DOSAGE FORMS AND STRENGTHS**

Shampoo, 1% (3)

**CONTRAINDICATIONS**

None (4)

**WARNINGS AND PRECAUTIONS**

- If signs of irritation occur, discontinue use. (5.1)
- Avoid contact with eyes. (5.1)
- Hair discoloration has been reported with ciclopirox use. (5.1)

**ADVERSE REACTIONS**

- The most frequently reported adverse reactions are pruritus, burning, and erythema. (6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact Encube Ethicals Private Limited at 1-833-285-4151, and/or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

**See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.**

**Revised: 1/2026**

---

**FULL PRESCRIBING INFORMATION: CONTENTS\***

**1 INDICATIONS AND USAGE**

**2 DOSAGE AND ADMINISTRATION**

**3 DOSAGE FORMS AND STRENGTHS**

**4 CONTRAINDICATIONS**

**5 WARNINGS AND PRECAUTIONS**

5.1 Local Effects

**6 ADVERSE REACTIONS**

6.1 CLINICAL TRIALS EXPERIENCE

6.2 POSTMARKETING EXPERIENCE

**8 USE IN SPECIFIC POPULATIONS**

8.1 PREGNANCY

8.3 NURSING MOTHERS

8.4 PEDIATRIC USE

8.5 GERIATRIC USE

## **11 DESCRIPTION**

## **12 CLINICAL PHARMACOLOGY**

- 12.1 MECHANISM OF ACTION
- 12.2 PHARMACODYNAMICS
- 12.3 PHARMACOKINETICS
- 12.4 MICROBIOLOGY

## **13 NONCLINICAL TOXICOLOGY**

- 13.1 CARCINOGENESIS & MUTAGENESIS & IMPAIRMENT OF FERTILITY

## **14 CLINICAL STUDIES**

## **16 HOW SUPPLIED**

## **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

Ciclopirox shampoo, 1 % is indicated for the topical treatment of seborrheic dermatitis of the scalp in adults.

### **2 DOSAGE AND ADMINISTRATION**

Ciclopirox shampoo is not for ophthalmic, oral, or intravaginal use.

Wet hair and apply approximately 1 teaspoon (5 mL) of ciclopirox shampoo to the scalp. Up to 2 teaspoons (10 ml) may be used for long hair. Lather and leave on hair and scalp for 3 minutes. A timer may be used. Avoid contact with eyes. Rinse off. Treatment should be repeated twice per week for 4 weeks, with a minimum of 3 days between applications.

If a patient with seborrheic dermatitis shows no clinical improvement after 4 weeks of treatment with ciclopirox shampoo, the diagnosis should be reviewed.

### **3 DOSAGE FORMS AND STRENGTHS**

Ciclopirox is a shampoo containing 1 % ciclopirox.

Each gram (equivalent to 0.96 mL) of ciclopirox shampoo contains 10 mg ciclopirox in a colorless and translucent shampoo base.

### **4 CONTRAINDICATIONS**

None.

### **5 WARNINGS AND PRECAUTIONS**

## **5.1 Local Effects**

If a reaction suggesting sensitivity or irritation occurs with the use of ciclopirox shampoo, treatment should be discontinued and appropriate therapy instituted.

Contact of ciclopirox shampoo with the eyes should be avoided. If contact occurs, rinse thoroughly with water.

In patients with lighter hair color, hair discoloration has been reported.

## **6 ADVERSE REACTIONS**

### **6.1 CLINICAL TRIALS EXPERIENCE**

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

In 626 subjects treated with ciclopirox shampoo twice weekly in the two pivotal clinical trials, the most frequent adverse events were increased itching in 1% of subjects, and application site reactions, such as burning, erythema, and itching, also in 1 % of subjects.

### **6.2 POSTMARKETING EXPERIENCE**

The following adverse reactions have been identified during post approval use of ciclopirox shampoo: hair discoloration and abnormal hair texture, alopecia, irritation and rash. Because these events are reported voluntarily from a population of uncertain size, it is not possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 PREGNANCY**

Teratogenic Effects: Pregnancy Category B

There are no adequate or well-controlled studies in pregnant women. Therefore, ciclopirox shampoo should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Oral embryofetal developmental studies were conducted in mice, rats, rabbits and monkeys. Ciclopirox or ciclopirox olamine was orally administered during the period of organogenesis. No maternal toxicity, embryotoxicity or teratogenicity were noted at the highest doses of 77, 125, 80 and 38.5 mg/kg/day ciclopirox in mice, rats, rabbits and monkeys, respectively (approximately 13, 42, 54 and 26 times the maximum recommended human dose based on body surface area comparisons, respectively).

Dermal embryofetal developmental studies were conducted in rats and rabbits with ciclopirox olamine dissolved in PEG 400. Ciclopirox olamine was topically administered during the period of organogenesis. No maternal toxicity, embryotoxicity or teratogenicity were noted at the highest doses of 92 mg/kg/day and 77 mg/kg/day ciclopirox in rats and rabbits, respectively (approximately 31 and 54 times the maximum

recommended human dose based on body surface area comparisons, respectively).

### **8.3 NURSING MOTHERS**

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when ciclopirox shampoo is administered to a nursing woman.

### **8.4 PEDIATRIC USE**

No clinical trials have been conducted in subjects younger than 16 years.


### **8.5 GERIATRIC USE**

In clinical trials, the safety and tolerability of ciclopirox shampoo in the population 65 years and older was comparable to that of younger subjects. Results of the efficacy analysis in those subjects 65 years and older showed effectiveness in 25 of 85 (29%) subjects treated with ciclopirox shampoo, and in 15 of 61 (25%) subjects treated with the vehicle; due to the small sample size, a statistically significant difference was not demonstrated. Other reported clinical experience has not identified differences in responses between the elderly and younger subjects, but greater sensitivity to adverse effects in some older individuals cannot be ruled out.

## **11 DESCRIPTION**

Ciclopirox shampoo 1 % contains the synthetic antifungal agent, ciclopirox for topical use. Each gram (equivalent to 0.96 ml) of ciclopirox shampoo contains 10 mg ciclopirox in a shampoo base consisting of disodium laureth sulfosuccinate, laureth-2, purified water, sodium chloride, and sodium lauryl sulfate.

Ciclopirox shampoo is a colorless, translucent solution. The chemical name for ciclopirox is 6-cyclohexyl-1-hydroxy-4-methyl-2(1H)-pyridone, with the empirical formula C<sub>12</sub>H<sub>17</sub>NO<sub>2</sub> and a molecular weight of 207.27. The GAS Registry Number is [29342-05-0]. The chemical structure is:



## **12 CLINICAL PHARMACOLOGY**

### **12.1 MECHANISM OF ACTION**

Ciclopirox is a hydroxypyridone antifungal agent although the relevance of this property for the indication of seborrheic dermatitis is not known. Ciclopirox acts by chelation of

polyvalent cations (Fe<sup>3+</sup> or Al<sup>3+</sup>), resulting in the inhibition of the metal-dependent enzymes that are responsible for the degradation of peroxides within the fungal cell.

## **12.2 PHARMACODYNAMICS**

The pharmacodynamics of ciclopirox shampoo are unknown.

## **12.3 PHARMACOKINETICS**

In a study in patients with seborrheic dermatitis of the scalp, application of 5 mL ciclopirox shampoo, 1 % twice weekly for 4 weeks, with an exposure time of 3 minutes per application, resulted in detectable serum concentrations of ciclopirox in 6 out of 18 patients. The serum concentrations measured throughout the dosing interval on Days 1 and 29 ranged from 10.3 ng/mL to 13.2 ng/ml. Total urinary excretion of ciclopirox was less than 0.5% of the administered dose.

## **12.4 MICROBIOLOGY**

Ciclopirox is fungicidal *in vitro* against *Malassezia furfur* (*Pityrosporum spp.*), *P. ovale*, and *P. orbiculare*.

The clinical significance of antifungal activity in the treatment of seborrheic dermatitis is not known.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 CARCINOGENESIS & MUTAGENESIS & IMPAIRMENT OF FERTILITY**

A 104-week dermal carcinogenicity study in mice was conducted with ciclopirox cream applied at doses up to 1.93% (100 mg/kg/day or 300 mg/m<sup>2</sup>/day). No increase in drug related neoplasms was noted when compared to control.

The following *In vitro* genotoxicity tests have been conducted with ciclopirox: evaluation of gene mutation in the Ames Salmonella and *E. coli* assays (negative); chromosome aberration assays in V79 Chinese hamster lung fibroblast cells, with and without metabolic activation (positive); chromosome aberration assays in V79 Chinese hamster lung fibroblast cells in the presence of supplemental Fe<sup>3+</sup>, with and without metabolic activation (negative); gene mutation assays in the HGPRT-test with V79 Chinese hamster lung fibroblast cells (negative); and a primary DNA damage assay (i.e., unscheduled DNA synthesis assay in A549 human cells) (negative). An *In vitro* cell transformation assay in BALB/c 3T3 cells was negative for cell transformation. In an *In vivo* Chinese hamster bone marrow cytogenetic assay, ciclopirox was negative for chromosome aberrations at a dosage of 5,000 mg/kg body weight.

A combined oral fertility and embryofetal developmental study was conducted in rats with ciclopirox olamine. No effect on fertility or reproductive performance was noted at the highest dose tested of 3.85 mg/kg/day ciclopirox (approximately 1.3 times the maximum recommended human dose based on body surface area comparisons).

## **14 CLINICAL STUDIES**

In two randomized, double-blind clinical trials, subjects 16 years and older with

seborrheic dermatitis of the scalp applied ciclopirox shampoo or its vehicle twice weekly for 4 weeks. Subjects who were immune- compromised, those with psoriasis or atopic dermatitis, women of childbearing potential not using adequate contraception, and pregnant or lactating women were excluded from the clinical trials. An evaluation of the overall status of the seborrheic dermatitis, the presence and severity of erythema or inflammation, and scaling, was made at Week 4, using a scale of 0 = none, 1 = slight, 2 = mild, 3 = moderate, 4 = pronounced, and 5 = severe. Effective treatment was defined as achieving a score of 0 (or a score of 1 if the baseline score was  $\geq 3$ ) simultaneously for status of the seborrheic dermatitis, erythema or inflammation, and scaling at Week 4.

Ciclopirox shampoo was shown to be statistically significantly more effective than vehicle in both trials. Efficacy results for the two trials are presented in Table 1 below.

**Table 1. Effective Treatment Rates at Week 4 in Trials 1 and 2**

	<b>Ciclopirox Shampoo</b>	<b>Vehicle</b>
<b>Study 1</b>	<b>220/380 (58%)</b>	<b>60/192 (31%)</b>
<b>Study 2</b>	<b>65/250 (26%)</b>	<b>32/249 (13%)</b>

Efficacy for African American subjects was not demonstrated, although only 53 African American subjects were enrolled in the two pivotal trials.

## **16 HOW SUPPLIED**

Ciclopirox shampoo, 1 % is colorless and translucent, and supplied in 120 mL plastic bottles (NOC 21922-105-52).

Discard unused product after initial treatment duration. Store between 15°C and 30°C (59° F and 86° F).

Keep out of reach of children.

## **17 PATIENT COUNSELING INFORMATION**

Advise the patient to read the FDA-approved patient labeling (Patient Information).

The patient should be instructed to:

- Use ciclopirox shampoo as directed by the physician. Avoid contact with the eyes. If contact occurs, rinse thoroughly with water. Ciclopirox shampoo is for external use on the scalp only. Do not swallow.
- Use ciclopirox shampoo for seborrheic dermatitis for the full treatment time even though symptoms may have improved. Notify the physician if there is no improvement after 4 weeks.
- Inform the physician if the area of application shows signs of increased irritation (redness, itching, burning, blistering, swelling, or oozing).

Manufactured by:

**Encube Ethicals Pvt. Ltd.**

Plot No. C-1, Madkaim  
Industrial Estate, Madkaim,  
Post: Mardol, Panda,  
Goa - 403 404, India.

Distributed by:

**Encube Ethicals, Inc.**

200 Meredith Drive,  
Suite 202, Durham,  
NC27713 USA

**Revision: 05/2025**

## **PATIENT INFORMATION**

Ciclopirox (sye" klo pir' ox) Shampoo

Important: For use on the scalp only. Do not get ciclopirox shampoo in your eyes, mouth, or vagina.

### **What is ciclopirox shampoo?**

Ciclopirox shampoo is a prescription medicine used on the scalp to treat adults with a skin condition called seborrheic dermatitis.

It is not known if ciclopirox shampoo is safe and effective in children under 16 years of age.

### **What should I tell my doctor before using ciclopirox shampoo?**

Before using ciclopirox shampoo, tell your doctor if you:

- have any other medical conditions
- are pregnant or plan to become pregnant. It is not known if ciclopirox shampoo will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if ciclopirox shampoo passes into your breast milk.
- are taking prescription and nonprescription medicines, vitamins, and herbal supplements.

Know the medicines you take. Keep a list of them to show your doctor and pharmacist when you get a new medicine.

### **How should I use ciclopirox shampoo?**

- Use ciclopirox shampoo exactly as your doctor tells you to use it.
- Wash your hair using ciclopirox shampoo 2 times each week for 4 weeks. There should be at least 3 days between each time you use ciclopirox shampoo.
- Use ciclopirox shampoo for 4 weeks even if your skin condition improves.
- Tell your doctor if your scalp condition is not getting better after you have used

ciclopirox shampoo for 4 weeks.

- Do not swallow ciclopirox shampoo.
- Avoid getting ciclopirox shampoo in your eyes. If ciclopirox shampoo gets into your eyes, rinse them well with water.

### **How should I apply ciclopirox shampoo?**

- Wet your hair and apply approximately 1 teaspoon of ciclopirox shampoo to your scalp. You may use up to 2 teaspoons of ciclopirox shampoo if you have long hair. Lather and leave ciclopirox shampoo on your hair and scalp for 3 minutes. You may use a timer.
- After 3 minutes have passed, rinse your hair and scalp.

### **What are the possible side effects of ciclopirox shampoo?**

**The most common side effects of ciclopirox shampoo include:** itching, burning, and redness of the scalp. Tell your doctor if you get any of these symptoms and they become worse or do not go away, or if you get blistering, swelling, or oozing in your scalp.

These are not all the possible side effects of ciclopirox shampoo. For more information, ask your doctor.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

### **How should I store ciclopirox shampoo?**

- Store ciclopirox shampoo at room temperature, between 59°F to 86°F (15°C to 30°C).
- Safely throw away any unused ciclopirox shampoo after you finish your treatment.

### **Keep ciclopirox shampoo and all medicines out of the reach of children.**

### **General information about ciclopirox shampoo**

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use ciclopirox shampoo for a condition for which it was not prescribed. Do not give ciclopirox shampoo to other people, even if they have the same symptoms you have. It may harm them.

You can ask your pharmacist or doctor for information about ciclopirox shampoo that is written for health professionals.

For more information about ciclopirox shampoo, call 1-833-285-4151.

### **What are the ingredients in ciclopirox shampoo?**

**Active ingredient:** ciclopirox

**Inactive ingredients:** disodium laureth sulfosuccinate, laureth-2, purified water, sodium chloride, and sodium laureth sulfate.

This Patient Information has been approved by the U.S. Food and Drug Administration.

Manufactured by:

**Encube Ethicals Pvt. Ltd.**

PlotNo. C-1, Madkaim

Industrial Estate, Madkaim,

Post: Mardol, Panda,

Goa - 403 404, India.

Distributed by:

**Encube Ethicals, Inc.**

200 Meredith Drive,

Suite 202, Durham,

NC 27713 USA

**Revision: 05/2025**

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

NDC 21922-**105**-52

**Ciclopirox Shampoo 1%**

For Topical Use Only


Not for Ophthalmic,

Oral or Intravaginal Use.


Keep Out of Reach of Children

**RX only**

120 mL



EBX3A/00  
Rev: 05/25



## CICLOPIROX

ciclopirox shampoo

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDG:21922-105
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CICLOPIROX (UNII: 19W019Z DRJ) (CICLOPIROX - UNII:19W019Z DRJ)	CICLOPIROX	1 g in 100 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>SODIUM LAURETH-2 SULFATE</b> (UNII: ZZQ59TY3KG)	
<b>DISODIUM LAURETH SULFOSUCCINATE</b> (UNII: D6DH1DTN7E)	
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	
<b>LAURETH-2</b> (UNII: D4D38LT1L5)	
<b>WATER</b> (UNII: 059QF0KO0R)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21922-105-52	1 in 1 CARTON	12/03/2025	
1		120 mL in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA209975	12/03/2025	

**Labeler** - Encube Ethicals, Inc. (116982244)

## Establishment

Name	Address	ID/FEI	Business Operations
Encube Ethicals Private Limited		725076298	pack(21922-105) , label(21922-105) , manufacture(21922-105) , analysis(21922-105)

Revised: 1/2026

Encube Ethicals, Inc.