ACID CONTROLLER- famotidine tablet, film coated H E B

HEB Acid Controller Drug Facts

Active ingredient (in each tablet)

Famotidine 20 mg

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- if you have kidney disease, except under the advice and supervision of a doctor
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 12 years and over:
- to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture and light

Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, lactose (monohydrate), magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, and titanium dioxide

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to Maximum Strength Pepcid® AC active ingredient

Maximum Strength

Acid Controller

Famotidine Tablets, 20 mg

Acid Reducer

Just One Tablet Prevents & Relieves Heartburn Due To Acid Indigestion

actual size

200 TABLETS





Drug Facts

Active ingredient Purpose (in each tablet)

Uses

- relieves heartburn associated with acid indigestion and sour stomach

 prevents heartburn associated with acid indigestion
- and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

 if you have kidney disease, except under the advice
- and supervision of a doctor
- with other acid reducers
- Ask a doctor before use if you have

 had heartburn over 3 months. This may be a sign of a
- more serious condition.

 hearthum with lightheadedness, sweating, or dizziness
 chest pain or shoulder pain with shortness of breath;
 sweating; pain spreading to arms, neck or shoulders;
- or lightheadedness

 frequent chest pain

 frequent wheezing, particularly with heartburn
 unexplained weight loss
- nausea or vomiting

Drug Facts (continued)

- Stop use and ask a doctor if
 your heartburn continues or worsens
 you need to take this product for more than 14 days
- If pregnant or breast-feeding, ask a health sional before use

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center stable outs:

Directions

- adults and children 12 years and over:
 to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause hearthurn
- do not use more than 2 tablets in 24 hours children under 12 years: ask a doctor

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
 store at 20°-25°C (68°-77°F)
 protect from moisture and light

Inactive ingredients carnauba wax, colloidal silicon dioxide, croscarmellose sodium, lactose (monohydrate), magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, and titanium dioxide

(Questions or comments?

Compare to Maximum Strength Pepcid® AC active Ingredient



Maximum Strength

Acid Controller

Famotidine Tablets, 20 mg



Just One Tablet Prevents & Relieves Heartburn Due to Acid Indigestion

200 TABLETS

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- · Do not eat before bedtime
- · Raise the head of your bed
- Avoid heartburn causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- · Eat slowly and avoid big meals
- · If overweight, lose weight
- Quit smoking


DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING

*This product is not manufactured or distributed by Johnson & Johnson • MERCK Consumer
Pharmaceuticals Co., distributor of Maximum Strength Pepcid® AC.

MADE WITH PRIDE & CARE FOR H-E-B® SAN ANTONIO, TX 78204

GLUTEN FREE







ACID CONTROLLER

famotidine tablet, film coated

Product Information

HUMAN OTC DRUG NDC:37808-042 Product Type Item Code (Source)

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength
l	FAMO TIDINE (UNII: 50 ZO 15,12Z8) (FAMO TIDINE - UNII: 50 ZO 15,12Z8)	FAMOTIDINE	20 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIO XIDE (UNII: ETJ7Z6 XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28 OL1HH48)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICRO CRYSTALLINE CELLULO SE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIO XIDE (UNII: 15FIX9 V2JP)	

Product Characteristics				
Color	WHITE	Score	no score	
Shape	ROUND	Size	8 m m	
Flavor		Imprint Code	L194	
Contains				

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:37808-042-82	1 in 1 CARTON	09/14/2016		
1		200 in 1 BOTTLE; Type 0: Not a Combination Product			
2	NDC:37808-042-63	1 in 1 CARTON	0 1/27/20 20		
2		25 in 1 BOTTLE; Type 0: Not a Combination Product			

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA077351	09/14/2016		

Labeler - HEB (007924756)

Revised: 1/2020 HE B