

**STONARHINI- chlorpheniramine maleate, phenylephrine tablet**  
**Sato Pharmaceutical Co., Ltd.**

-----  
**Stonarhini**

**Active ingredients** (In each tablet)

Chlorpheniramine Maleate 2 mg

Phenylephrine HCl 5 mg

**Purpose**

Chlorpheniramine Maleate     Antihistamine

Phenylephrine HCl         Nasal decongestant

**Uses**

Temporarily relieves these symptoms due to common cold, hay fever or other upper respiratory allergies (allergic rhinitis).

- sneezing         ■ runny nose         ■ itchy, watery eyes
- nasal congestion, stuffy nose         ■ itchy throat

**Warnings**

Enter section text here

**Do not use this product**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma     ■ heart disease     ■ thyroid disease
- diabetes     ■ high blood pressure
- difficulty in urination due to enlargement of the prostate

**Ask a doctor or pharmacist before use**

- if you are taking sedatives or tranquilizers

**When using this product**

- do not exceed recommended dosage. If nervousness, dizziness, or sleeplessness occur, discontinue use and consult a doctor.
- you may get drowsy.
- may cause excitability, especially in children.
- alcohol, sedatives and tranquilizers may increase the drowsiness effect.
- avoid alcoholic beverages.
- use caution when driving a motor vehicle or operating machinery.

**Stop use and ask a doctor if**

- symptoms do not improve within seven days
- symptoms are accompanied by fever

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt attention is critical even if you do not notice any signs or symptoms.

***Directions***

adults and children 12 years and over: 2 tablets every 4 hours, not to exceed 12 tablets in 24 hours

children under 12 years of age: ask a doctor

***Inactive ingredients*** aspartame, colloidal silicone dioxide, FD&C Red No. 40 aluminum Lake, flavor, hydroxypropyl cellulose, magnesium stearate, mannitol.



## STONARHINI

chlorpheniramine maleate, phenylephrine tablet

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49873-109
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)		CHLORPHENIRAMINE MALEATE	2 mg	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)		PHENYLEPHRINE HYDROCHLORIDE	5 mg	
Inactive Ingredients				
Ingredient Name			Strength	
ASPARTAME (UNII: Z0H242BBR1)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
HYDROXYPROPYL CELLULOSE (TYPE EL) (UNII: 8VAB711C5E)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
MANNITOL (UNII: 3OWL53L36A)				
ALUMINUM OXIDE (UNII: LMI26O6933)				
Product Characteristics				
Color	red (pale red)	Score	no score	
Shape	ROUND	Size	10mm	
Flavor	MENTHOL	Imprint Code	S10	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49873-109-01	4 in 1 CARTON	11/16/2001	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012		11/16/2001	

**Labeler** - Sato Pharmaceutical Co., Ltd. (690575642)

## Establishment

Name	Address	ID/FEI	Business Operations
Sato Pharmaceutical Co., Ltd.		715699133	manufacture(49873-109) , label(49873-109) , pack(49873-109)

