

**STONARHINI- chlorpheniramine maleate, phenylephrine tablet**  
**Sato Pharmaceutical Co., Ltd.**

-----  
**Stonarhini**

**Active ingredients** (In each tablet)

Chlorpheniramine Maleate 2 mg

Phenylephrine HCl 5 mg

**Purpose**

Chlorpheniramine Maleate    Antihistamine

Phenylephrine HCl        Nasal decongestant

**Uses**

Temporarily relieves these symptoms due to common cold, hay fever or other upper respiratory allergies (allergic rhinitis).

- sneezing        ■ runny nose        ■ itchy, watery eyes
- nasal congestion, stuffy nose        ■ itchy throat

**Warnings**

Enter section text here

**Do not use this product**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma    ■ heart disease    ■ thyroid disease
- diabetes    ■ high blood pressure
- difficulty in urination due to enlargement of the prostate

**Ask a doctor or pharmacist before use**

- if you are taking sedatives or tranquilizers

**When using this product**

- do not exceed recommended dosage. If nervousness, dizziness, or sleeplessness occur, discontinue use and consult a doctor.
- you may get drowsy.
- may cause excitability, especially in children.
- alcohol, sedatives and tranquilizers may increase the drowsiness effect.
- avoid alcoholic beverages.
- use caution when driving a motor vehicle or operating machinery.

**Stop use and ask a doctor if**

- symptoms do not improve within seven days
- symptoms are accompanied by fever

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt attention is critical even if you do not notice any signs or symptoms.

***Directions***

adults and children 12 years and over: 2 tablets every 4 hours, not to exceed 12 tablets in 24 hours

children under 12 years of age: ask a doctor

***Inactive ingredients*** aspartame, colloidal silicone dioxide, FD&C Red No. 40 aluminum Lake, flavor, hydroxypropyl cellulose, magnesium stearate, mannitol.



## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>ASPARTAME</b> (UNII: Z0H242BBR1)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>HYDROXYPROPYL CELLULOSE (TYPE EL)</b> (UNII: 8VAB711C5E)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>ALUMINUM OXIDE</b> (UNII: LMI26O6933)	

## Product Characteristics

<b>Color</b>	red (pale red)	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>	MENTHOL	<b>Imprint Code</b>	S10
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49873-109-01	4 in 1 CARTON	11/16/2001	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/16/2001	

**Labeler** - Sato Pharmaceutical Co., Ltd. (690575642)

## Establishment

Name	Address	ID/FEI	Business Operations
Sato Pharmaceutical Co., Ltd.		715699133	manufacture(49873-109) , label(49873-109) , pack(49873-109)

## Establishment

Name	Address	ID/FEI	Business Operations
Fuji Seal, Inc.		718023935	pack(49873-109) , label(49873-109)

Revised: 1/2025

Sato Pharmaceutical Co., Ltd.