

**COUNTERACT CHILDRENS PAIN RELIEVER- acetaminophen suspension
Melaleuca, Inc.**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

CounterAct Children's Pain Reliever Content of label

Active ingredient (in each 5 mL teaspoon)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Use temporarily:

- reduces fever
- relieves minor aches and pains due to:
 - the common cold
 - flu
 - headache
 - sore throat
 - toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Sore throat warning: If sore throat is severe, or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- do not give more than directed (see Liver warning)
- shake well before use
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup designed for use with this product. Do not use any other dosing

device.

- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give for more than 5 days unless directed by a doctor
- **this product does not contain directions or complete warnings for adult use**

Weight (lb)	Age (yr)	Dose (tsp or mL)
under 24	under 2 years	ask a doctor
24-35	2-3 years	1 tsp or 5 mL
36-47	4-5 years	1 1/2 tsp or 7.5 mL
48-59	6-8 years	2 tsp or 10 mL
60-71	9-10 years	2 1/2 tsp or 12.5 mL
72-95	11 years	3 tsp or 15 mL

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device/

Other information

- store at 20°-25°C (68°-77°F)
- **each teaspoon contains:** sodium 3 mg
- do not use if printed neckband is broken or missing

Inactive ingredients anhydrous citric acid, butylparaben, carboxymethylcellulose sodium, carrageenan, FD and C red no. 40, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose, propylene glycol, purified water, sodium benzoate, sorbitol solution

Questions or comments?

1-800-282-3000

Active ingredient (in each 5 mL = 1 teaspoon) Purpose
 Acetaminophen 160 mg.....Pain reliever/fever reducer

Uses temporarily:

- reduces fever
- relieves minor aches and pains due to:
 - the common cold
 - flu ■ headache ■ sore throat ■ toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

 These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- do not give more than directed (see Liver warning)
- shake well before using
- find right dose on chart. If possible, use weight to dose; otherwise, use age.
- use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.

RX
MELALEUCA PHARMACY

CounterAct®

SEE NEW WARNINGS INFORMATION

Children's
Pain Reliever



Acetaminophen Oral Suspension Liquid
Fever Reducer Pain Reliever

80 mg per
 1/2 teaspoon
 (160 mg per 5 mL)

4 FL OZ (118 mL)

**DO NOT USE
 IF PRINTED
 NECKBAND
 IS BROKEN
 OR MISSING**

- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give for more than 5 days unless directed by a doctor
- **this product does not contain directions or complete warnings for adult use**

Weight (lb)	Age (yr)	Dose (tsp or mL)
under 24	under 2 years	ask a doctor
24-35	2-3 years	1 tsp or 5 mL
36-47	4-5 years	1 1/2 tsp or 7.5 mL
48-59	6-8 years	2 tsp or 10 mL
60-71	9-10 years	2 1/2 tsp or 12.5 mL
72-95	11 years	3 tsp or 15 mL

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information ■ each teaspoon contains: sodium 3 mg ■ store at 20°-25°C (68°-77°F)

Inactive ingredients anhydrous citric acid, butylparaben, carboxymethylcellulose sodium, carrageenan, FD&C red no. 40, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose, propylene glycol, purified water, sodium benzoate, sorbitol solution


Questions or comments? 1-800-282-3000

Dist. by: Melaleuca, Inc.
 Idaho Falls, ID 83402-6003
 To order: 1-800-282-3000
 www.melaleuca.com
 #2255 08/09U
 100% Satisfaction Guarantee

CODE AREA
 • no varnish • no color

: 17526 MB F5

538H925/1



CounterAct®

EFB 839
175-002 EFB839-1

Children's
Pain Reliever
Acetaminophen

Fever Reducer/Pain Reliever

Drug Facts

Active ingredient Purpose
(in each 5 mL = 1 teaspoon)

Acetaminophen 160 mg... Pain reliever/fever reducer.

Uses

- temporarily:
 - reduces fever
 - relieves minor aches and pains due to:
 - the common cold
 - flu
 - headache
 - sore throat
 - toothache

Warnings

- Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:
 - more than 5 doses in 24 hours, which is the maximum daily amount
 - with other drugs containing acetaminophen
- Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- do not give more than directed (see Liver warning)
- shake well before using
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

Drug Facts (continued)

- use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give for more than 5 days unless directed by a doctor
- this product does not contain directions or complete warnings for adult use**

Weight (lb)	Age (yr)	Dose (tsp or mL)
under 24	under 2 years	ask a doctor
24-35	2-3 years	1 tsp or 5 mL
36-47	4-5 years	1 1/2 tsp or 7.5 mL
48-59	6-8 years	2 tsp or 10 mL
60-71	9-10 years	2 1/2 tsp or 12.5 mL
72-95	11 years	3 tsp or 15 mL

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each teaspoon contains: sodium 3 mg
- store at 20°-25°C (68°-77°F)
- do not use if printed neckband is broken or missing

Inactive ingredients anhydrous citric acid, butylparaben, carboxymethylcellulose sodium, carrageenan, FD&C red no. 40, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose, propylene glycol, purified water, sodium benzoate, sorbitol solution

(Questions or comments?)
1-800-282-3000



Dist. by: Melaleuca, Inc.
Idaho Falls, ID 83402-6003
To order: 1-800-282-3000
www.melaleuca.com
#2255 08/09U

NOT FOR RESALE IN CANADA

CounterAct®

SEE NEW WARNINGS INFORMATION



Children's
Pain Reliever

Acetaminophen
Oral Suspension Liquid

Fever Reducer/Pain Reliever

- ALCOHOL-FREE
- ASPIRIN-FREE
- IBUPROFEN-FREE

Recommended for fast, effective relief of children's pain and fever

80 mg per 1/2 teaspoon
(160 mg per 5 mL)

4 FL OZ (118 mL)



Read instructions carefully

CounterAct®

SEE NEW WARNINGS INFORMATION



Children's
Pain Reliever

Acetaminophen
Oral Suspension Liquid


Fever Reducer/Pain Reliever

- ALCOHOL-FREE
- ASPIRIN-FREE
- IBUPROFEN-FREE

Recommended for fast, effective relief of children's pain and fever

80 mg per 1/2 teaspoon
(160 mg per 5 mL)

4 FL OZ (118 mL)



Read instructions carefully



17526HBC5



LOT NO.

EXP.

Code Area

:17526 NB C5

acetaminophen suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:54473-098
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
BUTYLPARABEN (UNII: 3QP1I1U3FV8)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
CARRAGEENAN (UNII: 5C69YCD2YJ)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
HYDROXYETHYL CELLULOSE (4000 MPAS FOR 1% AQUEOUS SOLUTION) (UNII: ZYD53NBL45)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	red (Red)	Score	
Shape		Size	
Flavor	CHERRY (cherry)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54473-098-04	1 in 1 BOX		
1		5 mL in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	01/12/2010	

Revised: 3/2011

Melaleuca, Inc.