

**DAY TIME COLD AND FLU RELIEF NON DROWSY- acetaminophen,  
dextromethorphan hydrobromide, phenylephrine hydrochloride liquid  
Safeway, Inc.**

-----

**Drug Facts**

**Active ingredients (in each 15 mL)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

**Purpose**

**Pain reliever/fever reducer**

Cough suppressant

Nasal decongestant

**Uses**

temporarily relieves common cold/flu symptoms:

- minor aches and pains
- headache
- sore throat
- fever
- nasal congestion
- cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- adult takes more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount
- child takes more than 4 doses (15 mL each) in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks everyday while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- a sodium-restricted diet
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if**

you are taking the blood thinning drug warfarin.

**When using this product,**

**do not exceed recommended dosage.**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 5 days (children) or 7 days (adult)
- fever gets worse, or last more than 3 days
- redness or swelling is present
- new symptom occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center

(1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

- **do not take more than directed ((see Overdose warning)**
- do not take more than 4 doses in any 24-hours period
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product
- mL = milliliter

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children 4 to under 6 years	ask a doctor
children under 4 years	do not use

- **when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing**

### Other information

- **each 15 mL contains:** sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate

### Inactive ingredients

citric acid, FD&C yellow 6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

### Questions or comments?

Call **1-888-723-3929 Monday-Friday 7AM-6PM PST**

### Principal Display Panel

Compare to Vicks® DayQuil® active ingredients\*

Non-Drowsy

### Day Time

### Cold & Flu Relief

**Acetaminophen 325 mg** - Pain reliever/fever reducer

Dextromethorphan HBr 10mg - Cough suppressant

Phenylephrine HCl 5 mg - Nasal Decongestant

ORIGINAL FLAVOR

- Relieves aches, fever, sore throat, cough & nasal congestion
- Alcohol free
- Antihistamine free

For Ages 6 & Over

FL OZ (mL)

\*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and DayQuil® are registered trademarks of The Procter & Gamble Company.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER THE CAP IS BROKEN OR MISSING.**

DISTRIBUTED BY

BETTER LIVING BRANDS LLC

P.O. BOX 99, PLEASANTON, CA

94566-0009

[www.betterlivingbrandsLLC.com](http://www.betterlivingbrandsLLC.com)

**Product Label**

**Drug Facts****Active ingredients (in each 15 mL) Purposes**

Acetaminophen 325 mg.....Pain reliever/fever reducer  
 Dextromethorphan HBr 10 mg.....Cough suppressant  
 Phenylephrine HCl 5 mg.....Nasal decongestant

**Uses**

temporarily relieves common cold and flu symptoms  
 ■ minor aches and pains ■ headache ■ sore throat  
 ■ nasal congestion ■ fever  
 ■ cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if: ■ adult takes more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount ■ child takes more than 4 doses (15 mL each) in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash  
 If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

← PEEL HERE **Drug Facts** (continued under label)



Compare to  
 Vicks® DayQuil®  
 active ingredients\*

NDC 21130-866-08

# Non-Drowsy Daytime Cold & Flu Relief

Acetaminophen 325 mg - Pain Reliever/Fever Reducer  
 Dextromethorphan HBr 10 mg - Cough Suppressant  
 Phenylephrine HCl 5 mg - Nasal Decongestant

**ORIGINAL FLAVOR**

- Relieves aches, fever, sore throat, cough & nasal congestion
- Alcohol-free
- Antihistamine-free

For Ages 6 & Over

8 FL OZ (237 mL)

\*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and DayQuil® are registered trademarks of The Procter & Gamble Company.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

**PARENTS:**

Learn about teen medicine abuse.  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)



Scan here  
 for more  
 information

DISTRIBUTED BY  
 BETTER LIVING BRANDS LLC  
 P.O. BOX 99, PLEASANTON, CA  
 94566-0099  
 1-888-723-3929  
[www.betterlivingbrandsLLC.com](http://www.betterlivingbrandsLLC.com)

**OUR PROMISE**  
 QUALITY & SATISFACTION  
 100% GUARANTEED  
 ON YOUR MONEY BACK.

S5987 RD20022 PLD-03800 LB007385



3 2 1130 78888 1

**Drug Facts (continued)**

■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have** ■ liver disease  
 ■ heart disease ■ high blood pressure ■ thyroid disease  
 ■ diabetes ■ a sodium-restricted diet ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema  
 ■ cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use** if you are taking the blood thinning drug warfarin.

**When using this product, do not exceed recommended dosage.**

**Stop use and ask a doctor if** ■ nervousness, dizziness, or sleeplessness occur ■ pain, nasal congestion, or cough gets worse, or lasts more than 5 days (children) or 7 days (adult)  
 ■ fever gets worse, or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended

**Drug Facts (continued)**

dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ■ do not take more than directed (see Overdose warning) ■ do not take more than 4 doses in any 24-hour period ■ measure only with dosing cup provided. Do not use any other dosing device.  
 ■ keep dosing cup with product ■ mL = milliliter

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children 4 to under 6 years	ask a doctor
children under 4 years	do not use

■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

**Other information**

■ each 15 mL contains: sodium 12 mg  
 ■ store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients** citric acid, FD&C yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

**Questions or comments?**

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

## SIGNATURE CARE Non-Drowsy Daytime Cold & Flu Relief

### DAY TIME COLD AND FLU RELIEF NON DROWSY

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride liquid

**Product Information****Product Type**

HUMAN OTC DRUG

**Item Code (Source)**

NDC:21130-866

**Route of Administration**

ORAL

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH)	DEXTROMETHORPHAN	10 mg

(DEXTROMETHORPHAN - UNII:7355X3ROTS)	HYDROBROMIDE	in 15 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-866-12	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/31/2015	
2	NDC:21130-866-08	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/31/2015	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/31/2015	

**Labeler** - Safeway, Inc. (009137209)

Revised: 5/2024

Safeway, Inc.