

HISTEX- triprolidine hydrochloride syrup
Allegis Pharmaceuticals, LLC

HISTEX™ Syrup

Drug Facts

Active ingredients (in each 5 mL teaspoonful)

Tripolidine HCl 2.5 mg

Purpose

Antihistamine

Uses

temporarily relieves these symptoms due to hay fever (allergic rhinitis) or other upper respiratory allergies:

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes

Warnings

Do not exceed recommended dosage.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland

Ask a doctor before use if you are taking sedatives or tranquilizers

When using this product

- excitability may occur, especially in children
- may cause drowsiness
- alcohol, sedatives and tranquilizers may increase the drowsiness effect
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- new symptoms occur

If pregnant or breast-feeding, ask a health professional before use.

Keep out of the reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Do not exceed recommended dosage.

AGE	DOSE
Adults and Children 12 years of age and older:	1 teaspoonful (5 mL) every 4 to 6 hours, not to exceed 4 teaspoonfuls (20 mL) in 24 hours, or as directed by a doctor
Children 6 to under 12 years of age	$\frac{1}{2}$ teaspoonful (2.5 mL) every 4 to 6 hours, not to exceed 2 teaspoonfuls (10 mL) in 24 hours, or as directed by a doctor.
Children under 6 years of age	Consult a doctor

Other Information

Store at 15°-30° C (59°-86° F).

Tamper evident by foil seal under cap. Do not use if foil seal is missing or broken.

Dispense in a tight, light-resistant container with a child-resistant cap.

Inactive ingredients

Citric Acid, Glycerin, Propylene Glycol, Purified Water, Sodium Citrate, Sodium Saccharin, Sorbitol, Bubble Gum Flavor.

Questions? Comments?

Call 1-866-633-9033.

PRINCIPAL DISPLAY PANEL - 237 mL Bottle Label

NDC 28595-802-08

Antihistamine

HISTEX™

Syrup

Each teaspoonful (5 mL)

contains:

Tripolidine HCl 2.5 mg

Sugar-Free • Dye Free

Alcohol Free

Bubble Gum Flavor

8 fl oz (237 mL)

NDC 28595-802-08

Antihistamine

HISTEX™ Syrup

Each teaspoonful (5 mL) contains:
Tripolidine HCl 2.5 mg

Sugar-Free • Dye Free
Alcohol Free

Bubble Gum Flavor

8 fl oz (237 mL)

Drug Facts		Drug Facts (continued)
Active ingredient (in each 5 mL teaspoonful) Tripolidine HCl 2.5 mg	Purpose Antihistamine	Ask a doctor before use if you are taking sedatives or tranquilizers
Uses temporarily relieves these symptoms due to hay fever (allergic rhinitis) or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itching of the nose or throat ■ itchy, watery eyes		When using this product ■ excitability may occur, especially in children ■ may cause drowsiness ■ alcohol, sedatives and tranquilizers may increase the drowsiness effect ■ avoid alcoholic beverages ■ use caution when driving a motor vehicle or operating machinery
Warnings Do not exceed recommended dosage. Ask a doctor before use if you have ■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma ■ trouble urinating due to an enlarged prostate gland		Stop use and ask a doctor if ■ new symptoms occur If pregnant or breast-feeding, ask a health professional before use. Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)	Drug Facts (continued)								
Directions Do not exceed recommended dosage.	Tamper evident by foil seal under cap. Do not use if foil seal is missing or broken.								
<table border="1"><thead><tr><th>AGE</th><th>DOSE</th></tr></thead><tbody><tr><td>Adults and Children 12 years of age and older:</td><td>1 teaspoonful (5 mL) every 4 to 6 hours, not to exceed 4 teaspoonfuls (20 mL) in 24 hours, or as directed by a doctor</td></tr><tr><td>Children 6 to under 12 years of age</td><td>½ teaspoonful (2.5 mL) every 4 to 6 hours, not to exceed 2 teaspoonfuls (10 mL) in 24 hours, or as directed by a doctor.</td></tr><tr><td>Children under 6 years of age</td><td>Consult a doctor</td></tr></tbody></table>	AGE	DOSE	Adults and Children 12 years of age and older:	1 teaspoonful (5 mL) every 4 to 6 hours, not to exceed 4 teaspoonfuls (20 mL) in 24 hours, or as directed by a doctor	Children 6 to under 12 years of age	½ teaspoonful (2.5 mL) every 4 to 6 hours, not to exceed 2 teaspoonfuls (10 mL) in 24 hours, or as directed by a doctor.	Children under 6 years of age	Consult a doctor	Inactive ingredients Bubble gum flavor, citric acid, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin, sorbitol solution
AGE	DOSE								
Adults and Children 12 years of age and older:	1 teaspoonful (5 mL) every 4 to 6 hours, not to exceed 4 teaspoonfuls (20 mL) in 24 hours, or as directed by a doctor								
Children 6 to under 12 years of age	½ teaspoonful (2.5 mL) every 4 to 6 hours, not to exceed 2 teaspoonfuls (10 mL) in 24 hours, or as directed by a doctor.								
Children under 6 years of age	Consult a doctor								
Other information Store at 15°-30° C (59°-86° F).	Questions? Comments? Call 1-866-633-9033.								
	Manufactured for: Allegis Pharmaceuticals, LLC Canton, MS 39046 www.allegispharma.com Rev. 03/22								

HISTEX

triprolidine hydrochloride syrup

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:28595-802
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TRIPROLIDINE HYDROCHLORIDE (UNII: YAN7R5L890) (TRIPROLIDINE - UNII:2L8T9S52QM)	TRIPROLIDINE HYDROCHLORIDE	2.5 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM BENZOATE (UNII: OJ245FE5EU)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	BUBBLE GUM	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:28595-802-08	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/04/2014	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/04/2014	

Labeler - Allegis Pharmaceuticals, LLC (792272861)

