

**ASPIRIN- aspirin tablet, film coated**  
**Bi-Mart**

-----

***Active ingredient (in each tablet)***

**Aspirin 325 mg (NSAID)\***

**\*nonsteroidal anti-inflammatory drug**

***Purpose***

**Pain reliever/Fever reducer**

***Uses***

temporary relieves minor aches and pains due to:

- minor pain of arthritis
- headache
- muscle pain
- menstrual pain
- toothache
- temporarily reduces fever

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- asthma (wheezing)
- hives
- facial swelling
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have asthma
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

***Directions***

- drink a full glass of water with each dose
- **adults and children 12 years and over:** take 1 to 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours unless directed by a doctor
- **children under 12 years:** do not use unless directed by a doctor

## Other information

- store at 25°C (77°F); excursions permitted between 15-30°C (59-86°F)
- use by expiration date on package

## Inactive ingredients

Colloidal Silicon Dioxide, Corn Starch, Hydroxypropyl Methylcellulose, Microcrystalline Cellulose, Polyethylene Glycol, Polyvinyl Alcohol, Povidone, Talc, Titanium Dioxide

## Questions or comments?

1-844-912-4012

## Principal Display Panel

**BI-MART** NDC 37 835-577-25

COMPARE TO ACTIVE INGREDIENT IN  
GENUINE BAYER® ASPIRIN®

**ASPIRIN**  
(NSAID) 325 mg

PAIN RELIEVER/  
FEVER REDUCER

Actual Size **250** COATED TABLETS

**Drug Facts**

**Active ingredient (in each tablet)** Aspirin 325 mg (NSAID) \*nonsteroidal anti-inflammatory drug

**Uses** temporarily relieves: ■ minor pain of arthritis ■ headache ■ muscle pain ■ menstrual pain ■ toothache ■ temporarily reduces fever

**Warnings**  
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include: ■ asthma (wheezing) ■ hives ■ facial swelling ■ shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ take more or for a longer time than directed ■ have 3 or more alcoholic drinks every day while using this product

**Do not use** ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients

This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Genuine Bayer® Aspirin.

Included by:  
BI-MART  
201 Service Road  
Eugene, OR 97402  
971-25-EM 1525

0 171357 00175 0

**PEEL HERE FOR MORE DRUG FACTS**

**Drug Facts (continued)**

**Ask a doctor before use if**  
■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have asthma ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic

**Ask a doctor or pharmacist before use if you are taking a prescription drug for** ■ gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if**  
■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ ringing in the ears or a loss of hearing occurs ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions** ■ do not take more than directed ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor ■ children under 12 years: do not use unless directed by a doctor

**Other information**  
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package

**Inactive ingredients** Colloidal silicon dioxide, Corn starch, Hydroxypropyl methylcellulose, Microcrystalline cellulose, Polyethylene glycol, Polyvinyl alcohol, Povidone, Talc, Titanium dioxide.

**Questions or comments?** 1-844-912-4012

**STOP PEELING**

**ASPIRIN**

135700174 3

EXP. DATE: 04/2025  
LOT: 135700174  
NDC 37 835-577-01

**Drug Facts**

**Active ingredient (in each tablet)** Aspirin 325 mg (NSAID) \*nonsteroidal anti-inflammatory drug

**Uses** temporarily relieves: ■ minor pain of arthritis ■ headache ■ muscle pain ■ menstrual pain ■ toothache ■ temporarily reduces fever

**Warnings**  
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include: ■ asthma (wheezing) ■ hives ■ facial swelling ■ shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ take more or for a longer time than directed ■ have 3 or more alcoholic drinks every day while using this product

**Do not use** ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients

This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Genuine Bayer® Aspirin.

Included by:  
BI-MART  
201 Service Road  
Eugene, OR 97402  
971-25-EM 1525

0 171357 00175 0

**PEEL HERE FOR MORE DRUG FACTS**

<h2>Drug Facts (continued)</h2>	
<p><b>Do not use</b> ■ If you are allergic to aspirin or any other pain reliever/reducer          ■ If you have ever had an allergic reaction to this product or any of its ingredients</p>	
<p><b>Ask a doctor before you use it</b> ■ you are taking a diuretic          ■ stomach bleeding warning applies to you          ■ you have a history of stomach problems, such as heartburn          ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you have asthma</p>	
<p><b>Ask a doctor or pharmacist before you use if you are taking a prescription drug, for</b> ■ gout ■ diabetes ■ arthritis</p>	
<p><b>Stop use and ask a doctor if:</b> ■ you have an allergic reaction occurs.          Stop medical help right away.          ■ you experience any of the following signs of stomach bleeding:          ■ feel faint ■ vomit blood          ■ have bloody or black stools          ■ have stomach pain that does not get better          ■ ringing in the ears or a loss of hearing occurs          ■ stool gets worse or lasts more than 10 days          ■ liver gets worse or lasts more than 3 days          ■ redness or swelling is present ■ new symptoms occur          These could be signs of a serious condition.</p>	
<p><b>If pregnant or breast-feeding,</b> ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p>	
<p><b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.</p>	
<p><b>Directions</b> ■ do not take more than directed          ■ drink a full glass of water with each dose          ■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor          ■ children under 12 years: do not use unless directed by a doctor</p>	
<p><b>Other information</b> ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)          ■ use by expiration date on package</p>	
<p><b>Inactive Ingredients</b> Colloidal silicon dioxide; Corn starch; Hydroxyethylcellulose; Microcrystalline cellulose; Polyethylene glycol; Polyvinyl alcohol; Povidone; Talc; Titanium dioxide.</p>	
<p><b>Questions or comments? 1-844-912-4012</b></p>	
<p><b>STOP FEELING</b></p>	

## aspirin tablet, film coated

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Ingredient Name	Strength
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL</b> (UNII: 532B59J990)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	

<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)				
<b>Product Characteristics</b>				
<b>Color</b>	white	<b>Score</b>	no score	
<b>Shape</b>	ROUND	<b>Size</b>	10mm	
<b>Flavor</b>		<b>Imprint Code</b>	ASP325	
<b>Contains</b>				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:37835-577-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	08/01/2025	
2	NDC:37835-577-25	250 in 1 BOTTLE; Type 0: Not a Combination Product	08/01/2025	
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>		<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M013		08/01/2025	

**Labeler** - Bi-Mart (027630078)

**Registrant** - Bi-Mart (027630078)

Establishment			
Name	Address	ID/FEI	Business Operations
Elysium Pharmaceuticals Ltd		915664486	manufacture(37835-577)