

PENNITRACIN MD 50G- bacitracin methylenedisalicylate granule
Pharmgate Animal Health

Pennitracin MD 50 G[®]
(Bacitracin Type A medicated article)

For further manufacturing only.

See mixing directions, claims, cautions and warnings on back panel.

Restricted Drug (California) - Use Only as Directed.

Net wt. 50 LB (22.68 Kg)

Pennitracin MD 50 G[®]

(Bacitracin Type A medicated article)

Active Drug Ingredients: Each pound contains feed grade bacitracin methylenedisalicylate equivalent to 50 grams bacitracin			
Inactive Ingredients: Calcium Carbonate, Mineral Oil			
CAUTION: For Use in Dry Feeds only. Not for use in liquid medicated feeds.			
Directions For Use Approved concentrations of bacitracin in Type C medicated feeds.			
Species	Grams of bacitracin per ton	Indications For Use	Feeding Directions
Broiler and replacement chickens, growing turkeys, growing pheasants	4-50	For increased rate of weight gain and improved feed efficiency.	Feed as the sole ration throughout the feeding period.
Broiler and replacement chickens	50	For the prevention of mortality caused by necrotic enteritis associated with <i>Clostridium perfringens</i> .	Feed as the sole ration for 28 to 35 days, starting from the time chicks are placed for brooding.
Growing Quail	5-20	For increased rate of weight gain and improved feed efficiency in quail not over 5 weeks of age.	Feed as the sole ration through 5 weeks of age.
Beef steers and heifers fed in confinement for slaughter	70 mg/head/day	For reduction in the number of liver condemnations due to abscesses.	Feed as the sole ration throughout the feeding period.
	250 mg/head/day	For reduction in the number of liver condemnations due to abscesses.	Feed as the sole ration for 5 days then discontinue for subsequent 25 days, repeat the pattern during the feeding period.

.Mixing Directions

Prepare an intermediate preblend containing 5 grams of bacitracin per pound by mixing 1.0 lb of Pennitracin MD 50G with 9.0 lbs of unmedicated feed ingredient(s), then add 0.8 to 10 lbs of the intermediate preblend per ton of the finished feed to obtain the appropriate g/ton concentration of the Type C medicated feed.

Prepare Type B medicated feeds for chickens, turkeys, pheasants, or beef cattle containing 51 - 50,000 grams of bacitracin per ton by mixing 1.02 - 1,000 lbs of Pennitracin MD 50G with 1,998.98 - 1,000 lbs of unmedicated feed.

CAUTION: For Use in Dry Feeds only. Not for use in liquid medicated feeds.

WARNINGS

Withdrawal Periods:

No withdrawal period is required when used according to labeling.

User Safety Warnings:

Not for use in humans. Keep out of reach of children.

Store at or below 25°C (77°F)

Lot Number and Expiration Date: Both the Lot number and expiry date is provided on the front panel of the bag.

QUESTIONS/COMMENTS? Contact Pharmgate Animal Health LLC at 1-800-380-6099 or www.pharmgate.com. To report side effects, contact Pharmgate Animal Health LLC at 1-833-531-0114. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

Approved by FDA under NADA # 141-137

Pennitracin MD 50G® is a registered trademark of Pharmgate Inc.

Marketed by:

PHARMGATE ANIMAL HEALTH LLC

14040 Industrial Road

Omaha, NE 68144



PRINCIPAL DISPLAY PANEL - 22.68 Kg Bag

Pennitracin MD 50G®

(Bacitracin Type A medicated article)

For further manufacturing only.

See mixing directions, claims, cautions and warnings on back panel.

Restricted Drug (California) - Use Only as Directed.


Net wt. 50 LB (22.68 Kg)

Pharmgate

ANIMAL HEALTH

Pennitracin MD 50 G

(Bacitracin Type A medicated article)



For further manufacturing only.

See mixing directions, claims, cautions
and warnings on back panel.

Restricted Drug (California) - Use Only as Directed.

Net wt. 50 LB (22.68 Kg)



Pharmgate
ANIMAL HEALTH
Pennitracin MD 50 G

Pennitracin MD 50G[®]

(Bacitracin Type A medicated article)



Pharmgate
ANIMAL HEALTH

Pennitracin MD 50G[®]

Active Drug Ingredients	Each pound contains feed grade bacitracin methylglucoside tetrahydrate (Type C) bacitracin		
Inert Ingredients	Calcium Carbonate, Milled Cell		
Directions For Use			
Approved concentrations of bacitracin in Type C medicated feeds			
Species	Approved Feed Rate (ppm)	Indications For Use	Feeding Directions
Beef and replacement calves, growing heifers, growing steers	4.0	For improved rate of weight gain and improved feed efficiency	Feed as directed throughout the feeding period.
Beef and replacement calves	30	For the prevention of scours caused by specific bacteria associated with Clostridium perfringens.	Feed as directed for 21 to 28 days, starting from the time the calf is placed for weaning.
Growing Calf	5.0	For improved rate of weight gain and improved feed efficiency in quality lot over 2 weeks of age.	Feed as directed through 2 weeks of age.
Dry cows and heifers fed in confinement for slaughter	15 mg / head / day	For reduction of the number of live coccidia oocysts shed in manure.	Feed as directed throughout the feeding period.
	20 mg / head / day	For reduction of the number of live coccidia oocysts shed in manure.	Feed as directed for 14 days from the time the cow begins to dry, or for the entire dry period during the breeding period.
Mixing Directions			
Prepare an amount of feed containing 2 grams of bacitracin per pound by using 10 lbs of Pennitracin MD 50G (with 1 lb of unmedicated feed bags of 20 lbs) from 40 lbs of the same size pack (or per manufacturer's label) to distribute appropriately to the composition of the Type C medicated feed.			
Prepare Type B medicated feed (with 20 mg of bacitracin per lb) by using 1 lb of Pennitracin MD 50G (with 1 lb of unmedicated feed bags of 20 lbs) from 40 lbs of the same size pack (or per manufacturer's label) to distribute appropriately to the composition of the Type C medicated feed.			
CAUTION: For use in Dry Feeds only		Not for use in liquid medicated feeds.	
WARNINGS			
With drawal Periods:			
No withdrawal period is required when used as directed in the bag.			
User Safety Warnings:			
Not for use in human food products or feed.			
Store at below 77°F (25°C).			
Lot Number and Expiration Date: Both the Lot number and expiry date are provided on the front panel of the bag.			
CONTRIBUTOR COMMENTS: Contact Pharmgate Animal Health, LLC at 1-800-880-4999 or www.pharmgate.com. To report side effects, contact Pharmgate Animal Health, LLC at 1-800-880-4999. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VEED or http://www.fda.gov/reportmedwatch.			
Approved by FDA under NADA 141 H 419 Pennitracin MD 50G [®] is a registered trademark of Pharmgate, Inc.			



Marketed by:
Pharmgate Animal Health LLC
14040 Industrial Road
Omaha, Nebraska 68144



21180 3-03
Rev 1/22
Bag MD



22FG-GVC-00014B

GVC-14

PENNITRACIN MD 50G

bacitracin methylenedisalicylate granule

Product Information

Product Type	OTC TYPE A MEDICATED ARTICLE ANIMAL DRUG	Item Code (Source)	NDC:51429-054
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Bacitracin Methylenedisalicylate (UNII: JGV6OJ52FT) (BACITRACIN - UNII:58H6RWO52I)	Bacitracin Methylenedisalicylate	110 g in 1 kg

Inactive Ingredients

Ingredient Name	Strength
Calcium Carbonate (UNII: H0G9379FGK)	
Mineral Oil (UNII: T5L8T28FGP)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51429-054-05	22.68 kg in 1 BAG		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA141137	08/01/2018	

Labeler - Pharmgate Animal Health (833270817)**Registrant** - Pharmgate Inc. (079628671)