

**NIGHTTIME COLD AND FLU RELIEF- acetaminophen, dextromethorphan hbr and doxylamine succinate capsule, liquid filled
SALADO SALES INC.**

CVP Health Night Time Cold & Flu Relief liquid filled softgels

Active ingredients (in each Softgel)

Acetaminophen 325 mg
Dextromethorphan HBr 15 mg
Doxylamine succinate 6.25 mg

Purpose

Pain reliever/Fever reducer
Cough suppressant
Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever
- runny nose & sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 liquid caps 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleep

Ask a doctor before use if you have

- liver disease ■ glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, & tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center right away.

Quick medical attention is critical for adults as well as for children, even if you do not notice any signs or symptoms.

Directions

- take only as directed (see Warnings)
- do not exceed 4 doses per 24 hrs

| | |
|---------------------------------|-----------------------------------|
| adults & children 12 yrs & over | 2 softgels with water every 6 hrs |
| children 4 to under 12 yrs | ask a doctor |
| children under 4 yrs | do not use |

■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Other information

- Store at room temperature 20-25°C (68-77°F)
- Avoid excessive heat

Inactive ingredients

D&C yellow no. 10, FD&C blue no. 1, Gelatin, Glycerin, Polyethylene glycol, povidone, Propylene glycol, Purified water, Sorbitol sorbitan solution, Titanium dioxide

Questions?

Call weekdays from 9:30 AM to 4:30 PM EST at **1-877-798-5944**

CVP

COMPARE TO VICKS® NYQUIL™

HEALTH

COLD & FLU LIQUICAPS™

NIGHT TIME

ACTIVE INGREDIENTS

SOFTGELS

MULTI-SYMPOTOM COLD & FLU MEDICINE

RELIEVES:

- **ACETAMINOPHEN-PAIN RELIEVER/FEVER REDUCER**
HEADACHE

- **DEXTROMETHORPHAN HBr- COUGH SUPPRESSANT**
FEVER
- **DOXYLAMINE SUCCINATE- ANTIHISTAMINE**
SORE THROAT
- MINOR ACHES & PAINS

24 SOFTGELS
SNEEZING

NOSE

RUNNY

COUGH

Distributed by Consumer Value Products, Inc.

P.O. Box 6115, Temple, Texas 76502

CVPProducts.com

*This product is not manufactured or distributed by Procter & Gamble, owner of the registered trademark Vicks® NyQuil™ LiquiCaps™

6
1201240
3/22



NIGHT TIME SOFTGELS

MULTI-SYMPOTOM COLD & FLU MEDICINE

COMPARE TO
VICKS NYQUIL™
COLD & FLU LIQUICAPS™
ACTIVE INGREDIENTS



HEALTH

NIGHT TIME SOFTGELS

MULTI-SYMPOTOM COLD & FLU MEDICINE

- ACETAMINOPHEN - PAIN RELIEVER/FEVER REDUCER
- DEXTROMETHORPHAN HBr - COUGH SUPPRESSANT
- DOXYLAMINE SUCCINATE - ANTIHISTAMINE



ACTUAL
SIZE

24 SOFTGELS

RELIEVES:
HEADACHE
FEVER
SORE THROAT
MINOR ACHES & PAINS
SNEEZING
RUNNY NOSE
COUGH

Distributed by Consumer Value Products, Inc.
P.O. Box 6115, Temple, Texas 76502
CVPproducts.com

TAMPER EVIDENT: DO NOT USE IF
PACKAGE IS OPENED OR BLISTER
UNIT IS TORN, BROKEN OR SHOWS
ANY SIGNS OF TAMPERING

BX-137

Drug Facts

Active Ingredients (in each softgel) Purpose

Acetaminophen 325 mg.....Pain reliever/fever reducer
Dextromethorphan HBr 15 mg.....Cough suppressant
Doxylamine succinate 6.25 mg.....Antihistamine

Uses

temporarily relieves common cold/flu symptoms:
■ cough due to minor throat & bronchial irritation
■ sore throat ■ headache ■ minor aches & pains ■ fever ➤

■ runny nose & sneezing

Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

■ more than 8 Liquid Caps in 24 hrs, which is the maximum daily amount for this product ■ with other drugs containing acetaminophen

■ 3 or more alcoholic drinks daily while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

■ skin reddening ■ blisters ■ rash

Drug Facts (continued)

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

■ to make a child sleep

Ask a doctor before use if you have ■ liver disease ■ glaucoma

■ cough that occurs with too much phlegm (mucus)
■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
■ trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

■ taking sedatives or tranquilizers
■ taking the blood thinning drug warfarin

When using this product, do not use more than directed.

■ excitability may occur, especially in children
■ marked drowsiness may occur ■ avoid alcoholic drinks
■ be careful when driving a motor vehicle or operating machinery
■ alcohol, sedatives & tranquilizers may increase drowsiness

Stop use and ask a doctor if

■ pain or cough gets worse or lasts more than 7 days
■ fever gets worse or lasts more than 3 days

Drug Facts (continued)

■ redness or swelling is present ■ new symptoms occur
■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as children, even if you do not notice any signs or symptoms.

Directions

■ take only as directed (see Warnings)
■ do not exceed 4 doses per 24 hrs

| | |
|---------------------------------|------------------------------------|
| adults & children 12 yrs & over | 2 softgel with water every 6 hours |
|---------------------------------|------------------------------------|

| | |
|----------------------------|--------------|
| children 4 to under 12 yrs | ask a doctor |
|----------------------------|--------------|

| | |
|----------------------|------------|
| children under 1 yrs | do not use |
|----------------------|------------|

■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Other information

■ Store at room temperature 20-25°C (68-77°F)
■ avoid excessive heat

Inactive ingredients D&C Yellow # 10, FD&C Blue # 1, Gelatin, Glycerin, Polyethylene glycol, Povidone, Propylene glycol, Purified water, Sorbitol sorbitan solution, Titanium dioxide

Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944

*This product is not manufactured or distributed by Procter & Gamble, owner of the registered trademark Vicks® NyQuil™ LiquiCaps™

RES

NIGHTTIME COLD AND FLU RELIEF

acetaminophen, dextromethorphan hbr and doxylamine succinate capsule, liquid filled

Product Information

| | | | |
|-------------------------|----------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:57243-603 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------------------|----------|
| ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D) | ACETAMINOPHEN | 325 mg |
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 15 mg |
| DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL) | DOXYLAMINE SUCCINATE | 6.25 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|--------------------------------------|----------|
| D&C YELLOW NO. 10 (UNII: 35SW5USQ3G) | |

FD&C BLUE NO. 1 (UNII: H3R47K3TBD)

GELATIN (UNII: 2G86QN327L)

GLYCERIN (UNII: PDC6A3C00X)

POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)

POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)

WATER (UNII: 059QF0KO0R)

SORBITOL (UNII: 506T60A25R)

SORBITAN (UNII: 6092ICV9RU)

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

Product Characteristics

| | | | |
|-----------------|-------|---------------------|----------|
| Color | green | Score | no score |
| Shape | OVAL | Size | 20mm |
| Flavor | | Imprint Code | 603 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:57243-603-03 | 24 in 1 CARTON | 12/23/2025 | |
| 1 | | 12 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M012 | 12/23/2025 | |

Labeler - SALADO SALES INC. (009830555)

Revised: 12/2025

SALADO SALES INC.