

**LOW STRENGTH CHEWABLE ASPIRIN- aspirin tablet, chewable**  
**Geri-Care Pharmaceutical Corp**

-----  
**gc 911**

**Active ingredient (in each tablet)**

Aspirin 81 mg(NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever

**Uses**

for the temporary relief of minor aches and pains or as recommended by your doctor

**Warnings**

**Reye's Syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy Alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if your are allergic to aspirin or any other pain reliever/fever reducer.
- this product for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

### **Ask a doctor or pharmacist before use if you are you**

are taking a prescription drug for

gout  diabetes  arthritis

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding: -feel faint -vomit blood -have bloody or black stools -have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- fever gets worse or lasts more than 3 days
- any new symptoms occur
- ringing in the ears or loss of hearing occurs

### **If pregnant or breast-feeding,**

ask a health professional before use. it is especially important not to use aspirin at 20 weeks or later in

pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose , get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### **Directions**

- drink a full glass of water with each dose
- do not exceed recommended dose
- adults and children 12 years and older: take 4-8 tablets every 4 hours, as needed, not more than 48 tablets in 24 hours, or as directed by a doctor
- children under 12 years: ask a doctor

### **Other information**

- store at room temperature 15°C - 30°C (59°F - 86°F)

### **Inactive ingredients**

corn starch, dextrans, FD&C yellow #6 aluminum lake, orange flavor, saccharin sodium.

Questions or comments?

1-800-540-3765

Package label

**GERI-CARE**  
NDC 57896-911-36

**ADULT LOW DOSE CHEWABLE ASPIRIN 81 mg**  
PAIN RELIEVER (NSAID)

Compare to active ingredient in **BAYER® Low Dose Aspirin\***

**36 Tablets**

**Drug Facts**  
Tamper Evident: Do not use if imprinted seal under cap is missing or broken.  
**Active ingredient (in each tablet)** Aspirin 81 mg (NSAID)\*  
\*nonsteroidal anti-inflammatory drug  
**Purpose** Pain reliever

**Uses** for the temporary relief of minor aches and pains or as recommended by your doctor

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior or with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:  

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinner (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use** if you are allergic to aspirin or any other pain reliever/fever reducer  
 ■ this product for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor  
 ■ if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**  
 ■ stomach bleeding warning applies to you  
 ■ you have a history of stomach problems, such as heartburn  
 ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease  
 ■ you are taking a diuretic  
 ■ you have asthma

\* This product is not manufactured or distributed by the owner of the registered trademark Bayer®.  
Dist. by: Geri-Care Pharmaceuticals Corp.  
1295 Towbin Avenue, Lakewood, NJ 08701

REV 1123  
3 57896 91136 2  
PEEL HERE FOR MORE DRUG FACTS

**Drug Facts (continued)**

**Ask a doctor or pharmacist before use if you are taking a prescription drug for**  
 ■ gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if**  
 ■ an allergic reaction occurs. Seek medical help right away if you experience any of the following signs of stomach bleeding:  
 ■ feel faint ■ vomit blood ■ have bloody or black stool  
 ■ pain gets worse or lasts more than 10 days  
 ■ redness or swelling is present  
 ■ new symptoms occur  
 ■ ringing in the ears or loss of hearing occurs

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center. (1-800-222-1222) right away.

**Directions**  
 ■ drink a full glass of water with each dose  
 ■ do not exceed recommended dose  
 adults and children take 4-8 tablets every 4-12 hours, as needed, not more than 48 tablets in 24 hours, or as directed by a doctor  
 children under 12 ask a doctor

**Other information** ■ store at room temperature 15°C - 30°C (59°F - 86°F)

**Inactive ingredients:** corn starch, dextrates, FD&C yellow #6 aluminum lake, orange flavor, saccharin sodium.

**Questions or comments? 1-800-540-3765**

**STOP PEELING**

**LOW STRENGTH CHEWABLE ASPIRIN**

aspirin tablet, chewable

**Product Information**

|                                |                |                           |               |
|--------------------------------|----------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:57896-911 |
| <b>Route of Administration</b> | ORAL           |                           |               |

**Active Ingredient/Active Moiety**

| Ingredient Name  | Basis of Strength | Strength |
|--|-------------------|----------|
| ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E) | ASPIRIN           | 81 mg    |

## Inactive Ingredients

| Ingredient Name                      | Strength |
|--------------------------------------|----------|
| FD&C YELLOW NO. 6 (UNII: H77VEI93A8) |          |
| SACCHARIN SODIUM (UNII: SB8ZUX40TY)  |          |
| DEXTRATES (UNII: G263MI44RU)         |          |
| STARCH, CORN (UNII: O8232NY3SJ)      |          |

## Product Characteristics

|          |        |              |          |
|----------|--------|--------------|----------|
| Color    | orange | Score        | no score |
| Shape    | ROUND  | Size         | 7mm      |
| Flavor   | ORANGE | Imprint Code | AZ;013   |
| Contains |        |              |          |

## Packaging

| # | Item Code        | Package Description                               | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:57896-911-36 | 36 in 1 BOTTLE; Type 0: Not a Combination Product | 01/01/2013           |                    |

## Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M013                                     | 01/01/1995           |                    |

**Labeler** - Geri-Care Pharmaceutical Corp (611196254)

**Registrant** - Geri-Care Pharmaceutical Corp (611196254)