

GOOD SENSE ACID REDUCER- famotidine tablet, film coated
L. Perrigo Company

Perrigo Acid Reducer Drug Facts

Active ingredient (in each tablet)

Famotidine 10 mg

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
- to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **15 to 60 minutes before** eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture

Inactive ingredients

carnauba wax, hypromellose, iron oxide red, iron oxide yellow, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, pregelatinized starch, talc, titanium dioxide, triacetin

Questions or comments?

1-800-719-9260

Principal Display Panel

Original Strength

See New Warnings

Famotidine Tablets, 10mg

Acid Reducer

Actual Size

Just One Tablet!

-
Prevents & Relieves Heartburn Due to Acid Indigestion

Compare to active ingredient of Pepcid® AC

100% SATISFACTION GUARANTEED

30 Tablets

GOODSENSE®

Famotidine Tablets, 10 mg

Acid Reducer

DO NOT USE IF BLISTER
UNIT IS BROKEN OR TORN

JUST ONE TABLET prevents
and relieves heartburn due to
acid indigestion brought on by
eating and drinking certain
foods and beverages.

GOODSENSE®

NDC 0113-0141-65

Original Strength

See New Warnings

Famotidine Tablets, 10 mg

Acid Reducer

Actual
Size

Just One Tablet!

**Prevents & Relieves Heartburn
Due to Acid Indigestion**

Compare to active ingredient of
Pepcid® AC

100%
SATISFACTION
GUARANTEED

30 Tablets

14165 C2 C2

OPEN OTHER END

Drug Facts

Active ingredient (in each tablet) Purpose
Famotidine 10 mg.....Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain ■ kidney disease

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
 - to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
 - to prevent symptoms, swallow 1 tablet with a glass of water at any time from 15 to 60 minutes before eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F) ■ protect from moisture

Inactive ingredients camellia wax, hypromellose, iron oxide red, iron oxide yellow, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, pregelatinized starch, talc, titanium dioxide, triacetin

Tips for Managing Heartburn • Do not lie flat or bend over after eating • Do not wear tight-fitting clothing around the stomach • Do not eat before bedtime • Raise the head of your bed • Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables • Eat slowly and avoid big meals • If overweight, lose weight • Quit smoking

CONVENIENT RECLOSING TAB



GOOD SENSE ACID REDUCER

famotidine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0113-0141
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	10 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	

Product Characteristics

Color	PINK	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	L141
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0113-0141-65	30 in 1 CARTON	10/23/2009	

1	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075400	10/23/2009	

Labeler - L. Perrigo Company (006013346)

Revised: 2/2025

L. Perrigo Company