

**MCKESSON ORASOL GEL- benzocaine 20% gel**  
**McKesson Medical-Surgical**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**McKesson Orasol Gel**

**Drug Facts**

**Active ingredient**

Benzocaine 20% (w/w)

**Purpose**

Oral anesthetic

**Uses**

For the temporary relief of minor pain and sore mouth associated with toothache, minor dental procedures and irritations from dentures or orthodontic appliances.

**Warnings**

**For oral use only.**

**Do not exceed recommended dosage.**

**Methemoglobinemia warning:**

Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before.

Stop use and seek medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- rapid heart rate
- dizziness or lightheadedness
- headache
- shortness of breath
- fatigue or lack of energy

**Allergy alert:**

Do not use this product if you have a history of allergy to local anesthetics, such as procaine, butacaine, benzocaine, or other "caine" anesthetics. If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- for teething - any teething related indication will not be an appropriate use of this product
- in children under 2 years of age
- for more than 7 days unless directed by a doctor

**Stop use and ask a dentist or doctor if**

- sore mouth does not improve in 7 days
- irritation, pain, or redness persists or worsens
- swelling, rash, fever or other allergic reaction develops

**When using this product**

- avoid contact with eyes

**Keep out of reach of children.**

If accidentally swallowed get medical help or contact a Poison Control Center right away.

**Directions****Adults and children (2 years and older):**

Dry affected area and apply medication undiluted. Use up to 4 times daily, but not more than every two hours, or as directed by a dentist or doctor.

**Children under 2 years:**

Do not use

**Other information**

- do not use if packet is torn, cut or opened
- store at room temperature: 59° to 86°F (15° to 30°C)
- tamper evident sealed packets

**Inactive ingredients**

PEG 400, PEG 3350, peppermint oil, sodium saccharin, sorbic acid

**Questions or comments?** Call 1-800-777-4908

NDC 68599-8247-4

McKesson

Orasol Gel

Individual Packets

Benzocaine 20% | Oral Anesthetic

Temporary relief of minor pain and sore mouth.

NET WT 0.75 g

75 Packs Per Box

Do Not Reuse

MFR # 82474

**Drug Facts**

**Active ingredient (in each gram)**  
Benzocaine 20% (w/w)      **Purpose**  
Oral anesthetic

**Uses**  
For the temporary relief of minor pain and sore mouth associated with toothache, minor dental procedures and irritations from dentures or orthodontic appliances.

**Warnings**  
For oral use only.  
Do not exceed recommended dosage.

**Methemoglobinemia warning:** Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before.  
Stop use and seek immediate medical attention if you or a child in your care develops:  

- pale, gray, or blue colored skin (cyanosis)
- rapid heart rate
- dizziness or lightheadedness
- headache
- shortness of breath
- fatigue or lack of energy

**Allergy alert:** Do not use this product if you have a history of allergy to local anesthetics such as procaine, tetracaine, benzocaine, or other "caine" anesthetics. If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**  

- for teething — any teething related indication will not be an appropriate use for this product
- in children under 2 years of age
- for more than 7 days unless directed by dentist or doctor

**Stop use and ask a dentist or doctor if**  

- sore mouth does not improve in 7 days
- irritation, pain, or redness persists or worsens
- swelling, rash, hives or other allergic reaction develops

**When using this product**  

- avoid contact with eyes

**Drug Facts (continued)**

Keep out of the reach of children.  
If accidentally swallowed get medical help or contact a Poison Control Center right away.

**Directions**

**Adults and children: (2 years and older)** Dry affected area and apply medication undiluted. Use up to 4 times daily, but not more than every 2 hours or as directed by a dentist or doctor.

**Children under 2 years:** Do not use.

**Other information**  

- do not use if packet is torn, cut or opened
- store at room temperature: 59°-86°F (15°-30°C)
- tamper evident sealed packets

**Inactive ingredients**  
PEG 400, PEG 3350, peppermint oil, sodium saccharin, sorbic acid

**Questions or comments?** Call 1-800-777-4908

NDC 68599-8247-4

**MCKESSON**

**Orasol Gel**

INDIVIDUAL PACKETS  
BENZOCAINE 20% | ORAL ANESTHETIC

Temporary relief of minor pain and sore mouth.

NET WT **0.75 g**    **75** PACKS PER BOX    DO NOT REUSE    MFR # **82474**

NDC 68599-8247-4

**MCKESSON**

**Orasol Gel**

INDIVIDUAL PACKETS  
BENZOCAINE 20% | ORAL ANESTHETIC

Temporary relief of minor pain and sore mouth.

NET WT **0.75 g**    **75** PACKS PER BOX    DO NOT REUSE    MFR # **82474**

**Satisfaction Guaranteed**  
For complete details, please visit [mms.mckesson.com/mckesson-brands](http://mms.mckesson.com/mckesson-brands).

Distributed by  
McKesson Medical-Surgical Inc.  
Richmond, VA 23233  
PVN A0721  
Made in USA

MFR #  
**82474**

6 12479 25358 2

# MCKESSON ORASOL GEL

benzocaine 20% gel

Product Information			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:68599-8247
<b>Route of Administration</b>	TOPICAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
<b>BENZOCAINE</b> (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	200 mg in 1 g

Inactive Ingredients	
Ingredient Name	Strength
<b>POLYETHYLENE GLYCOL 3350</b> (UNII: G2M7P15E5P)	
<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)	
<b>PEPPERMINT OIL</b> (UNII: AV092KU4JH)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SORBIC ACID</b> (UNII: X045WJ989B)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68599-8247-4	75 in 1 BOX	10/04/2021	
1		0.75 g in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part356	10/04/2021	

**Labeler** - McKesson Medical-Surgical (023904428)

**Registrant** - Unifirst First Aid Corporation (832947092)

Revised: 10/2022

McKesson Medical-Surgical