

LIL DRUG STORE IBUPROFEN- ibuprofen tablet, coated
Lil Drug Store Products, Inc

Lil' Drug Store® Ibuprofen

Drug Facts

Active ingredient (in each tablet)

Ibuprofen USP, 200 mg (NSAID)¹

¹ nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - slurred speech
 - trouble breathing
 - leg swelling
 - weakness in one part or side of the body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and older:
 - take 1 tablet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 tablet, 2 tablets may be used
 - do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store between 20°-25°C (68°-77°F)

Inactive ingredients

carnauba wax, croscarmellose sodium, hypromellose, microcrystalline cellulose, polydextrose, polyethylene glycol, pregelatinized starch, red iron oxide, silicon dioxide, stearic acid, titanium dioxide

Questions or comments?

Call toll-free **1-877-507-6516 (M-F 8AM-4:30PM CST)**

Product distributed by:

Lil' Drug Store Products, Inc.
1201 Continental Place NE
Cedar Rapids, IA 52402

PRINCIPAL DISPLAY PANEL - 200 mg Tablet Bottle Carton

**Compare to the
Active Ingredient
in Advil® Tablets****

QUALITY

□ **GUARANTEED** □

**Ibuprofen
Tablets USP, 200 mg
Pain Reliever/Fever Reducer (NSAID)**

ACTUAL SIZE

50

**Coated
Tablets**

**Lil'
DrugStore®**

Ibuprofen

Tablets USP, 200 mg

Pain Reliever/Fever Reducer (NSAID)

50
Coated
Tablets



Compare to the
Active Ingredient
in Advil® Tablets**

Ibuprofen

Tablets USP, 200 mg

Pain Reliever/Fever Reducer (NSAID)



ACTUAL SIZE

50
Coated
Tablets



Product distributed by:
Lil' Drug Store Products, Inc.
1201 Continental Place NE
Cedar Rapids, IA 52402

PLD-A16XX
FC002499



3 66715 68204 8

Lot No:
Exp Date:

TAMPER-EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF
PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE
WARNINGS AND PRODUCT INFORMATION.

Drug Facts

Active ingredient (in each tablet) Purpose

Ibuprofen USP, Pain reliever/
200 mg (NSAID)*fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

- The chance is higher if you:
- are age 60 or older
 - have had stomach ulcers or bleeding problems
 - take a blood thinning (anticoagulant) or steroid drug
 - take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 - have 3 or more alcoholic drinks every day while using this product
 - take more or for a longer time than directed

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are


- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
- you have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - slurred speech
 - trouble breathing
 - leg swelling
 - weakness in one part or side of the body
- pain gets worse or lasts more than 10 days



Drug Facts (continued)

- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions • do not take more than directed

- the smallest effective dose should be used
- adults and children 12 years and older:
 - take 1 tablet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 tablet, 2 tablets may be used
 - do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store between 20°-25°C (68°-77°F)

Inactive ingredients carnauba wax, croscarmellose sodium, hypromellose, microcrystalline cellulose, polydextrose, polyethylene glycol, pregelatinized starch, red iron oxide, silicon dioxide, stearic acid, titanium dioxide

Questions or comments? Call toll-free 1-877-507-6516 (M-F 8AM-4:30PM CST)

**This product is not manufactured or distributed by Pfizer Consumer Healthcare, owner of the registered trademark Advil® Tablets.

68204C-US-02-17

LIL DRUG STORE IBUPROFEN

ibuprofen tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66715-6820(NDC:59726-235)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength		
Ibuprofen (UNII: WK2XYI10QM) (Ibuprofen - UNII:WK2XYI10QM)	Ibuprofen	200 mg		
Inactive Ingredients				
Ingredient Name	Strength			
CARNAUBA WAX (UNII: R12CBM0EIZ)				
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)				
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)				
Microcrystalline Cellulose (UNII: OP1R32D61U)				
Polydextrose (UNII: VH2XOU12IE)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
STARCH, CORN (UNII: O8232NY3SJ)				
FERRIC OXIDE RED (UNII: 1K09F3G675)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
Stearic Acid (UNII: 4ELV7Z65AP)				
Titanium Dioxide (UNII: 15FIX9V2JP)				
Product Characteristics				
Color	BROWN	Score	no score	
Shape	ROUND	Size	9mm	
Flavor		Imprint Code	57	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-6820-4	1 in 1 CARTON	02/23/2017	12/01/2021
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA076460	02/23/2017	12/01/2021	

Labeler - Lil Drug Store Products, Inc (093103646)

Revised: 12/2018

Lil Drug Store Products, Inc