TOPCARE NITE TIME COUGH- dextromethorphan hydrobromide, doxylamine succinate solution Topco Associates LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Topco Associates LLC. Nite Time Drug Facts

Active ingredients (in each 30 mL)

Dextromethorphan HBr 30 mg
Doxylamine succinate 12.5 mg

Purpose

Cough suppressant

Antihistamine

Uses

temporarily relieves cold symptoms:

- · cough due to minor throat and bronchial irritation
- runny nose and sneezing

Warnings

Do not use

 if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- · persistent or chronic cough as occurs with smoking, asthma, or emphysema
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

When using this product

- excitability may occur, especially in children
- may cause marked drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and as a doctor if

cough persists for more than 1 week, tends to recur, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 30 mL contains: sodium 32 mg
- store at 20-25°C (68-77°F)

Inactive ingredients

alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments?

1-888-423-0139

Principal Display Panel

TopCare_® health

COMPARE TO VICKS® NYQUIL® COUGH ACTIVE INGREDIENTS

ALL NIGHT COUGH RELIEF

Nite Time Cough

COUGH SUPPRESSANT - DEXTROMETHORPHAN HBr

ANTIHISTAMINE - DOXYLAMINE SUCCINATE


- Cough
- Sneezing
- Runny Nose

ALCOHOL 10%

12 FL OZ (355 mL)

CHERRY FLAVOR





ALCOHOL 10%

12 FL OZ (355 mL)

CHERRY FI AVOR

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

Drug Facts Active ingredients Purpose (in each 30 mL)

. Dextromethorphan HBr 30 mg.. Doxylamine succinate 12.5 mg...... Antihistamine

Uses temporarily relieves cold symptoms: ■ cough due to minor throat and bronchial irritation

runny nose and sneezing

Warnings

Do not use

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have glaucoma

- cough that occurs with too much phlegm (mucus) a breathing problem such as emphysema or chronic bronchitis
 persistent or chronic cough as occurs with smoking, asthma,
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking edatives or tranquilizers

When using this product

- excitability may occur, especially in children
- may cause marked drowsiness
 avoid alcoholic drinks
 be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsines

Stop use and ask a doctor if cough persists for more than 1 week, tends to recur, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

: 66840 88 F6

TOPCARE NITE TIME COUGH

dextromethorphan hydrobromide, doxylamine succinate solution

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:36800-668

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL	
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL	

Inactive Ingredients		
Ingredient Name	Strength	
ALCOHOL (UNII: 3K9958V90M)		
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)		
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)		
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3MJQ0SDW1A)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
WATER (UNII: 059QF0KO0R)		
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)		
SACCHARIN SODIUM (UNII: SB8ZUX40TY)		

Product Characteristics		
Color	RED (Dark Red)	Score
Shape		Size
Flavor	CHERRY	Imprint Code
Contains		

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-668- 38	296 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/03/2004	11/02/2011
2	NDC:36800-668- 40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/16/2012	

Marketing In	rketing Information		
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	12/10/2002	

Revised: 9/2022 Topco Associates LLC