## STAHIST AD- chlorcyclizine hydrochloride and pseudoephedrine hydrochloride tablet Magna Pharmaceuticals, Inc.

Stahist AD

**Drug Facts** 

| Active Ingredients         |         |
|----------------------------|---------|
| (in each immediate-release | tablet) |

Purpose

Chlorcyclizine HCl 25 mg

Pseudoephedrine HCl 60 mg

Antihistamine
Nasal Decongestant

#### Uses

temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes
- nasal congestion
- reduces swelling of nasal passages

#### **Warnings**

Do not exceed recommended dosage.

#### Do not use this product

 If you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for two weeks after stopping the MAOI drug. If you do not know if your prescription drug contains MAOI, ask a doctor or pharmacist before taking this product.

#### Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- heart disease
- high blood pressure
- thyroid disease
- diabetes melitus
- difficulty in urination due to enlargement of the prostate gland

#### Ask a doctor before use if you are taking sedatives or tranquilizers

#### When using this product

• excitability may occur, especially in children

- may cause drowsiness
- alcohol, sedatives and tranquilizers may increase drowsiness effect
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery

#### Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- symptoms do not improve within 7 days or are accompanied by fever
- new symptoms occur

If pregnant or breast-feeding, ask a health professional before use.

#### Keep out of reach of children.

In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.

#### **Directions**

#### Do not exceed recommended dosage.

| Adults and children 12    | 1 tablet by mouth every 6-8 hours, not to exceed 3 tablets in |
|---------------------------|---|
| years of age and over:    | 24 hours, or as directed by a doctor                          |
| Children 6 to under 12    | ½ tablet by mouth every 6-8 hours, not to exceed 1½ tablets   |
| years of age:             | in 24 hours, or as directed by a doctor                       |
| Children under 6 years of | Consult a doctor  |
| age                       | Consult a doctor  |

#### **Inactive ingredients**

Magnesium Stearate, Microcrystalline Cellulose, Sodium Starch Glycolate

#### **Questions or Comments?**

Call 1-888-206-5525 • www.magnaweb.com

Rev. 10/11

58407-0625-30

# Stahist AD

## **Antihistamine • Nasal Decongestant**

**Each tablet contains:** 

Chlorcyclizine HCl ..... 25 mg

Pseudoephedrine HCl ..... 60 mg



30 Tablets

Add image transcription here...

#### STAHIST AD

chlorcyclizine hydrochloride and pseudoephedrine hydrochloride tablet

#### **Product Information**

Product Type HUMAN OTC DRUG Item Code (Source) NDC:58407-625

Route of Administration ORAL

| Active Ingredient/Active Moiety  |                                  |          |  |
|--|----------------------------------|----------|--|
| Ingredient Name  | <b>Basis of Strength</b>         | Strength |  |
| CHLORCYCLIZINE HYDROCHLORIDE (UNII: NPB7A7874U) (CHLORCYCLIZINE - UNII:M26C4IP44P)   | CHLORCYCLIZ INE<br>HYDROCHLORIDE | 25 mg    |  |
| PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F) | PSEUDOEPHEDRINE<br>HYDROCHLORIDE | 60 mg    |  |

| Inactive Ingredients                                     |          |  |  |
|--|----------|--|--|
| Ingredient Name  | Strength |  |  |
| MAGNESIUM STEARATE (UNII: 70097M6I30)                    |          |  |  |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)            |          |  |  |
| SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2) |          |  |  |

| Product Characteristics |       |              |          |
|-------------------------|-------|--------------|----------|
| Color                   | white | Score        | 2 pieces |
| Shape                   | OVAL  | Size         | 16mm     |
| Flavor                  |       | Imprint Code | M625     |
| Contains                |       |              |          |

| Packaging |                      |  |                         |                       |
|-----------|----------------------|--|-------------------------|-----------------------|
| #         | Item Code            | Package Description                                    | Marketing Start<br>Date | Marketing End<br>Date |
| 1         | NDC:58407-625-<br>30 | 1 in 1 BOX   | 12/20/2011              |                       |
| 1         |                      | 30 in 1 BOTTLE; Type 0: Not a Combination Product      |                         |                       |
| 2         | NDC:58407-625-<br>06 | 6 in 1 BOX   | 12/20/2011              |                       |
| 2         | NDC:58407-625-<br>01 | 1 in 1 BLISTER PACK; Type 0: Not a Combination Product |                         |                       |
| 3         | NDC:58407-625-<br>01 | 1 in 1 BLISTER PACK; Type 0: Not a Combination Product | 09/01/2019              |                       |

| Marketing Information |   |                         |                       |
|-----------------------|---|-------------------------|-----------------------|
| Marketing<br>Category | Application Number or Monograph<br>Citation | Marketing Start<br>Date | Marketing End<br>Date |
| OTC Monograph Drug    | M012  | 12/20/2011              |                       |
|                       |   |                         |                       |

### Labeler - Magna Pharmaceuticals, Inc. (620988360)

Registrant - Wittman Pharma, Inc. (830980947)

Revised: 10/2023 Magna Pharmaceuticals, Inc.