

**LAXATIVE ENTERIC COATED- bisacodyl tablet, coated  
Dolgenercorp, Inc. (DOLLAR GENERAL & REXALL)**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Drug Facts**

**Active ingredient (in each tablet)**

Bisacodyl, USP 5 mg

**Purpose**

Stimulant laxative

**Use**

- for temporary relief of occasional constipation and irregularity
- this product generally produces a bowel movement in 6 to 12 hours

**Warnings**

**Do not use**

if you cannot swallow without chewing.

**Ask a doctor before use if you have**

- stomach pain, nausea or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

**When using this product**

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- it may cause stomach discomfort, faintness or cramps

**Stop use and ask a doctor if**

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

**If pregnancy or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- take with a glass of water

adults and children 12 years of age and over	take 1 to 3 tablets in a single daily dose
children 6 to under 12 years of age	take 1 tablet in a single daily dose
children under 6 years of age	ask a doctor

### Other information

- store between 20-25°C (68°-77°F)

### Inactive ingredients

acacia, anhydrous calcium sulfate, anhydrous lactose, carnauba wax, colloidal silicon dioxide, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, gelatin, iron oxide, iron oxide black, iron oxide yellow, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl acetate phthalate, povidone, shellac, sodium starch glycolate, stearic acid, sugar, talc, titanium dioxide

### Questions or comments?

Call **1-888-309-9030**

### Principal Display Panel

**Compare to the active ingredient of Dulcolax® Laxative\***

#### **Laxative Tablets**

Bisacodyl, 5 mg

Stimulant Laxative

Gentle, Predictable Overnight Relief

Tablet

Comfort Coated

\*This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals, Inc., distributor of Dulcolax® Laxative.

DISTRIBUTED BY OLD EAST MAIN CO.

100 MISSION RIDGE

GOODLETTSVILLE, TN 37072

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS  
TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

**Product Label**

Overnight Relief

# Laxative Tablets

Bisacodyl, 5 mg  
Stimulant Laxative



Compare to the  
active ingredient  
of Dulcolax®  
Laxative\*

# Laxative Tablets

Bisacodyl, 5 mg  
Stimulant Laxative

Gentle, Predictable Overnight Relief



25 Tablets  
Comfort Coated

Actual Tablet Size

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Lot No.:  
Exp. Date:

\*This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals, Inc., distributor of Dulcolax® Laxative.

DISTRIBUTED BY OLD EAST MAIN CO.  
100 MISSION RIDGE  
GOODLETTSVILLE, TN 37072  
**100% SATISFACTION GUARANTEED**  
(888) 309-9030

PLD-F23K FC007455



<p><b>Drug Facts</b></p> <p><b>Active ingredient (in each tablet)</b> Bisacodyl USP 5 mg.....Stimulant laxative</p> <p><b>Purpose</b></p>		<p><b>Use</b></p> <ul style="list-style-type: none"> <li>for temporary relief of occasional constipation and irregularity</li> <li>this product generally produces a bowel movement in 6 to 12 hours</li> </ul>							
<p><b>Warnings</b></p> <p>Do not use if you cannot swallow without chewing.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> <li>stomach pain, nausea or vomiting</li> <li>a sudden change in bowel habits that lasts more than 2 weeks</li> </ul> <p>When using this product</p> <ul style="list-style-type: none"> <li>do not chew or crush tablet(s)</li> <li>do not use within 1 hour after taking an antacid or milk</li> <li>it may cause stomach discomfort, faintness and cramps</li> </ul> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> <li>you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.</li> <li>you need to use a laxative for more than 1 week</li> </ul>		<p><b>Other information</b></p> <ul style="list-style-type: none"> <li>store between 20-25°C (68-77°F)</li> </ul>							
<p><b>Directions</b></p> <p>take with a glass of water</p> <table border="1"> <tr> <td>adults and children 12 years of age and over</td> <td>take 1 to 3 tablets in a single daily dose</td> </tr> <tr> <td>children 6 to under 12</td> <td>take 1 tablet in a single daily dose</td> </tr> <tr> <td>children under 6 years of age</td> <td>ask a doctor</td> </tr> </table>		adults and children 12 years of age and over	take 1 to 3 tablets in a single daily dose	children 6 to under 12	take 1 tablet in a single daily dose	children under 6 years of age	ask a doctor	<p><b>Inactive ingredients</b> acacia, anhydrous calcium sulfate, anhydrous lactose, carnauba wax, colloidal silicon dioxide, corn starch, D&amp;C yellow #10 aluminum lake, FD&amp;C yellow #6 aluminum lake, gelatin, iron oxide black, iron oxide yellow, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl acetate, phthalate, povidone, shellac, sodium starch glycolate, stearic acid, sugar, talc, titanium dioxide</p>	
adults and children 12 years of age and over	take 1 to 3 tablets in a single daily dose								
children 6 to under 12	take 1 tablet in a single daily dose								
children under 6 years of age	ask a doctor								
<p><b>Drug Facts (continued)</b></p> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>		<p><b>Questions or comments?</b> Call 1-888-309-9030</p>							

**DOLLAR GENERAL Laxative Tablets**

**LAXATIVE ENTERIC COATED**

bisacodyl tablet, coated

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55910-190
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	5 mg

**Inactive Ingredients**

Ingredient Name	Strength
ACACIA (UNII: 5C5403N26O)	
CALCIUM SULFATE ANHYDROUS (UNII: E934B3V59H)	

<b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)
<b>GELATIN</b> (UNII: 2G86QN327L)
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)
<b>FERROSFERRIC OXIDE</b> (UNII: XM0M87F357)
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)
<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)
<b>Polyvinyl Acetate Phthalate</b> (UNII: 58QVG85GW3)
<b>POVIDONE</b> (UNII: FZ989GH94E)
<b>SHELLAC</b> (UNII: 46N107B71O)
<b>SODIUM STARCH GLYCOLATE TYPE A CORN</b> (UNII: AG9B65PV6B)
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)
<b>SUCROSE</b> (UNII: C151H8M554)
<b>TALC</b> (UNII: 7SEV7J4R1U)
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)

### Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	6mm
<b>Flavor</b>		<b>Imprint Code</b>	TCL003
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-190-25	1 in 1 CARTON	06/20/2013	06/20/2024
1		25 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	06/20/2013	06/20/2024

**Labeler** - Dolgencorp, Inc. (DOLLAR GENERAL & REXALL) (068331990)