

IBUPROFEN THOMPSON- ibuprofen tablet, coated

Thompson Safety

Thompson Ibuprofen

Drug Facts

Active ingredient (in each tablet)

Ibuprofen 200 mg (NSAID*)

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

Temporarily relieves minor aches and pains associated with

- headache ■ toothache ■ backache ■ menstrual cramps
- common cold ■ muscular aches ■ minor arthritis pain

Temporarily reduces fever.

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives ■ skin reddening ■ asthma (wheezing) ■ facial swelling ■ rash ■ shock ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack or stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer

- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- redness or swelling is present in the painful area
- any new or unexpected symptoms occur

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not use more than directed**
- **the smallest effective dose should be used**

■ do not take longer than 10 days, unless directed by a doctor (see Warnings)

Adults and children:(12 years and older)

Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used.

Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

Children under 12 years:

Ask a doctor

Other information

- read all product information before using
- store at 68-77°F (20-25°C)
- avoid excessive heat 104°F (above 40°C)
- tamper evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

carnauba wax*, corn starch, hypromellose*, iron oxide red, lactose*, magnesium stearate*, microcrystalline cellulose*, polydextrose*, polyethylene glycol, povidone (K30)*, silicon dioxide, sodium starch glycolate, stearic acid, talc*, titanium dioxide

*may contain

Questions or comments? 877.506.4291

Ibuprofen

PainReliever/Fever Reducer (NSAID)

Thompson

Safety Works | Business Thrives™

Pull to Open

This package is for Households without Young Children.

Aches, Fever

Ibuprofen (NSAID) 200 mg

100 Tablets

Tamper Evident

Sealed Packets

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Purpose Ibuprofen 200 mg (NSAID) Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug</p> <p>Uses Temporarily relieves minor aches and pains associated with: ■ headache ■ toothache ■ backache ■ menstrual cramps ■ common cold ■ muscular aches ■ minor arthritis pain Temporarily reduces fever.</p> <p>Warnings Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ skin redness ■ asthma (wheezing) ■ facial swelling ■ rash ■ shock ■ dizziness If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)</p>	<p>Drug Facts (continued)</p> <p>■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed</p> <p>Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p> <p>Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery</p> <p>Ask a doctor before use if ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ stomach bleeding warning applies to you ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke ■ you are taking a diuretic</p> <p>Ask a doctor or pharmacist before use if you are ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug (continued on opposite panel)</p>	<p>IBUPROFEN</p> <p>PAIN RELIEVER/FEVER REDUCER (NSAID)</p> <p>THOMPSON SAFETY WORKS BUSINESS THRIVES</p> <p>Pull to Open</p> <p>THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.</p> <p>ACHES, FEVER IBUPROFEN (NSAID) 200 MG</p> <p>100 Tablets TAMPER EVIDENT SEALED PACKETS</p>	<p>Drug Facts (continued)</p> <p>When using this product ■ take with food or milk. If stomach upset occurs</p> <p>Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ redness or swelling is present in the painful area ■ any new or unexpected symptoms occur</p> <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p>	<p>Drug Facts (continued)</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions ■ do not use more than directed ■ the smallest effective dose should be used ■ do not take longer than 10 days, unless directed by a doctor (see Warnings)</p> <p>Adults and children (12 years and older) Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.</p> <p>Children under 12 years: Ask a doctor</p> <p>Other information ■ read all product information before using ■ store at 68-77°F (20-25°C) ■ avoid excessive heat 104°F (above 40°C) ■ tamper evident sealed packets ■ do not use any opened or torn packets (continued on opposite panel)</p>	<p>Drug Facts (continued)</p> <p>Inactive ingredients carnauba wax*, corn starch, hydroxypropylcellulose, iron oxide red, lactose*, magnesium stearate*, microcrystalline cellulose*, polydextrose*, polyethylene glycol, povidone (K-30)*, silicon dioxide, sodium starch glycolate, stearic acid, talc*, titanium dioxide *may contain</p> <p>Questions or comments? 877.506.4291</p> <p>THOMPSON SAFETY WORKS BUSINESS THRIVES</p> <p>Reorder C590833 Distributed by Thompson Safety Houston, TX 77073 877.506.4291 www.thompson-safety.com</p>
--	---	---	--	---	---

Ibuprofen
PainReliever/Fever Reducer (NSAID)
Thompson
Safety Works | Business Thrives™
Pull to Open
This package is for Households without Young Children.
Aches, Fever
Ibuprofen (NSAID) 200 mg
100 Tablets
Tamper Evident
Sealed Packets

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Purpose Ibuprofen 200 mg (NSAID) Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug</p> <p>Uses Temporarily relieves minor aches and pains associated with: ■ headache ■ toothache ■ backache ■ menstrual cramps ■ common cold ■ muscular aches ■ minor arthritis pain Temporarily reduces fever.</p> <p>Warnings Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ skin redness ■ asthma (wheezing) ■ facial swelling ■ rash ■ shock ■ dizziness If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)</p>	<p>Drug Facts (continued)</p> <p>■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed</p> <p>Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p> <p>Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery</p> <p>Ask a doctor before use if ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ stomach bleeding warning applies to you ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke ■ you are taking a diuretic</p> <p>Ask a doctor or pharmacist before use if you are ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug (continued on opposite panel)</p>	<p>IBUPROFEN</p> <p>PAIN RELIEVER/FEVER REDUCER (NSAID)</p> <p>THOMPSON SAFETY WORKS BUSINESS THRIVES</p> <p>Pull to Open</p> <p>THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.</p> <p>ACHES, FEVER IBUPROFEN (NSAID) 200 MG</p> <p>100 Tablets TAMPER EVIDENT SEALED PACKETS</p>	<p>Drug Facts (continued)</p> <p>When using this product ■ take with food or milk. If stomach upset occurs</p> <p>Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ redness or swelling is present in the painful area ■ any new or unexpected symptoms occur</p> <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p>	<p>Drug Facts (continued)</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions ■ do not use more than directed ■ the smallest effective dose should be used ■ do not take longer than 10 days, unless directed by a doctor (see Warnings)</p> <p>Adults and children (12 years and older) Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.</p> <p>Children under 12 years: Ask a doctor</p> <p>Other information ■ read all product information before using ■ store at 68-77°F (20-25°C) ■ avoid excessive heat 104°F (above 40°C) ■ tamper evident sealed packets ■ do not use any opened or torn packets (continued on opposite panel)</p>	<p>Drug Facts (continued)</p> <p>Inactive ingredients carnauba wax*, corn starch, hydroxypropylcellulose, iron oxide red, lactose*, magnesium stearate*, microcrystalline cellulose*, polydextrose*, polyethylene glycol, povidone (K-30)*, silicon dioxide, sodium starch glycolate, stearic acid, talc*, titanium dioxide *may contain</p> <p>Questions or comments? 877.506.4291</p> <p>THOMPSON SAFETY WORKS BUSINESS THRIVES</p> <p>Reorder C590833 Distributed by Thompson Safety Houston, TX 77073 877.506.4291 www.thompson-safety.com</p>
--	---	---	--	---	---

IBUPROFEN THOMPSON			
ibuprofen tablet, coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73408-809
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength

IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg		
Inactive Ingredients				
Ingredient Name		Strength		
FERRIC OXIDE RED (UNII: 1K09F3G675)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
POVIDONE K30 (UNII: U725QWY32X)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
STARCH, CORN (UNII: O8232NY3SJ)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)				
TALC (UNII: 7SEV7J4R1U)				
Product Characteristics				
Color	red (Reddish Brown)	Score	no score	
Shape	ROUND	Size	10mm	
Flavor		Imprint Code	G;2	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73408-809-33	50 in 1 BOX	10/04/2021	
1		2 in 1 PACKET; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA079174	10/04/2021		

IBUPROFEN THOMPSON			
ibuprofen tablet, coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73408-509
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength

IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg
---	-----------	--------

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STARCH, CORN (UNII: O8232NY3SJ)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
LACTOSE, UNSPECIFIED FORM (UNII: J2B2A4N98G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYDEXTROSE (UNII: VH2XOU12IE)	

Product Characteristics

Color	red (Reddish Brown)	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	G;2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73408-509-33	50 in 1 BOX	10/04/2021	
1		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075010	10/04/2021	

Labeler - Thompson Safety (080998015)

Registrant - Unifirst First Aid Corporation (832947092)