

**PEG-3350, ELECTROLYTES, AND ASCORBATE- polyethylene glycol 3350, sodium sulfate, sodium chloride, potassium chloride, sodium ascorbate, and ascorbic acid**  
**Oceanside Pharmaceuticals**

-----

**HIGHLIGHTS OF PRESCRIBING INFORMATION**

These highlights do not include all the information needed to use POLYETHYLENE GLYCOL-3350, ELECTROLYTES, AND ASCORBATE FOR ORAL SOLUTION safely and effectively. See full prescribing information for POLYETHYLENE GLYCOL-3350, ELECTROLYTES, AND ASCORBATE FOR ORAL SOLUTION.

**POLYETHYLENE GLYCOL-3350, ELECTROLYTES, AND ASCORBATE FOR ORAL SOLUTION (polyethylene glycol 3350, sodium sulfate, sodium chloride, potassium chloride, sodium ascorbate, and ascorbic acid for oral solution)**  
**Initial U.S. Approval: 2006**

----- **INDICATIONS AND USAGE** -----

Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution is an osmotic laxative indicated for cleansing of the colon as a preparation for colonoscopy in adults. (1)

----- **DOSAGE AND ADMINISTRATION** -----

Preparation and Administration:

- Two doses of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution are required for a complete preparation for colonoscopy, using a “Two-Day” preferred method or “One-Day” alternative method dosing regimen. (2.1)
- Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution must be reconstituted in water prior to ingestion. (2.1)
- Additional clear liquids must be consumed after each dose of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution in both dosing regimens. (2.1, 5.1)
- Do not take other laxatives while taking Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution. (2.1, 5.5)
- Do not take oral medications within 1 hour of starting each dose. (2.1)

Dosing Regimen:

- *Two-Day (Split-Dose) (Preferred Method):* Dose 1 the evening before the colonoscopy, and Dose 2 the morning of the colonoscopy (approximately 12 hours after the start of Dose 1, and at least 3 ½ hours prior to the colonoscopy). (2.2)
- *One-Day (Evening Only) (Alternative Method):* Dose 1 at least 3 ½ hours prior to bedtime the evening before the colonoscopy and Dose 2 approximately 1 ½ hours after starting Dose 1 the evening before the colonoscopy. (2.3)
- For complete information on dosing, preparation and administration see full prescribing information (2.1, 2.2, 2.3)

----- **DOSAGE FORMS AND STRENGTHS** -----

For Oral Solution: 2 pouches labeled Pouch A and 2 pouches labeled Pouch B. (3)

- Each Pouch A contains 100 grams of polyethylene glycol (PEG) 3350, 7.5 grams of sodium sulfate, 2.691 grams of sodium chloride, and 1.015 grams of potassium chloride. Pouch A contains lemon flavoring. (3)
- Each Pouch B contains 4.7 grams of ascorbic acid and 5.9 grams of sodium ascorbate. (3)

----- **CONTRAINDICATIONS** -----

- Gastrointestinal (GI) obstruction (4, 5.6)
- Bowel perforation (4, 5.6)
- Gastric retention (4)
- Ileus (4)
- Toxic colitis or toxic megacolon (4)
- Hypersensitivity to any ingredient in Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral

---

**WARNINGS AND PRECAUTIONS**

---

- Risk of fluid and electrolyte abnormalities: Encourage adequate hydration, assess concurrent medications, and consider laboratory assessments prior to and after use. (5.1, 7.1)
- Cardiac arrhythmias: Consider pre-dose and post-colonoscopy ECGs in patients at increased risk. (5.2)
- Seizures: Use caution in patients with a history of seizures and patients at increased risk of seizure, including medications that lower the seizure threshold. (5.3, 7.1)
- Patients with renal impairment or taking concomitant medications that affect renal function: Use caution, ensure adequate hydration and consider laboratory testing. (5.4, 7.1, 8.6)
- Colonic mucosal ulcerations: Consider potential for ulcerations when interpreting colonoscopy findings in patients with known or suspected inflammatory bowel disease. (5.5)
- Suspected GI obstruction or perforation: Rule out the diagnosis before administration. (5.6)
- Patients at risk for aspiration: Observe during administration. (5.7)
- Glucose-6-phosphate dehydrogenase deficiency (G-6-PD): Use with caution. (5.8)
- Risks in patients with phenylketonuria: Contains phenylalanine (5.9)
- Hypersensitivity reactions, including anaphylaxis: Inform patients to seek immediate medical care if symptoms occur. (5.10)

---

**ADVERSE REACTIONS**

---

Most common adverse reactions ( $\geq 5\%$ ) are:

- Two-Day (Split-Dose): malaise, nausea, abdominal pain, vomiting, and upper abdominal pain. (6.1)
- One-Day (Evening-Only): abdominal distension, anal discomfort, thirst, nausea, abdominal pain, sleep disorder, rigors, hunger, malaise, vomiting, and dizziness. (6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact Oceanside Pharmaceuticals at 1-800-321-4576 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

---

**DRUG INTERACTIONS**

---

Drugs that increase risk for fluid and electrolyte imbalance. (7.1)

**See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.**

**Revised: 10/2025**

---

**FULL PRESCRIBING INFORMATION: CONTENTS\***

**1 INDICATIONS AND USAGE**

**2 DOSAGE AND ADMINISTRATION**

2.1 Important Preparation and Administration Instructions

2.2 Two-Day Split-Dosing Regimen (Preferred Method)

2.3 One-Day Evening Only Dosing Regimen (Alternative Method)

**3 DOSAGE FORMS AND STRENGTHS**

**4 CONTRAINDICATIONS**

**5 WARNINGS AND PRECAUTIONS**

5.1 Serious Fluid and Electrolyte Abnormalities

5.2 Cardiac Arrhythmias

5.3 Seizures

5.4 Use in Patients with Renal Impairment

5.5 Colonic Mucosal Ulceration, Ischemic Colitis and Ulcerative Colitis

5.6 Use in Patients with Significant Gastrointestinal Disease

5.7 Aspiration

5.8 Glucose-6-Phosphate Dehydrogenase (G6PD) Deficiency

5.9 Risks in Patients with Phenylketonuria

5.10 Hypersensitivity Reactions

## **6 ADVERSE REACTIONS**

6.1 Clinical Studies Experience

6.2 Postmarketing Experience

## **7 DRUG INTERACTIONS**

7.1 Drugs That May Increase Risks due to Fluid and Electrolyte Abnormalities

7.2 Potential for Reduced Drug Absorption

7.3 Stimulant Laxatives

## **8 USE IN SPECIFIC POPULATIONS**

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Renal Impairment

## **10 OVERDOSAGE**

## **11 DESCRIPTION**

## **12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

## **14 CLINICAL STUDIES**

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

## **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution is an osmotic laxative indicated for cleansing of the colon as a preparation for colonoscopy in adults.

### **2 DOSAGE AND ADMINISTRATION**

#### **2.1 Important Preparation and Administration Instructions**

- Correct fluid and electrolyte abnormalities before treatment with Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution [see *Warnings and Precautions* (5.1)].
- Two doses of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution are required for a complete preparation for colonoscopy. The time interval between the two doses depends on the regimen prescribed and the planned timing of the colonoscopy procedure. [see *Dosage and Administration* (2.2, 2.3)].
- The “Split-Dose” is the preferred method and consists of two separate doses: the first dose is taken the evening before the colonoscopy, and the second dose is taken the next day, the morning of the day of the colonoscopy [see *Dosage and Administration* (2.2)].
- The “Evening Only” is the alternative method and consists of two separate doses:

both doses are taken in the evening before the day of the colonoscopy, with a minimum of 1.5 hours between the start of the first dose and the start of the second dose [see *Dosage and Administration (2.3)*].

- Both Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution dosing regimens require administration of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution using the mixing container provided to reconstitute the contents of Pouch A and B with water to the Fill Line. Do not reconstitute with other liquids and/or add starch-based thickeners to the mixing container [see *Warnings and Precautions (5.7)*].
- Additional clear liquids (including water) must be consumed in both dosing regimens [see *Dosage and Administration (2.2, 2.3)*, *Warnings and Precautions (5.1)*].
- Consume only clear liquids (no solid food) from the start of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution treatment until after the colonoscopy.
- Do not eat or drink alcohol, milk, anything colored red or purple or any other foods containing pulp material.
- Do not take other laxatives while taking Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution [see *Drug Interactions (7.3)*].
- Do not take oral medications within 1 hour before or after starting each dose of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution [see *Drug Interactions (7.2)*].
- Ensure completion of Dose 2, including all additional liquids, at least 2 hours before the colonoscopy.
- **Storage:** After reconstitution, store Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution in an upright position and keep refrigerated. Use within 24 hours after it is mixed in water.

## **2.2 Two-Day Split-Dosing Regimen (Preferred Method)**

The Two-Day Split-Dosing regimen is the preferred dosing method.

Instruct adult patients that on the day before the clinical procedure, they can consume breakfast, followed by a light lunch (no solid foods), and clear soup and/or plain yogurt for dinner, which must be completed at least 1 hour prior to the start of the first Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution dose.

Instruct adult patients to take two separate doses in conjunction with fluids as follows:

Dose 1 – In the evening before the colonoscopy, approximately 10 to 12 hours before Dose 2:

1. Empty the contents of 1 Pouch A and 1 Pouch B into the mixing container that comes with Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution.
2. Add lukewarm water to the Fill Line on the mixing container (32 fluid ounces). Do not add other ingredients to the Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution.
3. Thoroughly mix with a spoon or shake with lid on securely until the contents of Pouch A and B are completely dissolved.
4. Drink 8 ounces of the solution every 15 minutes. This should take about 1 hour. Be sure to drink all the solution.
5. Refill the mixing container halfway to the Fill Line (at least 16 ounces) with a clear liquid and drink all this liquid before going to bed.

Dose 2 - Take next morning, on the day of the colonoscopy, approximately 12 hours after the start of Dose 1 and at least 3 ½ hours prior to colonoscopy:

1. Empty the contents of 1 Pouch A and 1 Pouch B into the mixing container that comes with Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution.
2. Add lukewarm water to the Fill Line on the mixing container (32 fluid ounces). Do not add other ingredients to the Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution.
3. Thoroughly mix with a spoon or shake with lid on securely until the contents of Pouch A and B are completely dissolved.
4. Drink 8 ounces of the solution every 15 minutes. This should take about 1 hour. Be sure to drink all of the solution.
5. Refill the mixing container halfway to the Fill Line (at least 16 ounces) with a clear liquid and drink all this liquid at least 2 hours before the colonoscopy.
6. Consume additional water or clear liquids up to 2 hours before the colonoscopy or as prescribed by your healthcare provider. Then stop drinking liquids until after the colonoscopy.

Stop drinking Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution temporarily or drink each portion at longer intervals if severe bloating, abdominal discomfort or distention occurs, until these symptoms resolve.

### **2.3 One-Day Evening Only Dosing Regimen (Alternative Method)**

The One-Day Evening Only regimen is the alternative dosing method for patients for whom the Split-Dosing regimen is inappropriate.

Instruct adult patients that on the day before the clinical procedure, they can consume breakfast, followed by a light lunch (no solid foods), and clear soup and/or plain yogurt for dinner, which must be completed at least 1 hour prior to the start of the first Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution dose.

Instruct adult patients to take two separate doses in conjunction with fluids as follows:

Dose 1 - At least 3 ½ hours before bedtime the evening before the colonoscopy:

1. Empty the contents of 1 Pouch A and 1 Pouch B into the mixing container that comes with Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution.
2. Add lukewarm water to the Fill Line on the mixing container (32 fluid ounces). Do not add other ingredients to the Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution.
3. Thoroughly mix with a spoon or shake with lid on securely until the contents of Pouch A and B are completely dissolved.
4. Drink 8 ounces of the solution every 15 minutes. This should take about 1 hour. Be sure to drink all the solution.

Dose 2 - At least 1 ½ hours after starting Dose 1 on the evening before the colonoscopy:

1. Empty the contents of 1 Pouch A and 1 Pouch B into the mixing container that comes with Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution.
2. Add lukewarm water to the Fill Line on the mixing container (32 fluid ounces). Do

not add other ingredients to the Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution.

3. Thoroughly mix with a spoon or shake with lid on securely until the contents of Pouch A and B are completely dissolved.
4. Drink 8 ounces of the solution every 15 minutes. This should take about 1 hour. Be sure to drink all of the solution.
5. Refill the mixing container to the Fill Line (32 fluid ounces) with a clear liquid and drink all this liquid before going to bed.
6. Consume additional water or clear liquids up to 2 hours before the colonoscopy or as prescribed by your healthcare provider. *Then stop drinking liquids until after the colonoscopy.*

Stop drinking Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution temporarily or drink each portion at longer intervals if severe bloating, abdominal discomfort or distention occurs, until these symptoms resolve.

### **3 DOSAGE FORMS AND STRENGTHS**

Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution is supplied as a white to yellow powder for reconstitution and is available in a carton that contains 2 pouches labeled Pouch A and 2 pouches labeled Pouch B.

- Each Pouch A contains 100 grams of polyethylene glycol (PEG) 3350, 7.5 grams of sodium sulfate, 2.691 grams of sodium chloride, and 1.015 grams of potassium chloride. Pouch A contains lemon flavoring [*see Description (11)*].
- Each Pouch B contains 4.7 grams of ascorbic acid and 5.9 grams of sodium ascorbate.

### **4 CONTRAINDICATIONS**

Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution is contraindicated in the following conditions:

- Gastrointestinal (GI) obstruction [*see Warnings and Precautions (5.6)*]
- Bowel perforation [*see Warnings and Precautions (5.6)*]
- Gastric retention
- Ileus
- Toxic colitis or toxic megacolon
- Hypersensitivity to any ingredient in Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution [*see Warnings and Precautions (5.10) and Description (11)*]

### **5 WARNINGS AND PRECAUTIONS**

#### **5.1 Serious Fluid and Electrolyte Abnormalities**

Advise patients to hydrate adequately before, during, and after the use of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution. If a patient develops significant vomiting or signs of dehydration after taking Polyethylene Glycol-3350,

Electrolytes, and Ascorbate for Oral Solution, consider performing post-colonoscopy lab tests (electrolytes, creatinine, and BUN).

Bowel preparations can cause fluid and electrolyte disturbances, which can lead to serious adverse reactions including cardiac arrhythmias, seizures, and renal impairment [see *Adverse Reactions (6.2)*]. Correct fluid and electrolyte abnormalities before treatment with Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution. Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution should be used with caution in patients using concomitant medications that increase the risk of electrolyte abnormalities [such as diuretics, angiotensin converting enzyme (ACE) inhibitors or angiotensin receptor blockers (ARBs)] or in patients with known or suspected hyponatremia. Consider performing pre-dose and post-colonoscopy laboratory tests (sodium, potassium, calcium, creatinine, and BUN) in these patients [see *Drug Interactions (7.1)*].

## **5.2 Cardiac Arrhythmias**

There have been rare reports of serious arrhythmias (including atrial fibrillation) associated with the use of ionic osmotic laxative products for bowel preparation. These occur predominantly in patients with underlying cardiac risk factors and electrolyte disturbances. Use caution when prescribing Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution for patients at increased risk of arrhythmias (e.g., patients with a history of prolonged QT, uncontrolled arrhythmias, recent myocardial infarction, unstable angina, congestive heart failure, cardiomyopathy, or electrolyte imbalance). Consider pre-dose and post-colonoscopy ECGs in patients at increased risk of serious cardiac arrhythmias.

## **5.3 Seizures**

There have been rare reports of generalized tonic-clonic seizures and/or loss of consciousness associated with use of bowel preparation products in patients with no prior history of seizures. The seizure cases were associated with electrolyte abnormalities (e.g., hyponatremia, hypokalemia, hypocalcemia, and hypomagnesemia) and low serum osmolality. The neurologic abnormalities resolved with correction of fluid and electrolyte abnormalities.

Use caution when prescribing Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution for patients with a history of seizures and in patients at increased risk of seizure, such as patients taking medications that lower the seizure threshold (e.g., tricyclic antidepressants), patients withdrawing from alcohol or benzodiazepines, or patients with known or suspected hyponatremia [see *Drug Interactions (7.1)*].

## **5.4 Use in Patients with Renal Impairment**

Use Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution with caution in patients with renal impairment or patients taking concomitant medications that affect renal function (such as diuretics, ACE inhibitors, angiotensin receptor blockers, or nonsteroidal anti-inflammatory drugs) [see *Drug Interactions (7.1)*]. These patients may be at risk for renal injury. Advise these patients of the importance of adequate hydration before, during, and after use of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution, and consider performing pre-dose and post-colonoscopy laboratory tests (electrolytes, creatinine, and BUN) in these patients [see *Use in Specific Populations*].

(8.6)].

### **5.5 Colonic Mucosal Ulceration, Ischemic Colitis and Ulcerative Colitis**

Osmotic laxatives may produce colonic mucosal aphthous ulcerations, and there have been reports of more serious cases of ischemic colitis requiring hospitalization. Concurrent use of stimulant laxatives and Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution may increase the risk of mucosal ulceration or ischemic colitis and is not recommended. Consider the potential for mucosal ulcerations resulting from the bowel preparation when interpreting colonoscopy findings in patients with known or suspected inflammatory bowel disease.

### **5.6 Use in Patients with Significant Gastrointestinal Disease**

If gastrointestinal obstruction or perforation is suspected, perform appropriate diagnostic studies to rule out these conditions before administering Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution [see *Contraindications (4)*].

Use with caution in patients with severe ulcerative colitis.

### **5.7 Aspiration**

Patients with impaired gag reflex or other swallowing abnormalities are at risk for regurgitation or aspiration of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution. Observe these patients during the administration of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution. Use with caution in these patients.

Do not combine Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution with starch-based thickeners [see *Dosage and Administration (2.1)*]. Polyethylene glycol (PEG), a component of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution, when mixed with starch-thickened liquids reduces the viscosity of the starch-thickened liquid. When a PEG-based product used for another indication was mixed in starch-based pre-thickened liquids used in patients with dysphagia, thinning of the liquid occurred and cases of choking and potential aspiration were reported.

### **5.8 Glucose-6-Phosphate Dehydrogenase (G6PD) Deficiency**

Since Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution contains sodium ascorbate and ascorbic acid, Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution should be used with caution in patients with glucose-6-phosphate dehydrogenase (G6PD) deficiency, especially G6PD deficiency patients with an active infection, with a history of hemolysis, or taking concomitant medications known to precipitate hemolytic reactions.

### **5.9 Risks in Patients with Phenylketonuria**

Phenylalanine can be harmful to patients with phenylketonuria (PKU). Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution contains phenylalanine, a component of aspartame. Each Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution treatment contains 131 mg of phenylalanine (after hydrolysis of the aspartame molecule *in vivo* to aspartic acid and phenylalanine). Before prescribing Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution to a patient with PKU, consider the combined daily amount of phenylalanine from all sources, including

Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution.

## 5.10 Hypersensitivity Reactions

Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution contains polyethylene glycol (PEG) and lemon flavoring (containing citral, lime oil, maltodextrin, acacia gum, vitamin E, and other natural flavoring agents) and may cause serious hypersensitivity reactions including anaphylaxis, angioedema, rash, urticaria, and pruritus [see *Contraindications (4)*, *Adverse Reactions (6.2)* and *Description (11)*]. Inform patients of the signs and symptoms of anaphylaxis and instruct them to seek immediate medical care should signs and symptoms occur.

## 6 ADVERSE REACTIONS

The following serious or otherwise important adverse reactions for bowel preparations are described elsewhere in the labeling:

- Serious Fluid and Electrolyte Abnormalities [see *Warnings and Precautions (5.1)*]
- Cardiac Arrhythmias [see *Warnings and Precautions (5.2)*]
- Seizures [see *Warnings and Precautions (5.3)*]
- Patients with Renal Impairment [see *Warnings and Precautions (5.4)*]
- Colonic Mucosal Ulceration, Ischemic Colitis and Ulcerative Colitis [see *Warnings and Precautions (5.5)*]
- Patients with Significant Gastrointestinal Disease [see *Warnings and Precautions (5.6)*]
- Aspiration [see *Warnings and Precautions (5.7)*]
- Glucose-6-Phosphate Dehydrogenase (G6PD) Deficiency [see *Warnings and Precautions (5.8)*]
- Risks in Patients with Phenylketonuria [see *Warnings and Precautions (5.9)*]
- Hypersensitivity Reactions [see *Warnings and Precautions (5.10)*]

### 6.1 Clinical Studies Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

The safety of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution as a Two-Day Split-Dosing and One-Day Evening Only Dosing Regimen was evaluated in two randomized, active-controlled, multicenter, investigator-blinded clinical trials in adult patients scheduled to have an elective colonoscopy [see *Clinical Studies (14)*]. The safety analysis for Study 1 included 359 adult patients ranging in age from 18 to 88 years (mean age 59), with 52% female and 48% male patients. The safety analysis for Study 2 included 340 adult patients ranging in age from 21 to 76 years (mean age 53), with 53% male and 47% female patients.

Tables 1 and 2 display adverse reactions reported in at least 2% and 5% of patients in either treatment group in Study 1 and Study 2, respectively. Since diarrhea was considered as a part of the efficacy assessment, it was not defined as an adverse reaction in these trials.

**Table 1: Common Adverse Reactions\* in Patients Undergoing Colonoscopy in Study 1**

	<b>Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution Two-Day Split Dosing Regimen (N=180)</b>	<b>4 Liter PEG + Electrolytes Solution (N=179)</b>
<b>Malaise</b>	19%	18%
<b>Nausea</b>	14%	20%
<b>Abdominal pain</b>	13%	15%
<b>Vomiting</b>	8%	13%
<b>Upper abdominal pain</b>	6%	6%
<b>Dyspepsia</b>	3%	1%

\* Reported in at least 2% of patients in either treatment group

**Table 2: Common Adverse Reactions\*† in Patients Undergoing Colonoscopy in Study 2**

	<b>Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution One-Day Evening Only Dosing Regimen (N=169)</b>	<b>90 mL Oral Sodium Phosphate Solution (N=171)</b>
Abdominal distension	60%	41%
Anal discomfort	51%	52%
Thirst	47%	65%
Nausea	47%	47%
Abdominal pain	39%	32%
Sleep disorder	35%	29%
Rigors	34%	30%
Hunger	30%	71%
Malaise	27%	53%
Vomiting	7%	8%
Dizziness	7%	18%
Headache	2%	5%
Hypokalemia	0%	6%
Hyperphosphatemia	0%	6%

\* Reported in at least 5% of patients in either treatment group

† Patients were specifically asked about the occurrence of the following symptoms: shivering, anal irritations, abdominal bloating or fullness, sleep loss, nausea, vomiting, weakness, hunger sensation, abdominal cramps or pain, thirst sensation, and dizziness.

## 6.2 Postmarketing Experience

The following adverse reactions have been identified during post-approval use of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution or other PEG-based products. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

*Cardiovascular:* tachycardia, palpitations, hypertension, arrhythmia, atrial fibrillation, peripheral edema, asystole, acute pulmonary edema and syncope, and dehydration.

*Gastrointestinal:* upper gastrointestinal bleeding from a Mallory-Weiss tear, esophageal perforation [usually with gastroesophageal reflux disease (GERD)].

*Hypersensitivity reactions:* anaphylaxis (some of which were severe, including shock), rash, urticaria, pruritus, lip, tongue and facial swelling, dyspnea, chest tightness and throat tightness, rhinorrhea, dermatitis, fever, and chills [see *Warnings and Precautions (5.10)*].

*Nervous system:* tremor, seizure.

*Renal:* renal impairment and/or failure.

## **7 DRUG INTERACTIONS**

### **7.1 Drugs That May Increase Risks due to Fluid and Electrolyte Abnormalities**

Use caution when prescribing Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution for patients with conditions and/or who are using medications that increase the risk for fluid and electrolyte disturbances or may increase the risk of renal impairment, seizures, arrhythmias, or QT prolongation in the setting of fluid and electrolyte abnormalities [see *Warnings and Precautions (5.1, 5.2, 5.3, 5.4)*]. Consider additional patient evaluations as appropriate.

### **7.2 Potential for Reduced Drug Absorption**

Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution can reduce the absorption of other co-administered drugs. Administer oral medications at least 1 hour before the start of administration of each dose of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution [see *Dosage and Administration (2.1)*].

### **7.3 Stimulant Laxatives**

Concurrent use of stimulant laxatives and Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution may increase the risk of mucosal ulceration or ischemic colitis. Avoid use of stimulant laxatives (e.g., bisacodyl, sodium picosulfate) while taking Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution [see *Warnings and Precautions (5.5, 5.6)*].

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

## Risk Summary

There are no available data on Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution in pregnant women to inform a drug-associated risk for adverse developmental outcomes. Animal reproduction studies have not been conducted with Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution.

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

## **8.2 Lactation**

### Risk Summary

There are no data available on the presence of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution in human milk, the effects of the drug on the breastfed child, or the effects of the drug on milk production. The lack of clinical data during lactation precludes a clear determination of the risk of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution to a child during lactation; therefore, the developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution and any potential adverse effects on the breastfed child from Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution or from the underlying maternal condition.

## **8.4 Pediatric Use**

The safety and effectiveness of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution in pediatric patients have not been established.

## **8.5 Geriatric Use**

Of the 413 patients in clinical trials receiving Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution, 91 (22%) patients were aged 65 or older, while 25 (6%) patients were over 75 years of age. No overall differences in safety or effectiveness were observed between geriatric patients and younger patients, and other reported clinical experience has not identified differences in responses between geriatric patients and younger patients. However, elderly patients are more likely to have decreased hepatic, renal or cardiac function and may be more susceptible to adverse reactions resulting from fluid and electrolyte abnormalities [see *Warnings and Precautions (5.1)*].

## **8.6 Renal Impairment**

Use Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution with caution in patients with renal impairment or patients taking concomitant medications that may affect renal function [see *Drug Interactions (7.1)*]. These patients may be at risk for renal injury. Advise these patients of the importance of adequate hydration before, during and after the use of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution, and consider performing baseline and post-colonoscopy laboratory tests (electrolytes, creatinine, and BUN) in these patients [see *Warnings and Precautions*

(5.4)].

## **10 OVERDOSAGE**

Overdosage of more than the recommended dose of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution may lead to severe electrolyte disturbances, including hyponatremia and/or hypokalemia, as well as dehydration and hypovolemia, with signs and symptoms of these disturbances. Certain severe electrolyte disturbances may lead to cardiac arrhythmias, seizures, and renal failure [see *Warnings and Precautions* (5.1, 5.2, 5.3)]. Monitor for fluid and electrolyte disturbances and treat symptomatically.

## **11 DESCRIPTION**

Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution is an osmotic laxative consisting of 4 pouches (2 of Pouch A and 2 of Pouch B) containing white to yellow powder for reconstitution.

Each Pouch A contains 100 grams of polyethylene glycol (PEG) 3350, 7.5 grams of sodium sulfate, 2.691 grams of sodium chloride, and 1.015 grams of potassium chloride, plus the following excipients: aspartame (sweetener), acesulfame potassium (sweetener), and lemon flavoring containing citral, lime oil, maltodextrin, acacia gum, vitamin E, and other natural flavoring agents.

Pouch A contains 111.9 g of powder for oral solution.

Each Pouch B contains 4.7 grams of ascorbic acid and 5.9 grams of sodium ascorbate. Pouch B contains 10.6 g of powder for oral solution.

When 1 Pouch A and 1 Pouch B are dissolved together in water to a volume of 1 liter, Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution is an oral solution having a lemon taste.

The entire, reconstituted, 2-liter Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution colon preparation contains 200 grams of polyethylene glycol-3350, 15 grams of sodium sulfate, 5.38 grams of sodium chloride, 2.03 grams of potassium chloride, 9.4 grams of ascorbic acid, and 11.8 grams of sodium ascorbate plus the following excipients: aspartame (sweetener), acesulfame potassium (sweetener), and lemon flavoring.

A mixing container for reconstitution is enclosed.

Phenylketonurics: Contains Phenylalanine 131 mg per treatment.

## **12 CLINICAL PHARMACOLOGY**

### **12.1 Mechanism of Action**

The primary mode of action is osmotic action of polyethylene glycol 3350, sodium sulfate, sodium chloride, potassium chloride, sodium ascorbate, and ascorbic acid, which induce a laxative effect. The physiological consequence is increased water retention in the lumen of the colon, resulting in loose stools.

## 14 CLINICAL STUDIES

The colon cleansing efficacy and safety of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution was evaluated in two randomized, actively-controlled, multi-center, investigator-blinded trials in adult patients scheduled to have an elective colonoscopy.

In Study 1, patients were randomized to one of the following two colon preparation treatments: 1) 2 liters of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution with 1 additional liter of clear liquid split into two doses (during the evening before and the morning of the colonoscopy) and 2) 4 liters of polyethylene glycol plus electrolytes solution (4L PEG + E) split into two doses (during the evening before and the morning of the colonoscopy). Patients were allowed to have a morning breakfast, a light lunch, clear soup and/or plain yogurt for dinner. Dinner had to be completed at least one hour prior to initiation of the colon preparation administration.

The primary efficacy endpoint was the proportion of patients with effective colon cleansing as judged by blinded gastroenterologists on the basis of videotapes recorded during the colonoscopy.

The blinded gastroenterologists graded the colon cleansing twice (during introduction and withdrawal of the colonoscope) and the poorer of the two assessments was used in the primary efficacy analysis.

The efficacy analysis included 308 adult patients who had an elective colonoscopy. Patients ranged in age from 18 to 88 years old (mean age about 59 years old) with 52% female and 48% male patients. Table 3 displays the results.

**Table 3: Effectiveness of Overall Colon Cleansing of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution vs. 4 Liter Polyethylene Glycol plus Electrolytes Solution in Study 1**

	<b>Responders A* or B† (%)</b>	<b>C‡ (%)</b>	<b>D (%)</b>
<b>Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution (N=153)</b>	88.9	9.8	1.3
<b>4L PEG + E§ (N=155)</b>	94.8	4.5	0.6

\* A: colon empty and clean or presence of clear liquid, but easily removed by suction

† C: semisolid amounts of stool, only partially removable with a risk of incomplete visualization of the gut mucosa

‡ D: semisolid or solid amounts of stool; consequently colonoscopy incomplete or needed to be terminated.

§ 4L PEG + E is 4 Liter Polyethylene Glycol plus Electrolytes Solution.

4L PEG+E's responder rate was not significantly higher than Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution's responder rate.

In Study 2, patients were randomized to one of the following two colon preparation treatments: 1) 2 liters of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution with 1 additional liter of clear liquid in the evening prior to the colonoscopy and 2) 90 mL of oral sodium phosphate solution (90 mL OSPS) with at least 2 liters of additional clear liquid during the day and evening prior to the colonoscopy. Patients randomized to Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution therapy were allowed to have a morning breakfast; a light lunch; and clear soup and/or plain yogurt for dinner. Dinner had to be completed at least one hour prior to initiation of the colon preparation administration.

The primary efficacy endpoint was the proportion of patients with effective colon cleansing as judged by the colonoscopist and one blinded gastroenterologist (on the basis of videotapes recorded during the colonoscopy). In case of a discrepancy between the colonoscopist and the blinded gastroenterologist, a second blinded gastroenterologist made the final efficacy determination.

The efficacy analysis included 280 adult patients who had an elective colonoscopy. Patients ranged in age from 21 to 76 years old (mean age about 53 years old) with 47% female and 53% male patients. Table 4 displays the results.

**Table 4: Effectiveness of Overall Colon Cleansing of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution vs. 90 mL Oral Sodium Phosphate Solution in Study 2**

	<b>Responders A* or B† (%)</b>	<b>C‡ (%)</b>	<b>D§ (%)</b>
<b>Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution (N=137)</b>	73.0	23.4	3.6
<b>90 mL OSPS¶ (N=143)</b>	64.4	29.4	6.3

\* A: empty and clean or clear liquid (transparent, yellow, or green)

† B: brown liquid or semisolid remaining small amounts of stool, fully removable by suction or displaceable allowing a complete visualization of the underlying mucosa

‡ C: semisolid only partially removable/displaceable stools; risk of incomplete examination of the underlying mucosa

§ D: heavy and hard stool making the segment examination uninterpretable and, consequently, the colonoscopy needed to be terminated.

¶ OSPS is Oral Sodium Phosphate Solution.

Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution's responder rate was not significantly higher than OSPS's responder rate.

## 16 HOW SUPPLIED/STORAGE AND HANDLING

Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution (polyethylene glycol 3350, sodium sulfate, sodium chloride, potassium chloride, sodium ascorbate, and ascorbic acid for oral solution) is supplied as a white to yellow powder for reconstitution and is lemon flavored.

NDC 68682-201-75, Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution, single-use outer carton:

- Each outer carton contains a disposable mixing container with lid for reconstitution of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution, prescribing information and patient information, and one inner carton.
- Each inner carton contains 2 pouches labeled Pouch A and 2 pouches labeled Pouch B.

### Storage

Store carton/container at room temperature, between 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F). When reconstituted, store upright and keep solution refrigerated. Use within 24 hours [see *Dosage and Administration (2.1)*].

## 17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide and Instructions for Use).

Instruct patients:

- That two doses of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution are required for a complete preparation for colonoscopy either as a Split-Dose (2-Day), or Evening Only (1-Day) dosing regimen [see *Instructions for Use*].
- Not to take other laxatives while they are taking Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution.
- That Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution contains 131 mg of phenylalanine per treatment [see *Warnings and Precautions (5.9)*].
- That each pouch needs to be reconstituted in water before ingestion and that they should drink additional clear liquids. Examples of clear liquids can be found in the *Instructions for Use*.
- Not to take oral medications within one hour of starting each dose of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution.
- To follow the directions in the *Instructions for Use*, for either the Two-Day Split-Dosing or the One-Day Evening Only Dosing regimen, as prescribed.
- To consume additional clear liquids before, during, and after the use of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution to prevent dehydration [see *Warnings and Precautions (5.1)*].
- To contact their healthcare provider if they develop significant vomiting or signs of dehydration after taking Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution or if they experience altered consciousness or seizures [see *Warnings*].

and Precautions (5.1, 5.2, 5.3, 5.4)]

- Not to eat or drink alcohol, milk, anything colored red or purple or any other foods containing pulp material.
- To stop drinking Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution temporarily or drink each portion at longer intervals if they develop severe abdominal discomfort or distention until these symptoms diminish. If severe symptoms persist, tell patients to contact their healthcare provider.

**Distributed by:**

Oceanside Pharmaceuticals, a division of  
Bausch Health US, LLC  
Bridgewater, NJ 08807 USA

© 2025 Bausch Health Companies Inc. or its affiliates

9676405 70015788-02

**MEDICATION GUIDE**

**POLYETHYLENE GLYCOL-3350, ELECTROLYTES, AND ASCORBATE FOR ORAL SOLUTION**

**(polyethylene glycol 3350, sodium sulfate, sodium chloride, potassium chloride, sodium ascorbate, and ascorbic acid for oral solution)**

**Read this Medication Guide and Instructions for Use before your colonoscopy and again before you start taking Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution.**

**What is the most important information I should know about Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution? Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution and other bowel preparations can cause serious side effects, including:**

- **Serious loss of body fluid (dehydration) and changes in blood salts (electrolytes) in your blood. These changes can cause:**
  - o abnormal heartbeats that can cause death.
  - o seizures. This can happen even if you have never had a seizure.
  - o kidney problems.

**Your chance of having fluid loss and changes in body salts with Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution is higher if you:**

- have heart problems.
- have kidney problems.
- take water pills (diuretics), high blood pressure medicine, or non-steroidal anti-inflammatory drugs (NSAIDs).

**Tell your healthcare provider right away if you have any of these symptoms of serious loss of body fluid (dehydration) while taking Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution:**

- vomiting
- urinating less often than normal

- dizziness
- headache

**See “What are the possible side effects of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution?” for more information about side effects.**

### **What is Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution?**

Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution is a prescription medicine used by adults to clean the colon before a colonoscopy. Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution cleans your colon by causing you to have diarrhea (loose stools). Cleaning your colon helps your healthcare provider see the inside of your colon more clearly during your colonoscopy.

It is not known if Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution is safe and effective in children.

### **Who should not take Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution?**

Do not take Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution if your healthcare provider has told you that you have:

- a blockage in your intestine (bowel obstruction).
- an opening in the wall of your stomach or intestine (bowel perforation).
- problems with food and fluid emptying from your stomach (gastric retention).
- a problem with food moving too slowly through your intestines (ileus).
- a very dilated intestine (toxic megacolon).
- an allergy to any of the ingredients in Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution. See the end of this Medication Guide for a complete list of ingredients in Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution.

### **What should I tell my healthcare provider before taking Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution?**

Before taking Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution, tell your healthcare provider **about all of your medical conditions, including** if you:

- have problems with serious loss of body fluid (dehydration) and changes in blood salts (electrolytes).
- have heart problems.
- have seizures or take medicines for seizures.
- have kidney problems or take medicines for kidney problems.
- have stomach or bowel problems, including ulcerative colitis.
- have problems with swallowing, gastric reflux, or if you inhale food or fluid into your lungs when eating or drinking (aspirate).
- have a condition called glucose-6-phosphate dehydrogenase (G6PD) deficiency that destroys red blood cells.
- are withdrawing from alcohol or benzodiazepines.
- have phenylketonuria (PKU). Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution contains phenylalanine.
- are allergic to any of the ingredients in Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution.

- are pregnant or plan to become pregnant. It is not known if Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution will harm your unborn baby. Talk to your healthcare provider if you are pregnant.
- are breastfeeding or plan to breastfeed. It is not known if Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution passes into your breast milk. You and your healthcare provider should decide if you will take Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution while breastfeeding.

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

**Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution may affect how other medicines work.** Do not take medicines by mouth 1 hour before or after starting each dose of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution.

Especially tell your healthcare provider if you take:

- medicines to treat a blood salt (electrolyte) imbalance.
- medicines for blood pressure or heart problems.
- medicines for seizures (antiepileptics).
- medicines for kidney problems.
- water pills (diuretics).
- non-steroidal anti-inflammatory drugs (NSAIDs).
- laxatives. Do not take other laxatives while taking Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution.
- medicines for depression or other mental health problems.
- starch-based thickeners. For people who have trouble swallowing, do not mix Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution with starch-based thickeners.

Ask your healthcare provider or pharmacist for a list of these medicines if you are not sure if you are taking any of the medicines listed above.

Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

### **How should I take Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution?**

**See the Instructions for Use for dosing instructions.** You must read, understand, and follow these instructions to take Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution the right way.

- Take Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution exactly as your healthcare provider tells you to take it. Your healthcare provider will tell you to take the Two-Day Split-Dosing regimen option or the One-Day Evening Only Dosing regimen option.
- On the day before the procedure, you can have breakfast, followed by a light lunch (no solid foods), and dinner of clear soup with or without plain yogurt, or plain yogurt only. You must finish dinner at least 1 hour before the start of the first Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution dose.
- Drink only clear liquids before, during, and after you take Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution, until 2 hours before your colonoscopy to help prevent fluid loss (dehydration).
- Do not eat solid food while taking Polyethylene Glycol-3350, Electrolytes, and

Ascorbate for Oral Solution until after your colonoscopy.

- Do not eat or drink alcohol, milk, anything colored red or purple or any other foods with pulp.
- It is important for you to drink the additional amount of clear liquids listed in the **Instructions for Use**.
- You may have stomach-area (abdomen) bloating after your first dose of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution.
  - If you have severe stomach-area (abdomen) discomfort or bloating, stop drinking Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution for a short time or wait a longer time between each dose of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution until your stomach-area symptoms improve. If your stomach-area discomfort or bloating continues, tell your healthcare provider.
- If you take too much Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution, call your healthcare provider.

### **What are the possible side effects of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution?**

**Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution can cause serious side effects, including:**

- **Changes in certain blood tests.** Your healthcare provider may do blood tests after you take Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution to check your blood for changes. Tell your healthcare provider if you have any symptoms of too much fluid loss, including:
  - vomiting
  - heart problems
  - seizures
  - dizziness
  - kidney problems
  - dry mouth
  - feel faint, weak or lightheaded especially when you stand up (orthostatic hypotension)
- **Ulcers of the bowel or bowel problems (ischemic colitis).** Tell your healthcare provider right away if you have severe stomach-area (abdomen) pain or rectal bleeding.
- **Serious allergic reactions.** Symptoms of a serious allergic reaction may include:
  - skin rash
  - raised red patches on your skin (hives)
  - kidney problems
  - swelling of the face, lips, tongue and throat
  - itching

**The most common side effects of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution include:**

- o anal discomfort
- o sleep problems
- o vomiting
- o thirst
- o chills
- o indigestion
- o nausea
- o hunger
- o dizziness
- o stomach-area (abdomen pain or bloating) discomfort

These are not all the possible side effects of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

### **How should I store Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution?**

- Store Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution that has not been mixed with water at room temperature, between 68°F to 77°F (20°C to 25°C).
- Store Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution that has been mixed with water in an upright position in the refrigerator.
- Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution should be taken within 24 hours after it has been mixed with water.

### **Keep Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution and all medicines out of the reach of children.**

### **General information about the safe and effective use of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution.**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution for a condition for which it was not prescribed. Do not give Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution to other people, even if they are going to have the same procedure you are. It may harm them.

This Medication Guide summarizes the most important information about Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution. If you would like more information, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information that is written for health professionals.

For more information, call 1-800-321-4576.

### **What are the ingredients in Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution?**

#### **Active ingredients:**

Pouch A: polyethylene glycol (PEG) 3350, sodium sulfate, sodium chloride, potassium chloride.

Pouch B: ascorbic acid and sodium ascorbate.

#### **Inactive ingredients:**

Pouch A: aspartame, acesulfame potassium, and lemon flavoring including citral, lime oil, maltodextrin, acacia gum, vitamin E, and other natural flavoring agents.

**Distributed by:**

Oceanside Pharmaceuticals, a division of  
Bausch Health US, LLC  
Bridgewater, NJ 08807 USA

© 2025 Bausch Health Companies Inc. or its affiliates

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Revised: October 2025

9676405 70015788-02

**Instructions for Use**

POLYETHYLENE GLYCOL-3350, Electrolytes, and Ascorbate for Oral Solution (polyethylene glycol 3350, sodium sulfate, sodium chloride, potassium chloride, sodium ascorbate, and ascorbic acid for oral solution)

There are two different options for taking Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution. Your healthcare provider will tell you to take the Two-Day Split-Dosing Regimen option **or** the One-Day Evening Only Dosing Regimen option.

**Important Information on Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution:**

- You must drink all of the Dose 1 and Dose 2 of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution with either dosing regimen option. Make sure you finish Dose 2 at least 2 hours before your colonoscopy.
- After completing Dose 2, it is important that you drink additional clear liquids (including water), but you must stop drinking all liquids at least 2 hours before your colonoscopy.
- Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution is lemon flavored containing citral, lime oil, maltodextrin, acacia gum, vitamin E, and other natural flavoring agents.
- Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution **must** be mixed with water. **Do not** add any other ingredients to Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution.
- Do not mix Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution with starch-based thickeners.
- After you start taking Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution, you can only drink clear liquids (no solid foods) until after your colonoscopy. Examples of clear liquids include:
  - o water
  - o clear fruit juices without pulp including apple, white grape, or white cranberry
  - o strained limeade or lemonade
  - o coffee or tea (**do not** use any dairy or non-dairy creamer)
  - o clear broth
  - o clear soda

- o gelatin (without added fruit or topping, no red or purple)
- o popsicles (without pieces of fruit or pulp, no red or purple)
- Drink plenty of clear liquids before, during, and after you take Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution, up until 2 hours before your colonoscopy, to help prevent fluid loss (dehydration).
- **Do not** eat or drink anything within 2 hours before your colonoscopy.
- **Do not** eat or drink alcohol, milk, anything colored red or purple or containing pulp.
- **Do not** take other laxatives while taking Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution.
- **Do not** take any medicines by mouth (oral) within 1 hour before or after starting each dose of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution.
- **Do not** eat any solid food while taking Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution until after your colonoscopy.


To take each dose of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution, you will need:

- o Mixing Container that comes with Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution
- o One Pouch A
- o One Pouch B
- o Lukewarm water
- For the Two-Day Split-Dosing Regimen:
  - o On the day before the colonoscopy you can eat breakfast followed by a light lunch (no solid foods). For dinner you may have a clear soup with or without plain yogurt, or plain yogurt only.
  - o You must finish eating at least 1 hour before you start taking Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution.
  - o You must take the first dose between 10 to 12 hours before the second dose. The second dose must be taken at least 3 ½ hours before the colonoscopy.
  - o After you start taking Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution you can only drink clear liquids.
- For the One-Day Evening Only Dosing Regimen:
  - o On the day before the colonoscopy you can eat breakfast followed by a light lunch. For dinner you may have a clear soup with or without plain yogurt, or plain yogurt only.
  - o You must finish eating at least 1 hour before you start taking Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution.
  - o You must take the first dose 3 ½ hours before bedtime the evening before the colonoscopy.
  - o After you start taking Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution you can only drink clear liquids.
  - o Take the second dose 1 ½ hours after starting Dose 1.


### **Two-Day Split-Dosing Regimen Dosing Instructions**

**Dose 1 - Take this dose the evening before your colonoscopy (10 to 12 hours before Dose 2):**


Step 1: Empty the contents of **one** Pouch A and **one** Pouch B into the Mixing Container that comes with Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution.



Step 2: Add lukewarm water to the Fill Line on the Mixing Container. You will need at least 32 ounces.



Step 3: Mix to completely dissolve the medicine from Pouch A and Pouch B into the lukewarm water. To mix the solution, stir the medicine in the Mixing Container with a spoon, or close the lid and shake.




Step 4: Drink one 8 oz. (ounce) glass of the solution every 15 minutes. Be sure to drink all of the solution in the Mixing Container. It should take about 1 hour to drink all the liquid.

If you feel like you have severe stomach pain or discomfort you can stop taking Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution for a short period of time and then continue taking it or you can take smaller sips of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution so that you space out your dose longer than 1 hour. If you still have severe stomach pain, call your healthcare provider.




Step 5: Refill the Mixing Container with 16 oz. (at least halfway to Fill Line and enough for two 8 oz. glasses) of a clear liquid and drink all of this liquid before you go to bed.



**Dose 2 - Take this dose the next morning on the day of your colonoscopy (start at least 3 ½ hours before your colonoscopy):**

Step 1: Repeat steps 1 through 4 from Dose 1 of the Split-Dose (2-Day) instructions.

Step 2: Fill the Mixing Container with 16 oz. (at least halfway to Fill Line and enough for two 8 oz. glasses) of a clear liquid and drink all of this liquid at least 2 hours before your colonoscopy.




Step 3: Drink only clear liquids up to 2 hours before your colonoscopy or as prescribed by your healthcare provider. Then stop drinking liquids until after your colonoscopy.

### **One-Day Evening Only Dosing Regimen Instructions**

#### **Dose 1 (take at least 3 ½ hours before bedtime the evening before your colonoscopy):**


Step 1: Empty the contents of **one** Pouch A and **one** Pouch B into the Mixing Container that comes with Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution.



Step 2: Add lukewarm water to the Fill Line on the Mixing Container. You will need at least 32 ounces.



Step 3: Mix to completely dissolve the medicine from Pouch A and Pouch B into the lukewarm water. To mix the solution, stir the medicine in the Mixing Container with a spoon, or close the lid and shake.



Step 4: Drink one 8 oz. (ounce) glass of the solution every 15 minutes. Be sure to drink all of the solution in the Mixing Container. It should take about 1 hour to drink all the liquid.




If you feel like you have severe stomach pain or discomfort you can stop taking Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution for a short period of time and then continue taking it or you can take smaller sips of Polyethylene Glycol-3350, Electrolytes, and Ascorbate for Oral Solution so that you space out your dose longer than 1 hour. If you still have severe stomach pain, call your healthcare provider.

**Dose 2 (take about 1 ½ hours after starting Dose 1):**

Step 1: Repeat Steps 1 through 4 from Dose 1 of the Evening Only (1-Day) instructions.

Step 2: After you complete steps 1 through 4, fill the Mixing Container again to the Fill Line with clear liquid and drink all of this liquid before you go to bed.



Step 3: Drink only clear liquids up to 2 hours before your colonoscopy or as prescribed by your healthcare provider. Then stop drinking liquids until after your colonoscopy.

This Instructions for Use has been approved by the U.S. Food and Drug Administration.

**Distributed by:**

Oceanside Pharmaceuticals, a division of  
Bausch Health US, LLC  
Bridgewater, NJ 08807 USA

© 2025 Bausch Health Companies Inc. or its affiliates

Revised: October 2025

9676405 70015788-02

**PACKAGE LABEL PRINCIPAL DISPLAY PANEL - POLYETHYLENE GLYCOL-3350,  
Electrolytes, and Ascorbate for Oral Solution Outer Carton**

**Rx only**

**NDC 68682-201-75**


**POLYETHYLENE GLYCOL-3350,  
ELECTROLYTES, AND ASCORBATE  
FOR ORAL SOLUTION**

(Polyethylene Glycol-3350,  
Sodium Sulfate,  
Sodium Chloride, Potassium Chloride,  
Sodium Ascorbate and Ascorbic Acid  
for Oral Solution)

100 g/7.5 g/2.691 g/1.015 g/5.9 g/4.7 g

**OCEANSIDE  
Pharmaceuticals**

9676304 70016299-02



# PEG-3350, ELECTROLYTES, AND ASCORBATE

polyethylene glycol 3350, sodium sulfate, sodium chloride, potassium chloride, sodium ascorbate, and ascorbic acid kit

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:68682-201
---------------------	-------------------------	---------------------------	---------------

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68682-201-75	1 in 1 CONTAINER; Type 0: Not a Combination Product	10/01/2006	

## Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	2 POUCH	2 L
Part 2	2 POUCH	2 L

## Part 1 of 2

### POUCH A

polyethylene glycol 3350, sodium sulfate, sodium chloride, potassium chloride solution

## Product Information

Route of Administration ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>POLYETHYLENE GLYCOL 3350</b> (UNII: G2M7P15E5P) (POLYETHYLENE GLYCOL 3350 - UNII:G2M7P15E5P)	POLYETHYLENE GLYCOL 3350	100 g in 1 L
<b>SODIUM SULFATE</b> (UNII: 0YPR65R21J) (SODIUM SULFATE ANHYDROUS - UNII:36KCS0R750)	SODIUM SULFATE	7.5 g in 1 L
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X) (CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	2.69 g in 1 L
<b>POTASSIUM CHLORIDE</b> (UNII: 660YQ98I10) (POTASSIUM CATION - UNII:295O53K152)	POTASSIUM CHLORIDE	1.015 g in 1 L

## Inactive Ingredients

Ingredient Name	Strength
<b>ASPARTAME</b> (UNII: Z0H242BBR1)	
<b>LEMON</b> (UNII: 24RS0A988O)	
<b>ACESULFAME</b> (UNII: MA3UYZ6K1H)	
<b>CITRAL</b> (UNII: T7EU009VPP)	
<b>LEMON OIL, COLD PRESSED</b> (UNII: I9GRO824LL)	
<b>LIME OIL, COLD PRESSED</b> (UNII: UZH29XGA8G)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		1 L in 1 POUCH; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA authorized generic	NDA021881	10/01/2006	

## Part 2 of 2

### POUCH B

ascorbic acid, sodium ascorbate solution

## Product Information

Route of Administration ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASCORBIC ACID (UNII: PQ6CK8PD0R) (ASCORBIC ACID - UNII:PQ6CK8PD0R)	ASCORBIC ACID	4.7 g in 1 L
SODIUM ASCORBATE (UNII: S033EH8359) (ASCORBIC ACID - UNII:PQ6CK8PD0R)	ASCORBIC ACID	5.9 g in 1 L

## Inactive Ingredients

Ingredient Name	Strength
ASPARTAME (UNII: Z0H242BBR1)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
LEMON (UNII: 24RS0A988O)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		1 L in 1 POUCH; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA authorized generic	NDA021881	10/01/2006	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA021881	10/01/2006	

**Labeler** - Oceanside Pharmaceuticals (832011691)

## Establishment

Name	Address	ID/FEI	Business Operations
Novel Laboratories, Inc.		793518643	MANUFACTURE(68682-201)

## Establishment

Name	Address	ID/FEI	Business Operations
Norgine Limited		239828197	MANUFACTURE(68682-201)

