

**ASPIRIN- aspirin tablet, coated**  
**Wildman Business Group**

-----

**ACTIVE INGREDIENT**

***Active ingredient (in each tablet)***

Aspirin 325 mg (NSAID\*)

\*nonsteroidal anti-inflammatory drug

**PURPOSE**

***Purpose***

Pain reliever/fever reducer

**INDICATIONS & USAGE**

***Uses***

Temporarily relieves

- headache
- muscle pain
- minor pain of arthritis
- pain and fever of colds
- toothache
- menstrual pain

**WARNINGS**

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:**

This product contains an NSAID, which may cause severe stomach bleeding. The chance

is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcohol drinks every day while using this product
- take more or for a longer time than directed

## **DO NOT USE**

**Do not use** if you are allergic to aspirin or to any other pain reliever/fever reducer

## **ASK DOCTOR**

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

## **ASK DOCTOR/PHARMACIST**

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for

- gout
- diabetes
- arthritis

## **STOP USE**

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever lasts more than 3 days
- if ringing in the ears or loss of hearing occurs
- redness or swelling is present
- new symptoms appear

## **PREGNANCY OR BREAST FEEDING**

**If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

## **KEEP OUT OF REACH OF CHILDREN**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

## **DOSAGE & ADMINISTRATION**

### ***Directions***

- drink a full glass of water with each dose

**Adults and children: (12 years and over)** Take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours.

**Children under 12 years:** Consult a doctor

## **OTHER SAFETY INFORMATION**

### ***Other information***

- store at room temperature
- avoid excessive heat and humidity
- tamper evident sealed packets
- do not use any opened or torn packets

## **INACTIVE INGREDIENT**

### ***Inactive ingredients***

corn starch, hypromellose, polyethylene glycol, propylene glycol

## **QUESTIONS**

***Questions or comments?*** (888) 602-0288

## **Principal Display Panel - 250 Tablet Box Label**

**2501**

5 GRAIN

ASPIRIN

Pull to Open

For Temporary Relief of Minor Aches and Pains

The Provision

First Aid

Line™

BY WILDMAN

250 Tablets


2 Per Packet

# 5 GRAIN ASPIRIN

# 5 GRAIN ASPIRIN



*For Temporary  
Relief of  
Minor Aches  
and Pains*



250 Tablets  
2 Per Packet



*For Temporary Relief  
of Minor Aches  
and Pains*



250 Tablets

2 Per Packet

Manufactured for  
Provision by Wildman  
Warsaw, IN 46580



*For Temporary  
Relief of  
Minor Aches  
and Pains*



# 5 GRAIN ASPIRIN

<b>Drug Facts</b>		<b>Drug Facts (continued)</b>	
<b>Active ingredient (in each tablet)</b> <b>Purpose</b> Aspirin 325 mg (NSAID)*.....Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug		<b>Stop use and ask a doctor if</b> ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ■ feel faint    ■ vomit blood    ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever lasts more than 3 days ■ if ringing in the ears or loss of hearing occurs ■ redness or swelling is present ■ new symptoms appear	
<b>Uses</b> temporarily relieves ■ minor pain of arthritis    ■ headache    ■ muscle pain ■ toothache                    ■ pain and fever of colds ■ menstrual pain		<b>If pregnant or breast-feeding</b> , ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  <b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.	
<b>Warnings</b> Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives    ■ facial swelling    ■ asthma (wheezing)    ■ shock If an allergic reaction occurs, stop use and seek medical help right away.  Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcohol drinks every day while using this product ■ take more or for a longer time than directed  Do not use if you are allergic to aspirin or to any other pain reliever/fever reducer		<b>Directions</b> ■ drink a full glass of water with each dose	
<b>Ask a doctor before use if</b> ■ stomach bleeding warning applies to you ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma  Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout    ■ diabetes    ■ arthritis		<b>Adults and children: (12 years and older)</b> Take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours.  <b>Children under 12 years:</b> Consult a doctor	
		<b>Other information</b> ■ store at room temperature ■ avoid excessive heat and humidity ■ tamper evident sealed packets ■ do not use any opened or torn packets	
		<b>Inactive ingredients</b> corn starch, hypromellose, polyethylene glycol, propylene glycol	
		<b>Questions or comments?</b> (888) 602-0288 Manufactured for: Provision by Wildman Warsaw, IN 46580	

# 5 GRAIN ASPIRIN



Pull to Open

For Temporary Relief of Minor Aches and Pains



Retain carton for complete product information

250 Tablets

2 Per Packet

## ASPIRIN

aspirin tablet, coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:84269-2501
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

### Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

### Product Characteristics

<b>Color</b>	white (white)	<b>Score</b>	no score
<b>Shape</b>	ROUND (ROUND)	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	FR21
<b>Contains</b>			

<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84269-2501-2	125 in 1 BOX	07/01/2024	
1		2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:84269-2501-1	50 in 1 BOX	07/01/2024	
2		2 in 1 PACKET; Type 0: Not a Combination Product		

<b>Marketing Information</b>			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/01/2024	

**Labeler** - Wildman Business Group (016677338)

<b>Establishment</b>			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	MANUFACTURE(84269-2501)

<b>Establishment</b>			
Name	Address	ID/FEI	Business Operations
Medique Products		086911794	PACK(84269-2501)

<b>Establishment</b>			
Name	Address	ID/FEI	Business Operations
Prestige Packaging		080667761	PACK(84269-2501)

Revised: 7/2024

Wildman Business Group