

IBUPROFEN- ibuprofen tablet, coated
P & L Development, LLC

Drug Facts

Active ingredient (in each tablet)

Ibuprofen USP,
200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - minor pain of arthritis
 - toothache
 - backache
 - the common cold
 - menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - slurred speech
 - leg swelling
 - trouble breathing
 - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area

- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and older:
 - take 1 tablet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 tablet, 2 tablets may be used
 - do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store between 20°-25°C (68°-77°F). Avoid excessive heat above 40°C (104°F)

Inactive ingredients

colloidal silicon dioxide, corn starch, iron oxide red, macrogel/PEG, polyvinyl alcohol-part hydrolyzed, povidone K30, pregelatinized starch, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

†Compare to the active ingredient in Advil®

ibuprofen

tablets USP 200 mg

see new warnings

pain reliever / fever reducer (**NSAID**)

coated tablets

†This product is not manufactured or distributed by Pfizer Consumer Healthcare,

distributor of Advil®.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS
BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT
INFORMATION.**

Distributed by:

PL Developments

200 Hicks Street

Westbury, NY 11590

Product Label

Exp. Date: _____
 Lot No.: _____
 P.L.D.-A4468 F3004361 Product of India



Distributed by:
PL Developments
 200 Hicks Street
 Westbury, NY 11590

Drug Facts (continued)

- chest pain ■ slurred speech ■ leg swelling
- trouble breathing ■ weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

if pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions ■ do not take more than directed

- the smallest effective dose should be used
- adults and children 12 years and older:
- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- Store between 20° to 25°C (68° to 77°F). Avoid excessive heat above 40° (104°F).

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer

- right before or after heart surgery

Ask a doctor before use if ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ have bloody or black stools ■ feel faint ■ vomit blood

- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:

Drug Facts

Active ingredient (in each tablet) **Purpose**

Ibuprofen USP. 200 mg (NSAID)*Pain reliever/fever reducer
 *nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to: ■ headache ■ muscular aches ■ minor pain of arthritis ■ toothache ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives ■ facial swelling ■ asthma (wheezing)
- shock ■ skin reddening ■ rash ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you: ■ are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Drug Facts (continued)

Inactive ingredients colloidal silicon dioxide, corn starch, iron oxide red, macrogol/PEG, polyvinyl alcohol-part, hydrolyzed, povidone K30, pregelatinized starch, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments?
 Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil®.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.



Compare to active ingredient in Advil®
 NDC 59726-568-24

ibuprofen
 tablets USP 200 mg

see new warnings

pain reliever/
 fever reducer (NSAID)

24 coated tablets



READYinCASE Ibuprofen 200 mg Brown Tablets

IBUPROFEN

ibuprofen tablet, coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:59726-568
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE K30 (UNII: U725QWY32X)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
TALC (UNII: 7SEV7J4R1U)	

Product Characteristics

Color	brown	Score	no score
Shape	ROUND	Size	9mm
Flavor		Imprint Code	G2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59726-568-40	1 in 1 BOX	03/31/2017	04/26/2024
1		40 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:59726-568-24	1 in 1 BOX	03/31/2017	04/26/2024
2		24 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079174	03/31/2017	04/26/2024

Labeler - P & L Development, LLC (800014821)

