

**DEXTROSE- dextrose monohydrate injection, solution**  
**ICU Medical Inc.**

-----  
**70% Dextrose Injection, USP**  
Concentrated Dextrose in Water

**Pharmacy Bulk Package — Not For Direct Infusion.**  
**FOR USE ONLY WITH AUTOMATED COMPOUNDING DEVICES.**

**NOTE: These solutions are hypertonic - see *WARNINGS* and *PRECAUTIONS*.**

**Flexible Plastic Containers**

Rx

only

***DESCRIPTION***

70% Dextrose Injection, USP (concentrated dextrose in water) is a sterile, nonpyrogenic, hypertonic solution of Dextrose, USP in water for injection for intravenous administration after appropriate admixture or dilution.

The Pharmacy Bulk Package is a sterile dosage form which contains multiple single doses for use only in a pharmacy bulk admixture program.

The content and physical characteristics of the solutions are as follows:

<b>Solution Characteristics</b>	<b>70% Dextrose Injection, USP</b>
pH	4.0
pH range	3.2 - 6.5
Osmolarity (mOsmol/L) (calc.)	3532
Specific Gravity	1.236
Grams Dextrose/100 mL	70
kcal/100 mL*	238
Fill volume (mL)	2000

\* Caloric value calculated on the basis of 3.4 kcal/g of dextrose, hydrous.


The solutions contain no bacteriostat, antimicrobial agent or added buffer and are intended only for use as a single-dose injection following admixture or dilution.

This Pharmacy Bulk Package is intended only for use in the preparation of sterile, intravenous nutrient admixtures using automated compounding devices. The flexible plastic container is fabricated from a specially formulated polyvinylchloride. Water can permeate from inside the container into the overwrap but not in amounts sufficient to affect the solution significantly. Solutions in contact with the plastic container may leach out certain chemical components from the plastic in very small amounts; however, biological testing was supportive of the safety of the plastic container materials. Exposure to temperatures above 25°C/77°F during transport and storage will lead to

minor losses in moisture content. Higher temperatures lead to greater losses. It is unlikely that these minor losses will lead to clinically significant changes within the expiration period.

Dextrose Injection, USP is a parenteral fluid and nutrient replenisher.

Dextrose, USP is chemically designated D-glucose, monohydrate ( $C_6H_{12}O_6 \cdot H_2O$ ), a hexose sugar freely soluble in water. It has the following structural formula:



Water for Injection, USP is chemically designated H<sub>2</sub>O.

### **CLINICAL PHARMACOLOGY**

When administered intravenously, solutions containing carbohydrate in the form of dextrose restore blood glucose levels and provide calories. Carbohydrate in the form of dextrose may aid in minimizing liver glycogen depletion and exerts a protein sparing action. Dextrose injection undergoes oxidation to carbon dioxide and water.

Water is an essential constituent of all body tissues and accounts for approximately 70% of total body weight. Average normal adult daily requirement ranges from two to three liters (1.0 to 1.5 liters each for insensible water loss by perspiration and urine production).

Water balance is maintained by various regulatory mechanisms. Water distribution depends primarily on the concentration of electrolytes in the body compartments, and sodium (Na<sup>+</sup>) plays a major role in maintaining physiologic equilibrium.

### **INDICATIONS AND USAGE**

70% Dextrose Injection, USP (concentrated dextrose in water) in Pharmacy Bulk Packages is indicated for use with automated compounding devices for preparing intravenous nutritional admixtures in the pharmacy.

### **CONTRAINDICATIONS**

A concentrated dextrose solution should not be used when intracranial or intraspinal hemorrhage is present nor in the presence of delirium tremens if the patient is already dehydrated.

Dextrose injection without electrolytes should not be administered simultaneously with blood through the same infusion set because of the possibility that pseudoagglutination of red cells may occur.

## **WARNINGS**

Concentrated dextrose in water should be administered only after suitable dilution. Hypertonic dextrose solutions should be given slowly. Significant hyperglycemia and possible hyperosmolar syndrome may result from too rapid administration. The physician should be aware of the symptoms of hyperosmolar syndrome, such as mental confusion and loss of consciousness, especially in patients with chronic uremia and those with known carbohydrate intolerance.

The intravenous administration of these solutions can cause fluid and/or solute overloading resulting in dilution of serum electrolyte concentrations, overhydration, congested states or pulmonary edema.

**WARNING:** This product contains aluminum that may be toxic. Aluminum may reach toxic levels with prolonged parenteral administration if kidney function is impaired. Premature neonates are particularly at risk because their kidneys are immature, and they require large amounts of calcium and phosphate solutions, which contain aluminum.

Research indicates that patients with impaired kidney function, including premature neonates, who receive parenteral levels of aluminum at greater than 4 to 5 mcg/kg/day accumulate aluminum at levels associated with central nervous system and bone toxicity. Tissue loading may occur at even lower rates of administration.

### **FOR PERIPHERAL VEIN ADMINISTRATION**

Hypertonic dextrose solutions (above 5% concentration) should be given slowly, preferably through a small bore needle into a large vein, to minimize venous irritation.

### **FOR CENTRAL VENOUS ADMINISTRATION**

Concentrated dextrose should be administered via central vein after appropriate admixture or dilution when required.

## **PRECAUTIONS**

Electrolyte deficits, particularly in serum potassium and phosphate, may occur during prolonged use of concentrated dextrose solutions. Blood electrolyte monitoring is essential, and fluid and electrolyte imbalances should be corrected. Essential vitamins and minerals also should be provided as needed.

To minimize hyperglycemia and consequent glycosuria, it is desirable to monitor blood and urine glucose and if necessary, add insulin. When concentrated dextrose infusion is abruptly withdrawn, it is advisable to follow with the administration of 5% or 10% dextrose to avoid rebound hypoglycemia.

Aseptic technique is essential with the use of sterile preparations for compounding nutritional admixtures. Discard container within 4 hours of entering closure.

Solutions containing dextrose should be used with caution in patients with known subclinical or overt diabetes mellitus.

Care should be exercised to insure that the needle (or catheter) is well within the lumen of the vein and that extravasation does not occur.

Concentrated dextrose solutions should not be administered subcutaneously or intramuscularly.

Do not administer unless solution is clear and container is undamaged.

**Pregnancy Category C:** Animal reproduction studies have not been conducted with dextrose. It is also not known whether dextrose can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Dextrose should be given to a pregnant woman only if clearly needed.

### **Pediatric Use**

The safety and effectiveness in the pediatric population are based on the similarity of the clinical conditions of the pediatric and adult populations. In neonates or very small infants the volume of fluid may affect fluid and electrolyte balance. Caution should be exercised with low birth weight premature neonates, who are receiving dextrose concentrations of 10% or greater, as they are most susceptible to glucose intolerance and hyperglycemia. Frequent monitoring of serum glucose concentrations is required when dextrose is prescribed to pediatric patients, particularly neonates and low birth weight infants.

In very low birth weight infants, excessive or rapid administration of dextrose injection may result in increased serum osmolality and possible intracerebral hemorrhage.

### **ADVERSE REACTIONS**

Hyperosmolar syndrome, resulting from excessively rapid administration of concentrated dextrose may cause hypovolemia, dehydration, mental confusion and/or loss of consciousness.

Reactions which may occur because of the solution or the technique of administration include febrile response, infection at the site of injection, venous thrombosis or phlebitis extending from the site of injection, extravasation and hypervolemia.

If an adverse reaction does occur, discontinue the infusion, evaluate the patient, institute appropriate therapeutic countermeasures and save the remainder of the fluid for examination if deemed necessary.

### **OVERDOSAGE**

In the event of overhydration or solute overload during therapy, re-evaluate the patient and institute appropriate corrective measures. (See **WARNINGS** and **PRECAUTIONS**).

### **DOSAGE AND ADMINISTRATION**

Concentrated Dextrose in Water is administered by slow intravenous infusion (a) after admixture with amino acid solutions or (b) after dilution with other compatible IV fluids. Dosage should be adjusted to meet the requirements of each individual patient.

70% Dextrose Injection, USP in the 2000 mL flexible Pharmacy Bulk Package is designed for use with automated compounding devices for preparing intravenous nutritional admixtures. Dosages will be in accordance with the recommendation of the prescribing physician. 70% Dextrose Injection, USP are not intended for direct infusion. Admixtures

should be made by, or under the direction of, a pharmacist using strict aseptic technique under a laminar flow hood. Compounded admixtures may be stored under refrigeration for up to 24 hours. Administration of admixtures should be completed within 24 hours after removal from refrigeration.

The maximum rate at which dextrose can be infused without producing glycosuria is 0.5 g/kg of body weight/hr. About 95% of the dextrose is retained when infused at a rate of 0.8 g/kg/hr.

Clinical evaluation and periodic laboratory determinations are necessary to monitor changes in fluid balance, electrolyte concentrations, and acid base balance during prolonged parenteral therapy or whenever the condition of the patient warrants such evaluation.

As reported in the literature, the dosage and constant infusion rate of intravenous dextrose must be selected with caution in pediatric patients, particularly neonates and low birth weight infants, because of the increased risk of hyperglycemia/hypoglycemia.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. (See **PRECAUTIONS.**)

### **Drug Interactions**

Additives may be incompatible with the fluid withdrawn from this container. Consult with pharmacist, if available. When compounding admixtures, use aseptic technique, mix thoroughly and do not store.

Some opacity of the plastic due to moisture absorption during sterilization process may be observed. This is normal and does not affect the solution quality or safety. The opacity will diminish gradually.

### **Recommended Directions for Use of the Pharmacy Bulk Package Use Aseptic Technique**

1. During use, container must be stored, and all manipulations performed, in an appropriate laminar flow hood.
2. Remove cover from outlet port at bottom of container.
3. Insert piercing pin of transfer set and suspend unit in a laminar flow hood. Insertion of a piercing pin into the outlet port should be performed only once in a Pharmacy Bulk Package solution. Once the outlet site has been entered, the withdrawal of container contents should be completed promptly in one continuous operation. Should this not be possible, a maximum time of 4 hours from transfer set pin or implement insertion is permitted to complete fluid transfer operations; i.e., discard container no later than 4 hours after initial closure puncture.
4. Sequentially dispense aliquots of 70% Dextrose Injection, USP into IV containers using appropriate transfer set. During fluid transfer operations, the Pharmacy Bulk Package should be maintained under the storage conditions recommended in the labeling.

### **HOW SUPPLIED**

70% Dextrose Injection, USP are supplied as follows:

<b>NDC No.</b>	<b>Container</b>	<b>Concentration</b>	<b>Fill</b>
0409-7120-07	Flexible Pharmacy Bulk Package	70%	2000 mL
0990-7120-07			

ICU Medical is transitioning NDC codes from the "0409" to a "0990" labeler code. Both NDC codes are expected to be in the market for a period of time.

Store at 20 to 25°C (68 to 77°F). [See USP Controlled Room Temperature.] Protect from freezing.

Revised: June 2018

EN-4656

ICU Medical, Inc., Lake Forest, Illinois, 60045, USA

### **PRINCIPAL DISPLAY PANEL - 70 g Bag Label**

2000 mL

NDC 0990-7120-07

DEXTROSE

INJECTION,

USP

70%

Pharmacy Bulk Package —

Not For Direct Infusion.

MUST BE DILUTED.

EACH 100 mL CONTAINS DEXTROSE, HYDROUS, USP  
70 g IN WATER FOR INJECTION.

HYPERTONIC OSMOLARITY 3532 mOsmol/LITER (calc).

pH 4.0 (3.2 to 6.5) SPECIFIC GRAVITY 1.236

STERILE, NONPYROGENIC.

DOSAGE AND ADMINISTRATION: SEE PACKAGE  
INSERT.

CAUTION: DO NOT USE UNLESS SOLUTION IS CLEAR,  
CLOSURE IS INTACT, AND CONTAINER IS UNDAMAGED.

CHECK FOR MINUTE LEAKS BY SQUEEZING FIRMLY.

IF LEAKS ARE FOUND DISCARD CONTAINER AND

CONTENTS AS STERILITY MAY BE IMPAIRED. WITHIN

4 HOURS AFTER INITIAL ENTRY DISCARD CONTAINER

AND UNUSED CONTENTS.

DATE ENTERED:

TIME OF ENTRY:

STORE AT 20 TO 25°C (68 TO 77°F). [SEE USP  
CONTROLLED ROOM TEMPERATURE.] PROTECT FROM

FREEZING.

THIS PRODUCT CONTAINS NO MORE THAN 25 mcg/L  
OF ALUMINUM.

RX ONLY

70%

3

V

CONTAINS DEHP

IM-4368

ICU Medical, Inc., Lake Forest, Illinois, 60045, USA  
icumedical

2000 mL

NDC 0990-7120-07

# DEXTROSE INJECTION, USP



**Pharmacy Bulk Package —  
Not For Direct Infusion.  
MUST BE DILUTED.**

EACH 100 mL CONTAINS DEXTROSE, HYDROUS, USP  
70 g IN WATER FOR INJECTION.

HYPERTONIC OSMOLARITY 3532 mOsmol/LITER (calc).  
pH 4.0 (3.2 to 6.5) SPECIFIC GRAVITY 1.236  
STERILE, NONPYROGENIC.

**DOSAGE AND ADMINISTRATION:** SEE PACKAGE  
INSERT.

**CAUTION:** DO NOT USE UNLESS SOLUTION IS CLEAR,  
CLOSURE IS INTACT, AND CONTAINER IS UNDAMAGED.  
CHECK FOR MINUTE LEAKS BY SQUEEZING FIRMLY.  
IF LEAKS ARE FOUND DISCARD CONTAINER AND  
CONTENTS AS STERILITY MAY BE IMPAIRED. **WITHIN  
4 HOURS AFTER INITIAL ENTRY DISCARD CONTAINER  
AND UNUSED CONTENTS.**

DATE ENTERED: \_\_\_\_\_

TIME OF ENTRY: \_\_\_\_\_

STORE AT 20 TO 25°C (68 TO 77°F). [SEE USP  
CONTROLLED ROOM TEMPERATURE.] PROTECT FROM  
FREEZING.

THIS PRODUCT CONTAINS NO MORE THAN 25 mcg/L  
OF ALUMINUM.

Rx ONLY

1750 —  
1500 —  
1250 —  
1000 —  
750 —  
500 —  
250 —



CONTAINS DEHP

IM-4368



ICU Medical, Inc., Lake Forest, Illinois, 60045, USA



**icumedical**

**PRINCIPAL DISPLAY PANEL - 70 g Bag Overwrap Label**

TO OPEN — TEAR AT NOTCH

The overwrap is a moisture and oxygen barrier. Do not remove unit from overwrap until ready for use. Visually inspect overwrap for tears or holes. Use unit promptly when pouch is opened. Store at 20 to 25°C (68 to 77°F). [See USP Controlled Room Temperature.] Protect from freezing. See insert. After removing the overwrap, check for minute leaks by squeezing container firmly. If leaks are found, discard solution as sterility may be impaired.

COLOR VARIATION IN THE DEXTROSE INJECTION FROM PALE YELLOW TO YELLOW IS NORMAL AND DOES NOT ALTER EFFICACY.

Rx only

WR-0523

ICU Medical, Inc., Lake Forest, Illinois, 60045, USA

icumedical



(01)00309907120072



## TO OPEN — TEAR AT NOTCH

The overwrap is a moisture and oxygen barrier. Do not remove unit from overwrap until ready for use. Visually inspect overwrap for tears or holes. Use unit promptly when pouch is opened. Store at 20 to 25°C (68 to 77°F). [See USP Controlled Room Temperature.] Protect from freezing. See insert. After removing the overwrap, check for minute leaks by squeezing container firmly. If leaks are found, discard solution as sterility may be impaired.

**COLOR VARIATION IN THE DEXTROSE INJECTION FROM PALE YELLOW TO YELLOW IS NORMAL AND DOES NOT ALTER EFFICACY.**

**Rx** only

WR-0523

ICU Medical, Inc., Lake Forest, Illinois, 60045, USA

**icu**medical



### DEXTROSE

dextrose monohydrate injection, solution

#### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0990-7120
<b>Route of Administration</b>	INTRAVENOUS		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>DEXTROSE MONOHYDRATE</b> (UNII: LX22YL083G) (ANHYDROUS DEXTROSE - UNII:5SLOG7R0OK)	DEXTROSE MONOHYDRATE	70 g in 100 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>WATER</b> (UNII: 059QF0KO0R)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0990-7120-07	6 in 1 CASE	07/01/2019	
1		1 in 1 POUCH		
1		2000 mL in 1 BAG; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA019893	07/01/2019	

**Labeler** - ICU Medical Inc. (118380146)

Revised: 10/2025

ICU Medical Inc.