

## **LIDOPRO- lidocaine and menthol patch**

### **Clinic Pharma**

-----

### **LidoPro Patch**

#### **Active ingredients (in each patch)**

Lidocaine 4%

#### **Purpose**

External Anesthetic

#### **Active ingredients (in each patch)**

Menthol 1%

#### **Purpose**

External Analgesic

#### **Use**

Temporarily relieves minor pain

#### **Warnings**

For external use only

#### **Do not use**

- more than one patch on your body at a time
- on cut, irritated or swollen skin
- on puncture wounds
- for more than one week without consulting a doctor
- if you are allergic to any active or inactive ingredients
- if pouch is damaged or opened

#### **When using this**

- use only as directed
- read and follow all directions and warnings on this carton
- do not allow contact with the eyes
- do not use at the same time as other topical analgesics
- do not bandage tightly or apply local heat (such as heating pads) to the area of use
- do not microwave
- dispose of used patch in manner that always keeps product away from children and

pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

**Stop use and consult a doctor if**

- condition worsens
- redness is present
- irritation develops
- symptoms persist for more than 7 days or clear up and occur again within a few days
- you experience signs of skin injury, such as pain, swelling, or blistering where the product was applied

**If pregnant or breastfeeding,**

ask a health professional before use.

**Keep out of reach of children and pets.**

If swallowed, get medical help or contact a Poison Control Center 1-800-222-1222 right away.

**Directions**

**Adults and children 12 years of age and over:** clean and dry affected area, free of lotions, ointments, and creams. Carefully remove backing from patch starting at a corner. Apply sticky side of patch to affected area. Do not use more than one patch in a 12-hour period. Maximum 2 patches per day. Discard patch after single use.

**Children under 12 years of age:** consult a physician.

**Other information**

Store in a clean, dry place outside of direct sunlight. Protect from excessive moisture.

**Inactive ingredients**

dihydroxyaluminum aminoacetate, edetate disodium, glycerin, hydroxyacetophenone, kaolin, mineral oil, polyacrylic acid, polysorbate 80, povidone K90, propylene glycol, sodium polyacrylate, tartaric acid, titanium dioxide, water

**Questions or comments?**

Call (800) 224-2048 or email [info@clinicpharma.com](mailto:info@clinicpharma.com)

**Principal Display Panel**

NDC 83881-403-01

# LidoPro<sup>®</sup> Lidocaine 4% • Menthol 1% patch

## Pain Relieving Patch

- ✓ Deep Penetrating
- ✓ Long-lasting
- ✓ Soothing

1 PATCH

10 cm x 14 cm  
(3.9 in x 5.5 in)

### Drug Facts

| Active ingredients (in each patch) | Purpose             |
|------------------------------------|---------------------|
| Lidocaine 4%                       | External Anesthetic |
| Menthol 1%                         | External Analgesic  |

**Use** Temporarily relieves minor pain

#### Warnings

**For external use only**

**Do not use** • more than one patch on your body at a time • on cut, irritated or swollen skin • on puncture wounds • for more than one week without consulting a doctor • if you are allergic to any active or inactive ingredients • if pouch is damaged or opened

**When using this product** • use only as directed • read and follow all directions and warnings on the carton • do not allow contact with the eyes • do not use at the same time as other external analgesics • do not bandage tightly or apply local heat (such as heating pads) to the area of use • do not microwave • dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

**Stop use and consult a doctor if** • condition worsens • redness is present • irritation develops • symptoms persist for more than 7 days or clear up and occur again within a few days • you experience signs of skin injury, such as pain, swelling, or blistering where the product was applied

**If pregnant or breastfeeding**, ask a health professional before use.

**Keep out of reach of children and pets.** If swallowed, get medical help or contact a Poison Control Center 1-800-222-1222 right away.

#### Directions

**Adults 18 years of age and over:** clean and dry affected area, free of lotions, ointments, and creams. Carefully remove backing from patch starting at a corner. Apply sticky side of patch to affected area. Do not use more than one patch in a 12-hour period. Maximum 2 patches per day. Discard patch after single use.  
**Children under 18 years of age:** consult a physician.

#### Other information

Store in a clean, dry place outside of direct sunlight. Protect from excessive moisture.

#### Inactive ingredients

dihydroxyaluminum aminoacetate, edetate disodium, glycerin, hydroxyacetophenone, kaolin, mineral oil, polyacrylic acid, polysorbate 80, povidone K90, propylene glycol, sodium polyacrylate, tartaric acid, titanium dioxide, water

**Questions or comments?** Call (800) 224-2048 or email [info@clinicpharma.com](mailto:info@clinicpharma.com)

AF2.03



Manufactured For:  
**Clinic  
Pharma**  
Las Vegas, NV 89121  
[info@clinicpharma.com](mailto:info@clinicpharma.com)

Formulated and  
Designed in Nevada  
Patent Pending

LOT:  
EXP:

**LidoPro<sup>®</sup>**  
Lidocaine 4% • Menthol 1%  
*patch*

30 PATCHES

NDC 83881-403-30

**LidoPro<sup>®</sup>**  
Lidocaine 4% • Menthol 1%  
*patch*

**Pain Relieving Patch**

- ✓ Deep Penetrating
- ✓ Long-lasting
- ✓ Soothing

30 PATCHES

10 cm x 14 cm  
(3.9 in x 5.5 in)

**Drug Facts**

| Active ingredients (in each patch) | Purpose             |
|------------------------------------|---------------------|
| Lidocaine 4%                       | External Anesthetic |
| Menthol 1%                         | External Analgesic  |

**Use** Temporarily relieve minor pain

**Warnings**

For external use only

Do not use • more than one patch on your body at a time • on wet, irritated or swollen skin • on puncture wounds • for more than one week without consulting a doctor • if you are allergic to any active or inactive ingredients • if patch is damaged or opened

When using this product • use only as directed • read and follow all directions and warnings on this carton • do not allow contact with the eyes • do not use at the same time as other external analgesics • do not backage tightly or apply local heat (such as heating pad) to the area of use • do not microwave • dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

Stop use and consult a doctor if • condition worsens • redness is present • irritation develops • symptoms persist for more than 7 days or clear up and return again within a few days • you experience signs of skin injury, such as pain, swelling, or blistering where the product was applied.

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children and pets. If swallowed, get medical help or contact a Poison Control Center 1-800-222-1222 right away.

**Directions**

Adults 18 years of age and over: clean and dry affected area, free of lotions, ointments, and creams. Carefully remove backing from patch starting at a corner. Apply sticky side of patch to affected area. Do not use more than one patch in a 24-hour period. Maximum 2 patches per day. Discard patch after single use. Children under 18 years of age: consult a physician.

**Other information**

Store in a clean, dry place outside of direct sunlight. Protect from excessive moisture.

**Inactive ingredients**

glyceryltrimonitrate, acetaminophen, edetate disodium, glycerin, hydroxyacetophenone, kaolin, mineral oil, polyacrylic acid, polyethylene glycol, polyethylene glycol, sodium polyacrylate, tartaric acid, titanium dioxide, water.

**Questions or comments?**

Call 800-224-2048 or email: info@lidoapharma.com



Manufactured for  
**clinic  
Pharma**  
Las Vegas, NV 89121  
info@lidoapharma.com

Formulated and  
Designed in Nevada  
Heart Healthy

LOT:  
EXP:

Deep Penetrating Long-lasting Soothing™



## LIDOPRO

lidocaine and menthol patch

### Product Information

|                                |                |                           |               |
|--------------------------------|----------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:83881-403 |
| <b>Route of Administration</b> | TRANSDERMAL    |                           |               |

### Active Ingredient/Active Moiety

| Ingredient Name   | Basis of Strength | Strength      |
|---|-------------------|---------------|
| <b>LIDOCAINE</b> (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987) | LIDOCAINE         | 0.04 g in 1 g |
| <b>MENTHOL</b> (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)     | MENTHOL           | 0.01 g in 1 g |

### Inactive Ingredients

| Ingredient Name  | Strength |
|--|----------|
| <b>DIHYDROXYALUMINUM AMINOACETATE</b> (UNII: DO250MG0W6) |          |
| <b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)               |          |
| <b>GLYCERIN</b> (UNII: PDC6A3C0OX)                       |          |

|  |
|--|
| <b>HYDROXYACETOPHENONE</b> (UNII: G1L3HT4CMH)              |
| <b>KAOLIN</b> (UNII: 24H4NWX5CO)                           |
| <b>MINERAL OIL</b> (UNII: T5L8T28FGP)                      |
| <b>POLYACRYLIC ACID (250000 MW)</b> (UNII: 9G2MAD7J6W)     |
| <b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)                   |
| <b>POVIDONE K90</b> (UNII: RDH86HJV5Z)                     |
| <b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)                 |
| <b>SODIUM POLYACRYLATE (2500000 MW)</b> (UNII: 05I15JNI2J) |
| <b>TARTARIC ACID</b> (UNII: W4888I119H)                    |
| <b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)                 |
| <b>WATER</b> (UNII: 059QF0KO0R)                            |

### Packaging

| # | Item Code        | Package Description                                | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:83881-403-30 | 30 in 1 CARTON                                     | 03/11/2026           |                    |
| 1 |                  | 1 in 1 POUCH                                       |                      |                    |
| 1 |                  | 10 g in 1 PATCH; Type 0: Not a Combination Product |                      |                    |
| 2 | NDC:83881-403-15 | 15 in 1 CARTON                                     | 03/11/2026           |                    |
| 2 |                  | 1 in 1 POUCH                                       |                      |                    |
| 2 |                  | 10 g in 1 PATCH; Type 0: Not a Combination Product |                      |                    |
| 3 | NDC:83881-403-01 | 1 in 1 POUCH                                       | 03/11/2026           |                    |
| 3 |                  | 10 g in 1 PATCH; Type 0: Not a Combination Product |                      |                    |

### Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M017                                     | 03/11/2026           |                    |

**Labeler** - Clinic Pharma (119158469)

Revised: 3/2026

Clinic Pharma