

COLD MULTI-SYMP TOM NIGHTTIME- acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, and phenylephrine hydrochloride tablet, coated
Marc Glassman, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

MAR - 1139 - 2019-0901

Drug Facts

<i>Active ingredients (in each caplet)</i>	<i>Purpose</i>
Acetaminophen 325 mg	Pain reliever/fever reducer
Chlorpheniramine maleate 2 mg	Antihistamine
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these common cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - runny nose and sneezing
 - cough
 - sinus congestion and pressure
- helps clear nasal passages
- relieves cough to help you sleep
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache,

rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- high blood pressure
- diabetes
- heart disease
- thyroid disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)

adults and children 12 years and over	<ul style="list-style-type: none">take 2 caplets every 4 hoursswallow whole – do not crush, chew, or dissolvedo not take more than 10 caplets in 24 hours
children under 12 years	<ul style="list-style-type: none">ask a doctor

Other information

- store between 20-25C (68°-77°F) in a dry place
- retain carton for complete product information and warnings

Inactive ingredients

acesulfame potassium, colloidal silicon dioxide, croscarmellose sodium, FD&C blue #1 aluminum lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

PRINCIPAL DISPLAY PANEL

NDC 68998-139-02

Marc's

Compare to the active ingredients in Tylenol® Cold Multi-Symptom Nighttime
nighttime

Cold Max

Multi-Symptom

PAIN RELIEVER / FEVER REDUCER, COUGH SUPPRESSANT, NASAL DECONGESTANT,
ANTIHISTAMINE

Acetaminophen, Dextromethorphan HBr, Phenylephrine HCl, Chlorpheniramine Maleate

For Relief of:

Head + Body Aches

Fever + Sore Throat

Cough

Nasal Congestion

Runny Nose

Actual Size

24 COOL TASTE CAPLETS




Drug Facts (continued)
Other information
 magnesium stearate, microcrystalline cellulose, polyethylene

<p>Drug Facts (continued)</p> <p>Warnings</p> <p>■ breathing problem such as emphysema or chronic bronchitis</p> <p>■ Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin</p> <p>When using this product</p> <p>■ do not exceed recommended dosage</p> <p>■ excitability may occur, especially in children</p> <p>■ marked drowsiness may occur ■ avoid alcoholic drinks</p> <p>■ alcohol, sedatives, and tranquilizers may increase drowsiness</p> <p>■ be careful when driving a motor vehicle or operating machinery</p> <p>Stop use and ask a doctor if</p> <p>■ pain, nasal congestion, or cough gets worse or lasts more than 7 days</p> <p>■ fever gets worse or lasts more than 3 days</p> <p>■ redness or swelling is present ■ new symptoms occur</p> <p>■ cough comes back or occurs with rash or headache that lasts</p> <p>These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children.</p> <p>Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions</p> <p>■ do not take more than directed (see overdose warning)</p> <p>adults and children 12 years and over</p> <p>■ take 2 caplets every 4 hours</p> <p>■ swallow whole – do not crush, chew, or dissolve</p> <p>■ do not take more than 10 caplets in 24 hours</p> <p>children under 12 years</p> <p>■ ask a doctor</p>	<p>Drug Facts (continued)</p> <p>Warnings</p> <p>■ Liver warning: This product contains acetaminophen. (3250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours</p> <p>■ with other drugs containing acetaminophen</p> <p>■ 3 or more alcoholic drinks every day while using this product</p> <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:</p> <p>■ skin reddening ■ hives ■ rash</p> <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p>Do not use</p> <p>■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</p> <p>■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>■ if you have ever had an allergic reaction to this product or any of its ingredients</p> <p>Ask a doctor before use if you have ■ liver disease ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ glaucoma ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema</p> <p>■ cough that occurs with too much phlegm (mucus)</p>
<p>Drug Facts (continued)</p> <p>Uses ■ temporarily relieves these common cold/flu symptoms: ■ minor aches and pains</p> <p>■ headache ■ sore throat ■ nasal congestion</p> <p>■ runny nose and sneezing ■ cough</p> <p>■ sinus congestion and pressure</p> <p>■ helps clear nasal passages ■ temporarily reduces fever</p> <p>■ relieves cough to help you sleep</p>	<p>Drug Facts (continued)</p> <p>Active ingredients (in each caplet)</p> <p>Acetaminophen 325 mg... Pain reliever/fever reducer</p> <p>Chlorpheniramine maleate 2 mg... Antihistamine</p> <p>Dextromethorphan HBr 10 mg... Cough suppressant</p> <p>Phenylephrine HCl 5 mg... Nasal decongestant</p> <p>Purpose</p>

†This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol® Cold Multi-Symptom Nighttime.

DO NOT USE IF BLISTER UNITS ARE TORN OR BROKEN

Distributed by:
 Marc's Glassman, Inc.
 West 130th Street
 Cleveland, OH 44130



NDC 68998-139-02

†Compare to the active ingredients in Tylenol® Cold Multi-Symptom Nighttime



nighttime 

Cold Max

Multi-Symptom

PAIN RELIEVER / FEVER REDUCER, COUGH SUPPRESSANT, NASAL DECONGESTANT, ANTIHISTAMINE

NC

NC

NASAL DECONGESTANT, ANTIHISTAMINE

**Acetaminophen, Dextromethorphan HBr,
Phenylephrine HCl, Chlorpheniramine Maleate**

- For Relief of:
- Head + Body Aches
 - Fever + Sore Throat
 - Cough
 - Nasal Congestion
 - Runny Nose

Actual Size



24 COOL TASTE CAPLETS

N
C

COLD MULTI-SYMP TOM NIGHTTIME

acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, and phenylephrine hydrochloride tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68998-139
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
ALUMINUM OXIDE (UNII: LM26O6933)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	

STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	blue	Score	no score
Shape	OVAL	Size	17mm
Flavor	MINT	Imprint Code	AAA;1139
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68998-139-02	2 in 1 CARTON	01/27/2010	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	01/27/2010	

Labeler - Marc Glassman, Inc. (094487477)

Revised: 9/2019

Marc Glassman, Inc.