

**SENNA SYRUP- natural vegetable laxative liquid**  
**Method Pharmaceuticals**

-----  
**Senna Syrup**

**Active Ingredients**

Sennosides 8.8 mg

**Purpose**

Laxative

**Warning**

Do not use laxative products for longer than 1 week unless directed by a doctor.

**Uses**

- Relieves occasional constipation (irregularity)
- Generally causes bowel movement in 6 to 12 hours

**Ask a doctor before use if you**

- have stomach pain, nausea or vomiting
- notice a change in bowel habits that last over 2 weeks
- are **pregnant or breastfeeding.**

**Stop Use and ask a doctor**

if you have rectal bleeding or failure to have a bowel movement after use.

**Keep out of Reach of Children**

In case of accidental overdose, get medical help or contact a Poison Control Center immediately.

**Directions**

Shake well before using. Do not exceed maximum dose. Take at bedtime or as directed by a doctor.

**Store** at controlled room temperature 20-25°C(68-77°F). Excursions permitted 15-30° (59-86°F).

**Tamper Evident:** Do not use if foil seal over bottle opening is torn, broken or missing.

## Inactive Ingredients

Chocolate flavor, methylparaben, propylene glycol, propylparaben, purified water and sucrose

## Dosage

<b>Age</b>	<b>Usual Dose</b>	<b>Maximum Dose</b>
Adults and Children 12 years and older	2-3 teaspoons once a day	3 teaspoons twice a day
6-12 years	1 to 1 1/2 teaspoons once a day	1 1/2 teaspoons twice a day
2-6 years	1/2 to 3/4 teaspoon once a day	3/4 teaspoons twice a day
under 2 years	Consult a doctor	

## Senna Syrup

Manufactured for :

Method Pharmaceuticals, LLC

419 Bank St Ste 100 Southlake, TX 76092

Rev 01.0721



NDC 58657-518-08

### Senna Syrup

Natural Vegetable Laxative

**8.8 mg/5 mL**

Alcohol Free • Dye Free

8 fl. oz (237 mL)

Drug Facts		Drug Facts (continued)	
Active Ingredient (per 5 mL teaspoon) Purpose	Age	Usual Dose	Maximum Dose
Sennosides 8.8 mg.....Laxatives	Adults and children 12 years and older	2-3 teaspoons once a day	3 teaspoons twice a day
<b>Uses</b> • Relieves occasional constipation (irregularity) • Generally causes bowel movement in 6 to 12 hours	6 to 12 years	1 to 1 1/2 teaspoons once a day	1 1/2 teaspoons twice a day
	2 to 6 years	1/2 to 3/4 teaspoon once a day	3/4 teaspoon once a day
	Under 2 years	Consult a doctor	
<b>Warnings</b> Do not use laxative products for longer than 1 week unless directed by a doctor. Ask a doctor before use if you • have stomach pain, nausea or vomiting • notice a change in bowel habits that last over 2 weeks • are pregnant or breastfeeding. Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center immediately.	<b>Other Information</b> Store at controlled room temperature 20-25°C (68-77°F). Excursions permitted 15-30° (59-86°F). Tamper Evident: Do not use if foil seal over bottle opening is torn, broken or missing.		
	<b>Inactive Ingredients</b> Chocolate flavor, methylparaben, propylene glycol, propylparaben, purified water and sucrose		
<b>Directions</b> Shake well before using. Do not exceed maximum dose. Take at bedtime or as directed by a doctor.	<b>Manufactured for :</b> Method Pharmaceuticals, LLC 419 Bank St Ste 100 Southlake, TX 76092 Rev 01.0721		

0.875" x 1.125"  
UNVARNISHED AREA



N 58657 51808 0

## SENNAS SYRUP

natural vegetable laxative liquid

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:58657-518
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>SENNOSIDES</b> (UNII: 3FYP5M0IJX) (SENNOSIDES - UNII:3FYP5M0IJX)	SENNOSIDES	8.8 mg in 5 mL

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>WATER</b> (UNII: 059QF0K00R)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:58657-518-08	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/28/2021	

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M007	09/28/2021	

**Labeler** - Method Pharmaceuticals (060216698)

Revised: 10/2025

Method Pharmaceuticals