MIDOL COMPLETE- acetaminophen, caffeine, and pyrilamine maleate tablet Bayer HealthCare LLC.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Midol ® Complete

Drug Facts

Active ingredients (in each caplet)	Purpose
Acetaminophen 500 mg	Pain reliever
Caffeine 60 mg	Diuretic
Pyrilamine maleate 15 mg	Antihistamine

Uses

for the temporary relief of these symptoms associated with menstrual periods:

- cramps
- bloating
- water-weight gain
- headache
- backache
- muscle aches
- fatigue

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- glaucoma
- difficulty in urination due to enlargement of the prostate gland
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- you may get drowsy
- avoid alcoholic drinks
- excitability may occur, especially in children
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- limit the use of caffeine-containing medications, foods, or beverages because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat. The recommended dose of this product contains about as much caffeine as a cup of coffee.

Stop use and ask a doctor if

- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than the recommended dose
 - adults and children 12 years and older:
 - take 2 caplets with water
 - repeat every 6 hours, as needed
 - do not exceed 6 caplets per day
- children under 12 years: consult a doctor

Other information

store at room temperature

Inactive ingredients

carnauba wax, croscarmellose sodium, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, propylene glycol, shellac, titanium dioxide, triacetin

Questions or comments?

1-800-331-4536 (Mon-Fri 9AM - 5PM EST) or www.midol.com

PRINCIPAL DISPLAY PANEL - 40 Caplet Bottle Carton

VALUE SIZE

Midol _® Complete

Acetaminophen/Caffeine/Pyrilamine Maleate Pain Reliever/Diuretic/Antihistamine

MAXIMUM STRENGTH

MULTI-SYMPTOM RELIEF OF

CRAMPS, BLOATING FATIGUE, BACKACHE & HEADACHE

40 CAPLETS

MIDOL COMPLETE

acetaminophen, caffeine, and pyrilamine maleate tablet

Product Informat	ion						
Product T ype		HUMAN OTC DRUG	Item Cod	Item Code (Source)		NDC:0280-8005	
Route of Administrat	tion	ORAL					
Active Ingredient	/Active Moi	ety					
	- C	redient Name			Strength	Strengt	
ACETAMINOPHEN (UN	NII: 362O9ITL9I	D) (ACETAMINOPHEN - UNII:	362O9ITL9D)	PHEN	500 mg		
CAFFEINE (UNII: 3G6 A	5W338E) (CAFF	EINE - UNII:3G6A5W338E)		CAFFEINE		60 mg	
PYRILAMINE MALEATE (UNII: R35D29L3ZA) (PYRILAMINE - UNII: HPE31709TL) PYRILAMINE MALEATE					15 mg		
Inactive Ingredie	nts						
		Ingredient Name			S	trength	
CARNAUBA WAX (UNI	II: R12CBM0EIZ)						
CROSCARMELLOSE	SODIUM (UNII:	M28OL1HH48)					
FD&C BLUE NO. 2 (UI	NII: L06K8R7DQ	PK)					
HYPROMELLOSES (U	NII: 3NXW29V3	WO)					
MAGNESIUM STEARA	TE (UNII: 70097	7M6I30)					
CELLULOSE, MICRO	CRYSTALLINE	(UNII: OP1R32D61U)					
PROPYLENE GLYCOI	L (UNII: 6DC9Q	167V3)					
SHELLAC (UNII: 46 N10	7B710)						
TITANIUM DIO XIDE (P)					
TRIACETIN (UNII: XHX	(3C3X673)						
	•						
Product Characte			-				
Color	white			Score		no score	
Shape 	OVAL (oble	ong)		Size		17mm	
Flavor			Imprint Code		MIDOL	MIDOL	
Contains							
Packaging							
# Item Code		Package Description		Marketing Start Da	te Marketi	ng End Dat	
1 NDC:0280-8005-16	2 in 1 CARTON			02/10/2014			
1	8 in 1 BLISTER	PACK; Type 0: Not a Combir	ation Product				
2 NDC:0280-8005-24	4 3 in 1 CARTON			02/10/2014			
	8 in 1 BLISTER PACK; Type 0: Not a Combination Product						
3 NDC:0280-8005-40	1 in 1 CARTON			02/10/2014			
		; Type 0: Not a Combination					

Marketing Information							
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date				
OTC monograph not final	part343	02/10/2014					

Labeler - Bayer HealthCare LLC. (112117283)

Revised: 11/2017

Bayer HealthCare LLC.