

**COMBINATION PACK TOPCARE HEALTH DAYTIME SEVERE AND NITETIME  
SEVERE COLD AND FLU RELIEF SOFTGELS - 24 CT- acetaminophen,  
dextromethorphan hydrobromide, guaifenesin, doxylamine succinate,  
phenylephrine hydrochloride**

**Topco Associates LLC**

---

**COMBINATION PACK TopCare® health DayTime *SEVERE* and NiteTime *SEVERE*  
Cold & Flu Relief SOFTGELS**

**DAYTIME SEVERE COLD AND FLU RELIEF**

***Drug Facts***

***Active ingredients (in each softgel)***

Acetaminophen 325 mg  
Dextromethorphan HBr 10 mg  
Guaifenesin 200 mg  
Phenylephrine HCl 5 mg

***Purposes***

Pain reliever/fever reducer  
Cough suppressant  
Expectorant  
Nasal decongestant

***Uses***

- temporarily relieves common cold/flu symptoms:
  - nasal congestion
  - sinus congestion & pressure
  - cough due to minor throat & bronchial irritation
  - minor aches & pains
  - head ache
  - fever
  - sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 softgels in 24 hours, which is the maximum daily amount for this product

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions.

Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease • heart disease • high blood pressure
- thyroid disease • diabetes • glaucoma
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin; taking sedatives or tranquilizers

**When using this product, do not use more than directed.**

### **Stop use and ask a doctor if**

- you get nervous, dizzy, or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present • new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

### **Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away.

### **Directions**

- take only as directed
- do not exceed 8 softgels per 24 hrs

adults & children 12 yrs and over	2 softgels with water every 4 hrs
children 4 to under	ask a doctor

12 yrs	ask a doctor
children under 4 yrs	do not use

### **Other information**

- store at controlled room temperature between 20-25°C (68-77°F)
- protect from light, heat, and moisture

### **Inactive ingredients**

FD&C Yellow No. 6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, sorbitol sorbitan solution, titanium dioxide

### **Questions or comments?**

Call **1-888-423-0139**

## **NITETIME SEVERE COLD AND FLU RELIEF**

### **Drug Facts**

#### **Active ingredients (in each softgel)**

Acetaminophen 325 mg  
Dextromethorphan HBr 10 mg  
Doxylamine succinate 6.25 mg  
Phenylephrine HCl 5 mg

### **Purposes**

Pain reliever/fever reducer  
Cough suppressant  
Antihistamine  
Nasal decongestant

### **Uses**

- temporarily relieves common cold/flu symptoms:
  - nasal congestion
  - sinus congestion & pressure
  - cough due to minor throat & bronchial irritation
  - minor aches & pains
  - headache
  - fever
  - sore throat
  - runny nose & sneezing
  - reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

## **Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 softgels in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions.

Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

## **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

## **Ask a doctor before use if you have**

- liver disease • heart disease • high blood pressure
- thyroid disease • diabetes • glaucoma
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or occurs with smoking, asthma, chronic bronchitis, or emphysema

## **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

## **When using this product**

### **• do not use more than directed**

- excitability may occur, especially in children
- marked drowsiness may occur • avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

## **Stop use and ask a doctor if**

- you get nervous, dizzy, or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.  
These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control

Center right away.

### **Directions**

- take only as directed
- do not exceed 8 softgels per 24 hours

adults & children 12 yrs and over	2 softgels with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

### **Other information**

- store at controlled room temperature between 20-25°C (68-77°F)
- protect from light, heat, and moisture

### **Inactive ingredients**

D&C Yellow No. 10, FD&C Blue No. 1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, sorbitol sorbitan solution, titanium dioxide

### **Questions or comments?**

Call **1-888-423-0139**

COMPARE TO VICKS® DAYQUIL™ SEVERE COLD & FLU LIQUICAPS™ ACTIVE INGREDIENTS\*

### **MULTI-SYMPOTOM RELIEF**

### **Antihistamine Free**

COMPARE TO VICKS® NYQUIL™ SEVERE COLD & FLU LIQUICAPS™ ACTIVE INGREDIENTS\*

\*These products are not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademarks Vicks® DayQuil™ & NyQuil™ LiquiCaps™.

### **READ AND KEEP OUTER CARTON FOR COMPLETE PRODUCT WARNINGS AND INFORMATION**

### **DO NOT TAKE THESE PRODUCTS AT THE SAME TIME**

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.**

DISTRIBUTED BY TOPCO ASSOCIATES LLC  
ITASCA, IL 60143 ©TOPCO STVA0825

topcare@topco.com www.topcarebrand.com  
PRODUCT OF INDIA

## QUALITY GUARANTEED

This Topcare® product is laboratory tested to guarantee its highest quality. Your total satisfaction is guaranteed.

## Packaging



READ AND KEEP OUTER CARTON FOR COMPLETE PRODUCT WARNINGS AND INFORMATION.

DO NOT TAKE THESE PRODUCTS AT THE SAME TIME

### NITETIME SEVERE COLD AND FLU RELIEF

### DAYTIME SEVERE COLD AND FLU RELIEF

#### Drug Facts

##### Active ingredients (in each softgel)

Acetaminophen 325 mg.....Pain reliever/fever reducer  
Dextromethorphan HBr 10 mg.....Cough suppressant  
Doxylamine succinate 6.25 mg.....Antihistamine  
Phenylephrine HCl 5 mg.....Nasal decongestant

##### Purposes

- temporarily relieves common cold/flu symptoms:
  - nasal congestion
  - sinus congestion & pressure
  - cough due to minor throat & bronchial irritation
  - cough to help you sleep
  - minor aches & pains
  - headache
  - fever
  - sore throat
  - runny nose & sneezing
  - reduces swelling of nasal passages
  - temporarily restores freer breathing through the nose
  - promotes nasal and/or sinus drainage

#### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver

#### Drug Facts

##### Active ingredients (in each softgel)

Acetaminophen 325 mg.....Pain reliever/fever reducer  
Dextromethorphan HBr 10 mg.....Cough suppressant  
Guaiifenesin 200 mg.....Expectorant  
Phenylephrine HCl 5 mg.....Nasal decongestant

##### Purposes

- temporarily relieves common cold/flu symptoms:
  - nasal congestion
  - sinus congestion & pressure
  - cough due to minor throat & bronchial irritation
  - minor aches & pains
  - headache
  - fever
  - sore throat
  - reduces swelling of nasal passages
  - temporarily restores freer breathing through the nose
  - promotes nasal and/or sinus drainage
  - helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make

LOT:  
EXP:

damage may occur if you take

- more than 8 softgels in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.

coughs more productive.

#### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 softgels in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

#### Drug Facts (continued)

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

##### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

##### Ask a doctor before use if you have

- liver disease ■ heart disease ■ high blood pressure
- thyroid disease ■ diabetes ■ glaucoma
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema

##### Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

##### When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur ■ avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

#### Drug Facts (continued)

■ alcohol, sedatives, and tranquilizers may increase drowsiness

##### Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away.

##### Directions

- take only as directed
- do not exceed 8 softgels per 24 hours

adults & children 12 yrs and over	2 softgels with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

##### Other information

- store at controlled room temperature 20–25°C (68–77°F)
- protect from light, heat, and moisture

#### Drug Facts (continued)

##### Inactive ingredients

D&C Yellow No. 10, FD&C Blue No. 1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, sorbitol sorbitan solution, titanium dioxide

#### Drug Facts (continued)

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

##### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

##### Ask a doctor before use if you have

- liver disease ■ heart disease ■ high blood pressure
- thyroid disease ■ diabetes ■ glaucoma
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin; taking sedatives or tranquilizers**

##### When using this product, do not use more than directed.

##### Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days

#### Drug Facts (continued)

- fever gets worse or lasts more than 3 days
- redness or swelling is present ■ new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away.

##### Directions

- take only as directed
- do not exceed 8 softgels per 24 hrs

adults & children 12 yrs and over	2 softgels with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

##### Other information

- store at controlled room temperature 20–25°C (68–77°F)
- protect from light, heat, and moisture

##### Inactive ingredients

FD&C Yellow No. 6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, sorbitol sorbitan solution, titanium dioxide

##### Questions or comments? Call 1-888-423-0139

DISTRIBUTED BY TOPCO ASSOCIATES LLC

ITASCA, IL 60143 ©TOPCO STVA0825

QUESTIONS? 1-888-423-0139

topcare@topco.com www.topcarebrand.com

PRODUCT OF INDIA

**QUALITY GUARANTEED**

This product is manufactured in the USA.



Scan here for more

Questions or comments? Call 1-888-423-0139

This TopCare® product is laboratory tested to guarantee its highest quality. Your total satisfaction is guaranteed.

Information or call 1-888-423-0139

## COMBINATION PACK TOPCARE HEALTH DAYTIME SEVERE AND NITETIME SEVERE COLD AND FLU RELIEF SOFTGELS - 24 CT

acetaminophen, dextromethorphan hydrobromide, guaifenesin, doxylamine succinate, phenylephrine hydrochloride kit

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:76162-678
--------------	----------------	--------------------	---------------

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76162-678-24	1 in 1 CARTON; Type 0: Not a Combination Product	12/11/2025	

### Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BLISTER PACK	12
Part 2	1 BLISTER PACK	12

### Part 1 of 2

#### DAYTIME SEVERE COLD AND FLU RELIEF SOFTGELS

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride capsule, liquid filled

### Product Information

Item Code (Source)	NDC:76162-636
Route of Administration	ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

## Inactive Ingredients

Ingredient Name	Strength
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C00X)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6092ICV9RU)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

## Product Characteristics

Color	orange	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	A13
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/11/2025	

## Part 2 of 2

### NITETIME SEVERE COLD AND FLU RELIEF SOFTGELS

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride capsule, liquid filled

## Product Information

Item Code (Source)	NDC:76162-637
Route of Administration	ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362091TL9D) (ACETAMINOPHEN - UNII:362091TL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN, UNSPECIFIED</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C00X)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SORBITAN</b> (UNII: 6092ICV9RU)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	18mm
<b>Flavor</b>		<b>Imprint Code</b>	A07
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/11/2025	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/11/2025	

Revised: 12/2025

Topco Associates LLC