

**BASIC CARE DAYTIME COLD AND FLU NIGHTTIME COLD AND FLU-
acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine
hcl**

Amazon.com Services LLC

Amazon Daytime Cold & Flu Nighttime Cold & Flu Drug Facts

Active ingredients (in each softgel) - Nighttime Cold & Flu

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

Active ingredients (in each softgel) - Daytime Cold & Flu

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

Purpose - Nighttime Cold & Flu

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Purpose - Daytime Cold & Flu

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Uses - Nighttime Cold & Flu

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- minor aches and pains
- sore throat
- headache
- runny nose and sneezing
- fever

Uses - Daytime Cold & Flu

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- minor aches and pains
- sore throat
- headache
- nasal congestion
- fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have - Nighttime Cold & Flu

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough as occurs with smoking, asthma, or emphysema

- trouble urinating due to an enlarged prostate gland

Ask a doctor before use if you have - Daytime Cold & Flu

- liver disease
- diabetes
- heart disease
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are - Nighttime Cold & Flu

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

Ask a doctor or pharmacist before use if you are - Daytime Cold & Flu

taking the blood thinning drug warfarin

When using this product - Nighttime Cold & Flu

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

When using the product - Daytime Cold & Flu

do not use more than directed

Stop use and ask a doctor if - Nighttime Cold & Flu

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Stop use and ask a doctor if - Daytime Cold & Flu

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions - Nighttime Cold & Flu

- take only as directed – see overdose warning
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	2 softgels with water every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Directions - Daytime Cold & Flu

- take only as directed – see overdose warning
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	2 softgels with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients - Nighttime Cold & Flu

D&C yellow no. 10, edible ink*, FD&C blue no. 1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution *may contain this ingredient

Inactive ingredients - Daytime Cold & Flu

edible ink*, FD&C red no. 40, FD&C yellow no. 6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution *may contain this ingredient

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Multi-Symptom Relief

Compare to Vicks[®] DayQuil[®] Cold & Flu active ingredients

Non-Drowsy

Daytime Cold & Flu

Acetaminophen

Dextromethorphan HBr

Phenylephrine HCl

Pain Reliever/Fever Reducer

Nasal Decongestant

Cough Suppressant

Alcohol Free

Antihistamine Free

actual size

32 SOFTGELS

Multi-Symptom Relief

Compare to Vicks[®] NyQuil[®] Cold & Flu active ingredients

Nighttime Cold & Flu

Acetaminophen

Dextromethorphan HBr

Doxylamine Succinate

Pain Reliever/Fever Reducer

Antihistamine

Cough Suppressant

Alcohol Free

actual size

16 SOFTGELS



Multi-Symptom Relief

Compare to Vicks® DayQuil® Cold & Flu active ingredients**

Multi-Symptom Relief

Compare to Vicks® NyQuil® Cold & Flu active ingredients**

Non-Drowsy

Daytime Cold & Flu

Acetaminophen
Dextromethorphan HBr
Phenylephrine HCl

Pain Reliever/
Fever Reducer
Nasal Decongestant
Cough Suppressant

Alcohol Free
Antihistamine Free

actual size



32 SOFTGELS

Nighttime Cold & Flu

Acetaminophen
Dextromethorphan HBr
Doxylamine Succinate

Pain Reliever/
Fever Reducer
Antihistamine
Cough Suppressant



Alcohol Free

actual size



16 SOFTGELS

VISIBLE PRINT AREA



Nighttime Cold & Flu

Daytime Cold & Flu

<p>Drug Facts</p> <p>Active ingredients (in each softgel) Acetaminophen 325 mg.....Pain reliever/fever reducer Dextromethorphan HBr 15 mg.....Cough suppressant Doxylamine succinate 6.25 mg.....Antihistamine</p> <p>Purpose Pain reliever/fever reducer Cough suppressant Antihistamine</p> <p>Uses temporarily relieves common cold/flu symptoms: cough due to minor throat and bronchial irritation minor aches and pains sore throat headache runny nose and sneezing fever</p> <p>Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin redness blisters rash If a skin reaction occurs, stop use and seek medical help right away.</p>		<p>Drug Facts</p> <p>Active ingredients (in each softgel) Acetaminophen 325 mg.....Pain reliever/fever reducer Dextromethorphan HBr 10 mg.....Cough suppressant Phenylephrine HCl 5 mg.....Nasal decongestant</p> <p>Purpose Pain reliever/fever reducer Cough suppressant Nasal decongestant</p> <p>Uses temporarily relieves common cold/flu symptoms: cough due to minor throat and bronchial irritation minor aches and pains sore throat headache nasal congestion fever</p> <p>Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin redness blisters rash If a skin reaction occurs, stop use and seek medical help right away.</p>	
---	--	--	--

<p>Drug Facts (continued)</p> <p>Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p>Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. if you have ever had an allergic reaction to this product or any of its ingredients</p> <p>Ask a doctor before use if you have liver disease glaucoma cough that occurs with too much phlegm (mucus) a breathing problem such as asthma or chronic bronchitis persistent or chronic cough as occurs with smoking, asthma, or emphysema trouble urinating due to an enlarged prostate gland</p> <p>Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers taking the blood thinning drug warfarin</p> <p>When using this product excitability may occur, especially in children marked drowsiness may occur avoid alcohol drinks be careful when driving a motor vehicle or operating machinery alcohol, sedatives, and tranquilizers may increase drowsiness</p> <p>Stop use and ask a doctor if pain or cough gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days redness or swelling is present new symptoms occur cough comes back or occurs with rash or headache that lasts These could be signs of a serious condition.</p>		<p>Drug Facts (continued)</p> <p>Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p>Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. if you have ever had an allergic reaction to this product or any of its ingredients</p> <p>Ask a doctor before use if you have liver disease diabetes heart disease high blood pressure thyroid disease trouble urinating due to an enlarged prostate gland cough that occurs with too much phlegm (mucus) persistent or chronic cough as occurs with smoking, asthma, or emphysema</p> <p>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin</p> <p>When using this product do not use more than directed</p> <p>Stop use and ask a doctor if you get nervous, dizzy or sleepless pain, nasal congestion or cough gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days redness or swelling is present new symptoms occur cough comes back or occurs with rash or headache that lasts These could be signs of a serious condition.</p>	
---	--	---	--

<p>Drug Facts (continued)</p> <p>If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions take only as directed - see overdose warning do not exceed 4 doses per 24 hrs</p> <table border="1"> <tr> <td>adults & children 12 yrs & over</td> <td>2 softgels with water every 6 hrs</td> </tr> <tr> <td>children 4 to under 12 yrs</td> <td>ask a doctor</td> </tr> <tr> <td>children under 4 yrs</td> <td>do not use</td> </tr> </table> <p>Other information store at 20-25°C (68-77°F)</p> <p>Inactive ingredients D&C yellow no. 10, edible ink, FD&C blue no. 1, gelatin, glycerin, polyethylene glycol, polydioxane, polyethylene glycol, purified water, sorbitol sorbitan solution *may contain this ingredient</p> <p>Questions or comments? 1-800-719-9260</p>		adults & children 12 yrs & over	2 softgels with water every 6 hrs	children 4 to under 12 yrs	ask a doctor	children under 4 yrs	do not use	<p>Drug Facts (continued)</p> <p>If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions take only as directed - see overdose warning do not exceed 4 doses per 24 hrs</p> <table border="1"> <tr> <td>adults & children 12 yrs & over</td> <td>2 softgels with water every 4 hrs</td> </tr> <tr> <td>children 4 to under 12 yrs</td> <td>ask a doctor</td> </tr> <tr> <td>children under 4 yrs</td> <td>do not use</td> </tr> </table> <p>Other information store at 20-25°C (68-77°F)</p> <p>Inactive ingredients edible ink, FD&C red no. 40, FD&C yellow no. 6, gelatin, glycerin, polyethylene glycol, polydioxane, propylene glycol, purified water, sorbitol sorbitan solution *may contain this ingredient</p> <p>Questions or comments? 1-800-719-9260</p>		adults & children 12 yrs & over	2 softgels with water every 4 hrs	children 4 to under 12 yrs	ask a doctor	children under 4 yrs	do not use
adults & children 12 yrs & over	2 softgels with water every 6 hrs														
children 4 to under 12 yrs	ask a doctor														
children under 4 yrs	do not use														
adults & children 12 yrs & over	2 softgels with water every 4 hrs														
children 4 to under 12 yrs	ask a doctor														
children under 4 yrs	do not use														

58072 BNC-1

CODE AREA

OPEN OTHER END

MADE IN CANADA

DISTRIBUTED BY:
 AMAZON.COM SERVICES LLC
 410 TERRY AVENUE N.
 SEATTLE, WA 98109

© 2020 AMAZON.COM INC. OR ITS AFFILIATES. ALL RIGHTS RESERVED.

DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN

**These products are not manufactured or distributed by Procter & Gamble (distributor of Wicac® DayQuil® Cold & Flu and Micsac® NyQuil® Cold & Flu)

Gluten Free
 www.StopMedicineAbuse.org

PARENTS:
 Learn about how medicines abuse

CONVENIENT RECYCLING TAB

BASIC CARE DAYTIME COLD AND FLU NIGHTTIME COLD AND FLU				
acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl kit				
Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72288-580	
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72288-580-72	1 in 1 CARTON; Type 0: Not a Combination Product	08/26/2021	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	8 BLISTER PACK	16
Part 2	16 BLISTER PACK	32

Part 1 of 2

BASIC CARE NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled

Product Information

Item Code (Source)	NDC:72288-056
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	

Product Characteristics

Color	GREEN (clear)	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	056
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72288-056-00	8 in 1 CARTON		
1		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012		

Part 2 of 2

BASIC CARE DAYTIME COLD AND FLU

acetaminophen, dextromethorphan hbr, phenylephrine hcl capsule, liquid filled

Product Information

Item Code (Source)	NDC:72288-994
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	

Product Characteristics			
Color	ORANGE	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	L994
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72288-994-00	16 in 1 CARTON		
1		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/26/2021	

Labeler - Amazon.com Services LLC (128990418)

Revised: 11/2024

Amazon.com Services LLC