

LIDOCAINE- maximum strength pain reliever cream
Shinpharma Inc

Dr. Numb Maximum Strength Pain Reliever

Active ingredient

Lidocaine 5%

Purpose

Anorectal (Hemorrhoidal)

Uses

For temporary relief of local discomfort, itching, soreness or burning in the perianal area associated with anorectal disorders

Warnings

For external use only

When using this product

- do not exceed the recommended daily dosage unless directed by a doctor
- do not put into the rectum by using fingers or any mechanical device or applicator

Stop use and ask a doctor if

- allergic reaction occurs
- rectal bleeding occurs
- redness, irritation, swelling, pain or other symptoms begin or increase
- condition worsens or does not improve within 7 days

Keep Out Of Reach Of Children

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Adults and Children 12 years and older: When practical, cleanse the affected area with mild soap and warm water and rinse thoroughly. Gently dry by patting or blotting with toilet tissue or a soft cloth before application of this product. Apply to the affected area up to 6 times a day. Children under 12 years: Consult a doctor.

Other Information

- always keep the tube tightly closed
- store at temperatures not exceeding 15°C - 30°C (59°F - 86°F)
- protect from freezing

Inactive Ingredients

Benzyl Alcohol, Carbopol, Lecithin, Propylene Glycol, Tocopheryl Acetate, Water

Questions or comments?

Contact our dedicated Customer Care at +1(877)786 2001 or email us at support@shinpharma.com

Principal Display Panel

Dr. Numb

Maximum Strength Pain Reliever

Prompt, Soothing Relief from Painful Burning, Itching and Discomfort

NDC 69903-005-5

Dr. Numb[®]

Lidocaine Cream

For External Use Only

Maximum Strength Pain Reliever

5g

| | |
|--|---|
| <p>Drug Facts</p> <p>Active ingredient Lidocaine 5% Anorectal (Hemorrhoidal)</p> <p>Purpose</p> <p>Uses For temporary relief of local discomfort, itching, pain, soreness or burning in the perianal area associated with anorectal disorders</p> <p>Warnings For external use only</p> <p>When using this product</p> <ul style="list-style-type: none"> ■ do not exceed the recommended daily dosage unless directed by a doctor ■ do not put into the rectum by using fingers or any mechanical device or applicator <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ allergic reaction occurs ■ rectal bleeding occurs ■ redness, irritation, swelling, pain or other symptoms begin or increase ■ condition worsens or does not improve within 7 days <p>KEEP OUT OF REACH OF CHILDREN. If swallowed, get medical help or contact a Poison Control Center right away. ▶</p> | <p>Drug Facts (continued)</p> <p>Directions</p> <ul style="list-style-type: none"> ■ Adults and Children 12 years and older: When practical, cleanse the affected area with mild soap and warm water and rinse thoroughly. Gently dry by patting or blotting with toilet tissue or a soft cloth before application of this product. Apply to the affected area up to 6 times a day. <p>Children under 12 years: Consult a doctor</p> <p>Other Information</p> <ul style="list-style-type: none"> ■ always keep the tube tightly closed ■ store at temperatures not exceeding 15°C – 30°C (59°F - 86°F) ■ protect from freezing <p>Inactive Ingredients Benzyl Alcohol, Carbopol, Lecithin, Propylene Glycol, Tocopheryl Acetate, Water</p> <p>Questions or comments? Contact our dedicated Customer Care at +1(877) 786 2001 or email us at support@shinpharma.com</p> <p>Imported/Distributed by: Shinpharma Inc, 1124 Fir Avenue, Blaine, Washington, USA 98230</p> |
|--|---|



| | | | |
|--|--|---------------------------|-----------------|
| LIDOCAINE | | | |
| maximum strength pain reliever cream | | | |
| Product Information | | | |
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:69903-005 |
| Route of Administration | TOPICAL | | |
| Active Ingredient/Active Moiety | | | |
| | Ingredient Name | Basis of Strength | Strength |
| | LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987) | LIDOCAINE | 5 g in 100 g |
| Inactive Ingredients | | | |
| | Ingredient Name | | Strength |
| | WATER (UNII: 059QF0KO0R) | | |

CARBOMER HOMOPOLYMER, UNSPECIFIED TYPE (UNII: 0A5MM307FC)

LECITHIN, SUNFLOWER (UNII: 834K0WOS5G)

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)

ALPHA-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0)

BENZYL ALCOHOL (UNII: LKG8494WBH)

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:69903-005-05 | 1 in 1 BOX | 10/19/2021 | |
| 1 | | 50 g in 1 TUBE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M015 | 10/19/2021 | |

Labeler - Shinpharma Inc (248552403)

Revised: 4/2024

Shinpharma Inc