

**BAC-STOP NPE FREE- iodine liquid
IBA**

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

BAC-STOP

ACTIVE INGREDIENTS :

Iodine.....0.50%

(Provides A Full .5% Available Iodine as used

Inert Ingredients

INERT INGREDIENTS:

Skin Conditioners.....2.00%

Other Inerts.....95.50%

(Includes Iodine Stabilizers, Buffering Agents and Surfactants)

TOTAL:.....100.00%

pH of 5.5 at time of manufacture

This product contains highly effective levels of free iodine for fast kill of mastitis causing organisms.

GENERAL RECOMMENDATIONS

1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
2. When solution in the teat dip cup becomes dirty and/or noticeably loses color, discard remaining contents and refill cup with fresh teat dip.
3. Do not use this product for cleaning or sanitizing equipment.
4. Wash teat dip cups after each milking.

CAUTION: KEEP OUT OF REACH OF CHILDREN.

AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY.

FIRST AID PROCEDURES

EYE CONTACT: IMMEDIATELY FLUSH EYE WITH LARGE AMOUNTS OF FRESH, TEPID WATER FOR 15 MINUTES. HOLD EYELID OPEN TO ENSURE COMPLETE IRRIGATION OF EYE AND LID TISSUES. TILT HEAD TO THE SIDE AND IRRIGATE THE EYE FROM THE

BRIDGE OF THE NOSE TO THE OUTSIDE OF THE FACE.

SKIN CONTACT: FOR READY-TO-USE PRODUCTS, WASH ANY SENSITIVE AREAS.

INGESTIONS: DO NOT INDUCE VOMITING. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. DRINK LARGE AMOUNTS OF WATER. IF VOMITING OCCURS KEEP AIRWAYS OPEN. KEEP HEAD LOWER THAN THE HIPS TO PREVENT ASPIRATION INTO THE LUNGS.

ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION

FOR MEDICAL EMERGENCIES CALL 1-800-424-9300

RECOMMENDED PROCEDURES FOR TEAT SANITATION

A. PRE-MILKING (PRE-DIPPING)

1. Remove visible soils from the teats.
2. Observe foremilk by stripping two or three streams of milk into a strip cup.
3. Dip, foam or spray the cow's teats full length with BAC-STOP NPE FREE Teat Dip.
4. Wait approximately 30 seconds.
5. Remove all teat dip by thoroughly drying the teats, teat orifices and udder (if foaming or spraying) with a clean, single service paper towel(s)

to avoid contamination of milk.

6. Attach milking unit.

B. POST-MILKING (POST-DIPPING)

1. Immediately after milking, dip or spray all teats full length with BAC-STOP NPE FREE Teat Dip. Ensure good coverage of all sides of the teats and teat orifices.

UNIVERSITY TESTED AND PASSED USING

NATIONAL MASTITIS COUNCIL PROTOCOLS

PROVEN EFFECTIVE IN REDUCING

NEW INTRAMAMMARY INFECTIONS.

WARNING

DO NOT DILUTE THIS PRODUCT.

DO NOT RETURN TEAT DIP LEFT IN THE TEAT DIPPER TO STORAGE CONTAINER.


ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER CONDITIONS WHICH MAY CAUSE DAMAGE TO THE TEATS.

PROTECT THIS PRODUCT FROM FREEZING.

Not For Human Use

Bac-Stop NPE Free

NDC 29983-1014-4



BAC-STOP

SANITIZING TEAT DIP

NPE
FREE

NPE
FREE

A Highly Effective Pre-Milking and Post-Milking Iodine Teat Dip
Contains Triple Emollients For Excellent Skin Conditioning

ACTIVE INGREDIENT:
Iodine 0.50%
(Provides A Full .5% Available Iodine As Used)

INERT INGREDIENTS:
Skin Conditioners 2.00%
Other Inerts 97.50%
(Includes Iodine Stabilizers, Buffering Agents and Surfactants)
TOTAL: 100.00%

pH of 5.5 at time of manufacture

This product contains highly effective levels of free iodine for fast kill of mastitis causing organisms.

GENERAL RECOMMENDATIONS

1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
2. When solution in the teat dip cup becomes dirty and/or noticeably loses color, discard remaining contents and refill cup with fresh teat dip.
3. Do not use this product for cleaning or sanitizing equipment.
4. Wash teat dip cups after each milking.

**CAUTION: KEEP OUT OF REACH OF CHILDREN.
AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY.**


FIRST AID PROCEDURES

EYE CONTACT: IMMEDIATELY FLUSH EYE WITH LARGE AMOUNTS OF FRESH, TEPID WATER FOR 15 MINUTES. HOLD EYELID OPEN TO ENSURE COMPLETE IRRIGATION OF EYE AND LID TISSUES. TILT HEAD TO THE SIDE AND IRRIGATE THE EYE FROM THE BRIDGE OF THE NOSE TO THE OUTSIDE OF THE FACE.

SKIN CONTACT: FOR READY-TO-USE PRODUCTS, WASH ANY SENSITIVE AREAS.

INGESTION: DO NOT INDUCE VOMITING. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. DRINK LARGE AMOUNTS OF WATER. IF VOMITING OCCURS, KEEP AIRWAYS OPEN. KEEP HEAD LOWER THAN THE HIPS TO PREVENT ASPIRATION INTO THE LUNGS.

ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION
FOR MEDICAL EMERGENCIES CALL 1-800-424-9300



PATENT
PENDING

RECOMMENDED PROCEDURES FOR TEAT SANITATION

A. PRE-MILKING (PRE-DIPPING)

1. Remove visible soils from the teats.
2. Observe foremilk by stripping two or three streams of milk into a strip cup.
3. Dip, foam or spray the cow's teats full length with **BAC-STOP NPE FREE Teat Dip**.
4. Wait approximately 30 seconds.
5. Remove all teat dip by thoroughly drying the teats, teat orifices and udder (if foaming or spraying) with a clean, single service paper towel(s) to avoid contamination of milk.
6. Attach milking unit.

B. POST-MILKING (POST-DIPPING)

1. Immediately after milking, dip or spray all teats full length with **BAC-STOP NPE FREE Teat Dip**. Ensure good coverage of all sides of the teats and teat orifices.

WARNING

DO NOT DILUTE THIS PRODUCT.
DO NOT RETURN TEAT DIP LEFT IN THE TEAT DIPPER TO STORAGE CONTAINER.
ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER CONDITIONS WHICH MAY CAUSE DAMAGE TO THE TEATS.
PROTECT THIS PRODUCT FROM FREEZING.

Not For Human Use

A PART OF THE COMPLETE IBA SANITATION PROGRAM

EXP. DATE: LOT NO.: NET CONTENTS: 5 GALLONS (18.925 LITERS) IBA STOCK # 260633 121W

MANUFACTURED EXCLUSIVELY FOR IBA INC., SANITATION DIVISION, 103 GILMORE DRIVE, SUTTON, MA 01590

BAC-STOP NPE FREE

iodine liquid

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:29983-1014
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	0.5 kg in 100 kg

Inactive Ingredients

Ingredient Name	Strength

SODIUM HYDROXIDE (UNII: 55X04QC32I)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:29983-1014-4	19.12 kg in 1 PAIL		
2	NDC:29983-1014-6	57.36 kg in 1 DRUM		
3	NDC:29983-1014-8	210.31 kg in 1 DRUM		
4	NDC:29983-1014-9	1051.54 kg in 1 CELLO PACK		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		11/15/2018	

Labeler - IBA (019494160)

Registrant - IBA (019494160)

Establishment

Name	Address	ID/FEI	Business Operations
Morgan Gallacher Inc. DBA Custom Chemical Formulators Inc,		028311595	manufacture, api manufacture, pack

Revised: 12/2023

IBA