

**VANACOF- chlophedianol hydrochloride, dexchlorpheniramine maleate, and pseudoephedrine hydrochloride liquid**  
**GM Pharmaceuticals, INC**

-----  
**VanaCof**®

***Drug Facts***

***Active ingredients (in each 5 mL teaspoonful)***

Chlophedianol Hydrochloride 12.5 mg

Dexchlorpheniramine Maleate 1 mg

Pseudoephedrine Hydrochloride 30 mg

***Purpose***

Cough Suppressant

Antihistamine

Nasal Decongestant

***Uses***

- temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:
  - runny nose
  - sneezing
  - itching of the nose or throat
  - itchy, watery eyes
  - nasal congestion
  - reduces swelling of nasal passages
- cough due to minor throat and bronchial irritation

***Warnings***

**Do not exceed recommended dosage.**

**Do not use this product**

If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product

**Ask a doctor before use if you have**

- a cough that lasts or is chronic such as occurs with smoking, asthma or emphysema
- a cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- glaucoma
- heart disease
- high blood pressure
- thyroid disease
- diabetes

**Ask a doctor before use if you are**

taking sedatives or tranquilizers.

**When using this product**

- excitability may occur, especially in children
- may cause marked drowsiness
- alcohol, sedatives, and tranquilizers may increase the drowsiness effect
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by a fever, rash or persistent headache. A persistent cough may be a sign of serious condition.
- new symptoms occur

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

**Do not exceed recommended dosage.**

adults and children 12 years of age and over:	2 teaspoonfuls every 6 hours, not to exceed 8 teaspoonfuls in 24 hours.
children 6 to under 12 years of age:	1 teaspoonful every 6 hours, not to exceed 4 teaspoonfuls in 24 hours.
children under 6 years of age:	consult a doctor.

***Other information***

- this package is child resistant

- store at room temperature of 68°-86°F (20°-30°C) with excursions of 59°-86°F (15°-30°C)

***Inactive ingredients***

Citric acid anhydrous, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sorbitol solution, sucralose

***Questions or Comments?***

Call 1-888-535-0305 9a.m. – 5p.m. CST

**PRINCIPAL DISPLAY PANEL**

NDC 58809-999-01

VanaCof<sup>®</sup>

Each 5 mL (1 TEASPOONFUL) CONTAINS:

Chlophedianol Hydrochloride ..... 12.5 mg

Dexchlorpheniramine Maleate ..... 1 mg

Pseudoephedrine Hydrochloride ..... 30mg

NDC 58809-999-01

**VANACOF<sup>®</sup>**

EACH 5 mL (1 TEASPOONFUL) CONTAINS:

Chlrophedianol Hydrochloride .... 12.5 mg  
Dexchlorpheniramine Maleate ..... 1 mg  
Pseudoephedrine Hydrochloride .... 30 mg

**Cough Suppressant  
Antihistamine  
Nasal Decongestant**

**Sugar Free, Alcohol Free,  
Dye Free**

**Tamper evident by foil seal under cap.  
Do not use if foil seal is broken or missing.**

**THIS BOTTLE IS NOT TO BE DISPENSED TO CONSUMER.  
Dispense in a tight, light-resistant container with a  
child-resistant cap.**



US Patent # 9,463,191  
US Patent # 9,050,289

**GM Pharmaceuticals, Inc.**

**16 fl. oz. (473 mL)**

## Drug Facts

### Active ingredients Purpose in each 5 mL (1 teaspoonful)

Chlrophedianol Hydrochloride 12.5 mg..Cough Suppressant  
Dexchlorpheniramine Maleate 1 mg.....Antihistamine  
Pseudoephedrine Hydrochloride 30 mg..Nasal Decongestant

### Uses

- temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:
  - runny nose
  - sneezing
  - itching of the nose or throat
  - itchy, watery eyes
  - nasal congestion
  - reduces swelling of nasal passages
- cough due to minor throat and bronchial irritation

### Warnings

**Do not exceed recommended dosage.**

**Do not use this product** if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product

### Ask a doctor before use if you have

- a cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema
- a cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- glaucoma
- heart disease
- high blood pressure
- thyroid disease
- diabetes

**Ask a doctor before use if you are** taking sedatives or tranquilizers

Distributed by: GM Pharmaceuticals, Inc.  
Arlington, TX 76018  
201046-01 Rev.0725

**PEEL**

**Drug Facts** (continued)**When using this product**

- excitability may occur, especially in children
- may cause marked drowsiness
- alcohol, sedatives, and tranquilizers may increase the drowsiness effect
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by a fever, rash or persistent headache. A persistent cough may be a sign of a serious condition.
- new symptoms occur

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- mL= milliliter

Age	Dose
adults and children 12 years of age and over:	2 teaspoonfuls (10 mL) every 6 hours, not to exceed 8 teaspoonfuls (40 mL) in 24 hours.
children 6 to under 12 years of age:	1 teaspoonful (5 mL) every 6 hours, not to exceed 4 teaspoonfuls (20 mL) in 24 hours.
children under 6 years of age:	consult a doctor.

**Other information** ■ this packaging is child-resistant.  
 ■ store at room temperature of 68°-86°F (20°-30°C) with excursions of 59°-86°F (15°-30°C)

**Inactive ingredients** citric acid anhydrous, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sorbitol solution, sucralose

**Questions?** Call 1-888-535-0305 9a.m. - 5p.m. CST

**VANACOF**

chlorphedianol hydrochloride, dexchlorpheniramine maleate, and pseudoephedrine hydrochloride liquid

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:58809-999
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>CHLOPHEDIANOL HYDROCHLORIDE</b> (UNII: 69QQ58998Y) (CHLOPHEDIANOL - UNII:42C50P12AP)	CHLOPHEDIANOL HYDROCHLORIDE	12.5 mg in 5 mL
<b>DEXCHLORPHENIRAMINE MALEATE</b> (UNII: B10YD955QW) (DEXCHLORPHENIRAMINE - UNII:3Q9Q0B929N)	DEXCHLORPHENIRAMINE MALEATE	1 mg in 5 mL
<b>PSEUDOEPHEDRINE HYDROCHLORIDE</b> (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	30 mg in 5 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

## Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	TUTTI FRUTTI	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58809-999-02	12 in 1 TRAY	04/22/2008	
1		15 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:58809-999-01	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	04/22/2008	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/22/2008	

**Labeler** - GM Pharmaceuticals, INC (793000860)

Revised: 10/2025

GM Pharmaceuticals, INC