

MECLIZINE- meclizine hcl 25mg tablet, chewable
PD-Rx Pharmaceuticals, Inc.

Drug Facts

Active ingredient (in each tablet)

Meclizine HCl 25mg

Purpose

Antiemetic

Uses

prevents and treats nausea, vomiting or dizziness associated with motion sickness

Warnings

Do not use in children under 12 years of age unless directed by a doctor.

Do not take this product unless directed by a doctor if you have

- glaucoma
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland

Do not take this product if you are taking sedatives or tranquilizers, without first consulting your doctor.

When using this product

- do not exceed recommended dosage
- may cause drowsiness
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222)

Directions

- doage should be taken one hour before travel starts

Adults and children 12 years and over	take 1 or 2 tablets once daily or as directed by doctor
--	--

Other information

- **Tamper Evident:** do not use if seal is broken or missing from bottle.
- store at 20°-25°C (68°-77°F)

Inactive ingredients

Crosscarmellose sodium, dextrose, FD&C Red#40, raspberry flavor, magnesium stearate, microcrystalline cellulose, silicon dioxide, sodium saccharine, stearic acid

Questions or comments?

(866) 562-2756 (Mon - Fri, 8AM - 4PM EST)

MECLIZINE

CHEWABLE

Raspberry Flavor

<table border="1"> <tr> <th colspan="2">Drug Facts</th> </tr> <tr> <th>Active ingredient (in each tablet)</th> <th>Purpose</th> </tr> <tr> <td>Meclizine HCl 25mg</td> <td>Antiemetic</td> </tr> <tr> <th colspan="2">Uses</th> </tr> <tr> <td colspan="2">prevents and treats nausea, vomiting or dizziness associated with motion sickness</td> </tr> <tr> <th colspan="2">Warnings</th> </tr> <tr> <td colspan="2">Do not use for children under 12 years of age unless directed by a doctor. Do not take unless directed by a doctor if you have - glaucoma - trouble urinating due to an enlarged prostate gland - a breathing problem such as emphysema or chronic bronchitis. Do not take if you are taking sedatives or tranquilizers, without first consulting your doctor.</td> </tr> <tr> <td colspan="2">When using this product - do not exceed recommended dosage - drowsiness may occur - alcohol, sedatives, and tranquilizers may increase drowsiness - avoid alcoholic drinks - be careful when driving a motor vehicle or operating machinery</td> </tr> <tr> <td colspan="2">If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (800) 222-1222</td> </tr> <tr> <th colspan="2">Directions</th> </tr> <tr> <td colspan="2">- dosage should be taken 1 hour before travel starts</td> </tr> <tr> <td>Adults and children 12 years and over</td> <td>Take 1 or 2 Tablets once daily or as directed by doctor</td> </tr> <tr> <th colspan="2">Other information</th> </tr> <tr> <td colspan="2">- store at room temperature 20°- 25°C (68°- 77°F)</td> </tr> <tr> <th colspan="2">Inactive ingredients</th> </tr> <tr> <td colspan="2">Crosscarmellose sodium, dextrose, FD&C Red #40, flavor, magnesium stearate, microcrystalline cellulose, silicon dioxide, sodium saccharine, stearic acid</td> </tr> <tr> <td colspan="2">Questions? Adverse drug event call (866) 562-2756 (M-F, 8AM-4PM EST).</td> </tr> </table>	Drug Facts		Active ingredient (in each tablet)	Purpose	Meclizine HCl 25mg	Antiemetic	Uses		prevents and treats nausea, vomiting or dizziness associated with motion sickness		Warnings		Do not use for children under 12 years of age unless directed by a doctor. Do not take unless directed by a doctor if you have - glaucoma - trouble urinating due to an enlarged prostate gland - a breathing problem such as emphysema or chronic bronchitis. Do not take if you are taking sedatives or tranquilizers, without first consulting your doctor.		When using this product - do not exceed recommended dosage - drowsiness may occur - alcohol, sedatives, and tranquilizers may increase drowsiness - avoid alcoholic drinks - be careful when driving a motor vehicle or operating machinery		If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (800) 222-1222		Directions		- dosage should be taken 1 hour before travel starts		Adults and children 12 years and over	Take 1 or 2 Tablets once daily or as directed by doctor	Other information		- store at room temperature 20°- 25°C (68°- 77°F)		Inactive ingredients		Crosscarmellose sodium, dextrose, FD&C Red #40, flavor, magnesium stearate, microcrystalline cellulose, silicon dioxide, sodium saccharine, stearic acid		Questions? Adverse drug event call (866) 562-2756 (M-F, 8AM-4PM EST).		<p>NDC 43063-804-12</p>  <p>Meclizine Chewable 25 mg</p> <p>Raspberry Flavor</p> <ul style="list-style-type: none"> • Motion sickness • Nausea • Vomiting <p>12 Tablets</p> <p>TAMPER EVIDENT: DO NOT USE IF SEAL IS BROKEN OR MISSING FROM BOTTLE.</p>	<p>Marketed and Packaged by: PD-Rx Pharmaceuticals, Inc Oklahoma City, OK 73127 1-405-942-3040 v.8.19.0</p>  <p>3 43063 80412 4</p>  <p>GTIN: 00343063804124 SNO: K19E010002 EXP: 04/2021 LOT: K19E01</p>
Drug Facts																																				
Active ingredient (in each tablet)	Purpose																																			
Meclizine HCl 25mg	Antiemetic																																			
Uses																																				
prevents and treats nausea, vomiting or dizziness associated with motion sickness																																				
Warnings																																				
Do not use for children under 12 years of age unless directed by a doctor. Do not take unless directed by a doctor if you have - glaucoma - trouble urinating due to an enlarged prostate gland - a breathing problem such as emphysema or chronic bronchitis. Do not take if you are taking sedatives or tranquilizers, without first consulting your doctor.																																				
When using this product - do not exceed recommended dosage - drowsiness may occur - alcohol, sedatives, and tranquilizers may increase drowsiness - avoid alcoholic drinks - be careful when driving a motor vehicle or operating machinery																																				
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (800) 222-1222																																				
Directions																																				
- dosage should be taken 1 hour before travel starts																																				
Adults and children 12 years and over	Take 1 or 2 Tablets once daily or as directed by doctor																																			
Other information																																				
- store at room temperature 20°- 25°C (68°- 77°F)																																				
Inactive ingredients																																				
Crosscarmellose sodium, dextrose, FD&C Red #40, flavor, magnesium stearate, microcrystalline cellulose, silicon dioxide, sodium saccharine, stearic acid																																				
Questions? Adverse drug event call (866) 562-2756 (M-F, 8AM-4PM EST).																																				

MECLIZINE

meclizine hcl 25mg tablet, chewable

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:43063-804(NDC:16103-387)
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
MECLIZINE HYDROCHLORIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570)	MECLIZINE HYDROCHLORIDE	25 mg

Inactive Ingredients	
Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
DEXTROSE (UNII: IY9XDZ35W2)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
RASPBERRY (UNII: 4N14V5R27W)	

Product Characteristics			
Color	pink (LIGHT PINK COLOR)	Score	2 pieces
Shape	ROUND (ROUND TABLET)	Size	8mm
Flavor		Imprint Code	PH051
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:43063-804-06	6 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/23/2018	
2	NDC:43063-804-12	12 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/01/2018	
3	NDC:43063-804-20	20 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/01/2018	
4	NDC:43063-804-30	30 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/01/2018	
5	NDC:43063-804-01	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/18/2018	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M009	02/01/2018	

Labeler - PD-Rx Pharmaceuticals, Inc. (156893695)

Registrant - PD-Rx Pharmaceuticals, Inc. (156893695)

Establishment

Name	Address	ID/FEI	Business Operations
PD-Rx Pharmaceuticals, Inc.		156893695	repack(43063-804)

Revised: 3/2025

PD-Rx Pharmaceuticals, Inc.