


CLEMASTINE FUMARATE- clemastine fumarate tablet Genus Lifesciences

CLEMASTINE FUMARATE TABLETS, USP

Rx only

DESCRIPTION

Clemastine belongs to the benzhydryl ether group of antihistaminic compounds. The chemical name is (+)-(2*R*)-2-[2-[[*R*]-*p*-Chloro- α -methyl- α -phenylbenzyl]-oxy]ethyl]-1-methylpyrrolidine fumarate (1:1).



$C_{21}H_{26}ClNO \cdot C_4H_4O_4$

M.W. 459.97

Each tablet for oral administration contains 2.68 mg of clemastine fumarate, USP.

Inactive ingredients: colloidal silicon dioxide, corn starch, lactose monohydrate, povidone, pregelatinized starch and stearic acid.

CLINICAL PHARMACOLOGY

Clemastine fumarate is an antihistamine with anticholinergic (drying) and sedative side effects. Antihistamines appear to compete with histamine for cell receptor sites on effector cells. The inherently long duration of antihistaminic effects of clemastine fumarate has been demonstrated in wheal and flare studies. In normal human subjects who received histamine injections over a 24-hour period, the antihistaminic activity of clemastine reached a peak at 5 to 7 hours, persisted for 10 to 12 hours and, in some cases, for as long as 24 hours. Pharmacokinetic studies in man utilizing ³H and ¹⁴C labeled compound demonstrates that: clemastine is rapidly and nearly completely absorbed from the gastrointestinal tract, peak plasma concentrations are attained in 2 to 4 hours, and urinary excretion is the major mode of elimination.

INDICATIONS AND USAGE

Clemastine Fumarate Tablets USP, 2.68 mg are indicated for the relief of symptoms associated with allergic rhinitis such as sneezing, rhinorrhea, pruritus, and lacrimation.

Clemastine Fumarate Tablets USP, 2.68 mg are also indicated for the relief of mild, uncomplicated allergic skin manifestations of urticaria and angioedema. It should be noted that clemastine fumarate is indicated for the dermatologic indications at the 2.68 mg dosage level only.

CONTRAINDICATIONS

Usage in Nursing Mothers

Because of the higher risk of antihistamines for infants generally and for newborns and prematures in particular, antihistamine therapy is contraindicated in nursing mothers.

Usage in Lower Respiratory Disease

Antihistamines *should* not be used to treat lower respiratory tract symptoms including asthma. Antihistamines are also contraindicated in the following conditions:

Hypersensitivity to clemastine fumarate or other antihistamines of similar chemical structure.

Monoamine oxidase inhibitor therapy (see Drug Interactions section).

WARNINGS

Antihistamines should be used with considerable caution in patients with: narrow angle glaucoma, stenosing peptic ulcer, pyloroduodenal obstruction, symptomatic prostatic hypertrophy, and bladder neck obstruction.

Usage in Children

Safety and efficacy of clemastine fumarate have not been established in children under the age of 12 years.

Usage in Pregnancy

Experience with this drug in pregnant women is inadequate to determine whether there is exists a potential for harm to the developing fetus.

Usage with CNS Depressants

Clemastine has additive effects with alcohol and other CNS depressants (hypnotics, sedatives, tranquilizers, etc.).

Usage in Activities Requiring Mental Alertness

Patients should be warned about engaging in activities requiring mental alertness such as driving a car or operating appliances, machinery, etc.

Usage in the Elderly (approximately 60 years or older)

Antihistamines are more likely to cause dizziness, sedation, and hypotension in elderly patients.

PRECAUTIONS

Clemastine fumarate should be used with caution in patients with: history of bronchial asthma, increased intraocular pressure, hyperthyroidism, cardiovascular disease, and hypotension.

Drug Interactions

MAO inhibitors prolong and intensify the anticholinergic (drying) effects of antihistamines.

ADVERSE REACTIONS

Transient drowsiness, the most common adverse reaction associated with clemastine fumarate, occurs relatively frequently and may require discontinuation of therapy in some instances.

Antihistaminic Compounds

It should be noted that the following reactions have occurred with one or more antihistamines and, therefore, should be kept in mind when prescribing drugs belonging to this class, including clemastine. The most frequent adverse reactions are italicized.

1. **General:** *Urticaria, drug rash, anaphylactic shock, photosensitivity, excessive perspiration, chills, dryness of the mouth, nose, and throat.*
2. **Cardiovascular System:** *Hypotension, headache, palpitations, tachycardia, extrasystoles.*
3. **Hematologic System:** *Hemolytic anemia, thrombocytopenia, agranulocytosis.*
4. **Nervous System:** *Sedation, sleepiness, dizziness, disturbed coordination, fatigue, confusion, restlessness, excitation, nervousness, tremor, irritability, insomnia, euphoria, paresthesias, blurred vision, diplopia, vertigo, tinnitus, acute labyrinthitis, hysteria, neuritis, convulsions.*
5. **GI System:** *Epigastric distress, anorexia, nausea, vomiting, diarrhea, constipation.*
6. **GU System:** *Urinary frequency, difficult urination, urinary retention, early menses.*
7. **Respiratory System:** *Thickening of bronchial secretions, tightness of chest and wheezing, nasal stuffiness.*

OVERDOSAGE

Antihistamine overdosage reactions may vary from central nervous system depression to stimulation.

Stimulation is particularly likely in children. Atropine-like signs and symptoms: dry mouth; fixed, dilated pupils; flushing; and gastrointestinal symptoms may also occur.

If vomiting has not occurred spontaneously the conscious patient should be induced to vomit. This is best done by having him drink a glass of water or milk after which he should be made to gag. Precautions against aspiration must be taken, especially in infants and children. If vomiting is unsuccessful gastric lavage is indicated within 3 hours after ingestion and even later if large amounts of milk or cream were given beforehand. Isotonic and 1/2 isotonic saline is the lavage solution of choice.

Saline cathartics, such as milk of magnesia, by osmosis draw water into the bowel and therefore, are valuable for their action in rapid dilution of bowel content.

Stimulants should *not* be used.

Vasopressors may be used to treat hypotension.

DOSAGE AND ADMINISTRATION

DOSAGE SHOULD BE INDIVIDUALIZED ACCORDING TO THE NEEDS AND RESPONSE OF THE PATIENT.

Clemastine Fumarate Tablets 2.68 mg

The maximum recommended dosage is one tablet three times daily. Many patients respond favorably to a single dose which may be repeated as required, but not to exceed three tablets daily.

HOW SUPPLIED

Clemastine Fumarate Tablets USP, 2.68 mg are white, round, scored tablets debossed with "NH" and "268". Tablets are packaged in bottles of 30 (NDC 64950-268-03) and 100 (NDC 64950-268-10)

Store at 20° to 25°C (68° to 77°F)

[See USP Controlled Room Temperature].

Dispense in a tight, light-resistant container as defined in the USP, with a child-resistant closure (as required).

Manufactured For:
Genus Lifesciences Inc.
Allentown, PA 18102

Rev. 08/2024

PRINCIPAL DISPLAY PANEL - 2.68 mg Tablet Bottle Label


NDC 64950-268-10

Clemastine
Fumarate
Tablets, USP

2.68 mg

Rx only
100 Tablets

Genus
Lifesciences Inc.



N
3 64950 26810 2

20689326 Rev. 08/2024

Manufactured for:
 Genus Lifesciences Inc
 Allentown, PA 18102

**KEEP THIS AND ALL
 MEDICATIONS OUT OF THE
 REACH OF CHILDREN.**

Dispense in a tight,
 light-resistant container as
 defined in the USP, with a
 child-resistant closure
 (as required).

Temperature:
 Store at 20°C to 25°C (68°F to 77°F)
 [See USP Controlled Room
 Temperature].

Usual Adult Dosage: One tablet
 one to three times daily. See
 package insert for full prescribing
 information.

Each tablet contains 2.68 mg of
 clemastine fumarate, USP.

NDC 64950-268-10

Clemastine Fumarate Tablets, USP

2.68 mg

Rx only 100 Tablets



CLEMASTINE FUMARATE			
clemastine fumarate tablet			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:64950-268
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
Clemastine Fumarate (UNII: 19259EGQ3D) (CLEMASTINE - UNII:95QN29S1ID)	Clemastine Fumarate	2.68 mg	
Inactive Ingredients			
Ingredient Name	Strength		
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
STARCH, CORN (UNII: O8232NY3SJ)			
LACTOSE, UNSPECIFIED FORM (UNII: J2B2A4N98G)			
POVIDONE K30 (UNII: U725QWY32X)			
STEARIC ACID (UNII: 4ELV7Z65AP)			
Product Characteristics			
Color	WHITE	Score	2 pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	NH;268
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:64950-268-03	30 in 1 BOTTLE; Type 0: Not a Combination Product	11/20/2024	
2	NDC:64950-268-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	11/20/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA073283	11/20/2024	

Labeler - Genus Lifesciences (113290444)

Revised: 2/2025

Genus Lifesciences