

**TREDA ULTRA- bismuth subsalicylate tablet**  
**Laboratorios Sanfer, S.A. de C.V.**

-----  
**Treda Ultra 44-749**

***Active ingredient (in each caplet)***

Bismuth subsalicylate 525 mg

***Purpose***

Upset stomach reliever/antidiarrheal

***Uses***

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
  - indigestion
  - fullness
  - gas
  - nausea
  - heartburn
  - belching

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use**

if you have

- an ulcer
- a bleeding problem
- bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool

## **Ask a doctor or pharmacist before use if you are**

taking any drug for

- gout
- diabetes
- arthritis
- anticoagulation (thinning the blood)

## **When using this product**

a temporary, but harmless, darkening of the stool and/or tongue may occur

## **Stop use and ask a doctor if**

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

## **If pregnant or breast-feeding,**

ask a health professional before use.

## **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

## ***Directions***

- **do not take more than directed**
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- swallow with water; do not chew
- adults and children 12 years and over: 1 caplet every 1/2 to 1 hour as needed. Do not exceed 8 caplets in 24 hours.
- do not use for more than 2 days unless directed by a doctor
- use until diarrhea stops, but not more than 2 days
- children under 12 years: ask a doctor

## ***Other information***

- **each caplet contains:** calcium 45 mg, salicylate 206 mg, sodium 3 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- avoid excessive heat
- see end flap for expiration date and lot number

## ***Inactive ingredients***

calcium carbonate, corn starch, D&C red #27 aluminum lake, D&C red #30 aluminum lake, magnesium stearate, mannitol, microcrystalline cellulose, povidone, silicon dioxide, sodium starch glycolate, stearic acid

**Questions or comments?**

**Call 1-(833) 604-1084** (English and Spanish) 8:00 AM-5:00 PM CST, Monday-Friday

**Principal display panel**

**Treda®**

**Ultra**

**Bismuth Subsalicylate 525 mg**

**UPSET STOMACH RELIEVER**

**ANTIDIARRHEAL**

**RELIEVES:**

**Diarrhea**

**Nausea**

**Heartburn**

**Upset Stomach**

**Indigestion**

*Actual Size*

**24**

**caplets**

**sanfer®**

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

50844

ORG032174908

Manufactured for: Laboratorios Sanfer S.A. de C.V.

Bld. Adolfo López Mateos No. 314

Col. Tlacopac, CDMX, 01049. México

This product is produced in compliance with all United States regulatory and quality requirements for pharmaceutical product.

**Treda<sup>®</sup> ULTRA**  
Bismuth Subsalicylate 525 mg  
Subsalicilato de Bismuto 525 mg

**UPSET STOMACH RELIEVER  
ALIVIA EL MALESTAR ESTOMACAL  
ANTI-DIARRHEAL  
ANTI-DIARRHEICO**

**24 caplets / comprimidos**

**Treda<sup>®</sup> ULTRA**  
Bismuth Subsalicylate 525 mg  
Subsalicilato de Bismuto 525 mg

**UPSET STOMACH RELIEVER  
ALIVIA EL MALESTAR ESTOMACAL  
ANTI-DIARRHEAL  
ANTI-DIARRHEICO**

**RELIEVES:**  
ALIVIA:  
Diarrhea  
Diarrea  
Nausea  
Náuseas

**Heartburn  
Acidez  
Upset Stomach  
Malestar estomacal  
Indigestion  
Indigestión**

**Actual Size  
Tamaño Actual**

**24 caplets  
comprimidos**

**sanfer<sup>®</sup>**

**CONSERVE EL EMPAQUE PARA OBTENER LA INFORMACIÓN COMPLETA DEL PRODUCTO**

**Datos Farmacológicos**  
**Beneficio del ingrediente activo (en cada comprimido)**  
Subsalicilato de Bismuto 525 mg  
Alivia el malestar estomacal / anti-diarrea

**Usos**  
■ Diarrea del viajero ■ Diarrea ■ Malestar estomacal provocado por la ingestión excesiva de alimentos y bebidas como:  
■ indigestión ■ sensación de hinchazón ■ gases ■ náuseas ■ vómitos ■ acidez

**Advertencias**  
Consulte las advertencias completas sobre el Síndrome de Reye en el empaque: los niños y adolescentes enfermos o convalecientes por varicela o síntomas similares a los de la gripe no deben usar este producto. Si al usar este producto observa cambios de comportamiento como náuseas y vómitos, consulte a su médico porque estos síntomas podrían ser un signo del síndrome de Reye, una enfermedad rara pero grave.

**Advertencia de alergias: Contiene salicilatos.**  
No usar:  
■ si es alérgico a los salicilatos (como la aspirina).  
■ si está usando otros productos con salicilatos.

**No usar si tiene:**  
■ úlcera ■ problemas de sangrado ■ heces con sangre o negras

**Consulte a un médico si presenta:**  
■ fiebre ■ morecos en los brazos

**Consulte a su médico o farmacéutico antes de usarlo, si está tomando algún medicamento para:**  
■ aliviar la sangre (anticagregante) ■ gota ■ diabetes ■ zóonias

Este producto puede causar oscurecimiento temporal (sentido de las heces) y la diarrea.

**Suspenda su uso y consulte a su médico:**  
■ si empeoran los síntomas ■ si experimenta zumbido en los oídos o pérdida de la audición ■ si la diarrea se prolonga por más de 2 días ■ si está embarazada o en período de lactancia, consulte a su médico antes de usar este medicamento.

**Manténgase fuera del alcance de los niños.** En caso de sobredosis, busque atención médica o comuníquese inmediatamente con el Centro de Toxicología.

**Datos Farmacológicos (continuación)**  
**Indicaciones**  
■ No tomar más de lo indicado.  
■ Beber abundantes líquidos para ayudar a prevenir la deshidratación causada por la diarrea.  
■ Tomar el comprimido con agua; no masticar.  
■ Adultos y niños mayores de 12 años: 1 comprimido cada 12 a 1 hora según sea necesario. No tomar más de 8 comprimidos en 24 horas. ■ No usar por más de 2 días, excepto si indica un médico. ■ Usar hasta que se detenga la diarrea, pero no por más de 2 días. ■ Niños menores de 12 años: consultar al médico.

**Información adicional**  
■ Cada comprimido contiene: calcio 45 mg, salicilato 206 mg, sodio 3 mg.

**SELLO DE SEGURIDAD: NO USAR SI NO TIENE O ESTÁ ROTO EL SELLO DE SEGURIDAD IMPRESO BAJO LA TAPA.**  
■ Almacenar a 25 °C (77 °F); permite oscilaciones entre 15 °C y 30 °C (entre 59 ° y 86 °F).  
■ No exponer al calor extremo. ■ Consultar la fecha de caducidad y el número de lote en la perla inferior. ■ Usar antes de la fecha de caducidad indicada en el envase.

**Ingredientes inactivos:** carbonato de calcio, almidón de maíz, laca de aluminio rojo #27 D&C, laca de aluminio rojo #30 D&C, estearato de magnesio, manitol, celulosa microcristalina, povidona, dióxido de silicio, almidón glabro de sodio, ácido estéarico.

**¿Tiene preguntas o comentarios?**  
Lláme al 1-(833) 604-1084 de 9:00 a.m. a 9:00 p.m. hora del centro, de lunes a viernes (Inglés-Español).

**EVIDENCIA DE MANIPULACIÓN: NO UTILIZAR SI EL SELLO DE SEGURIDAD IMPRESO DEBAJO DE LA TAPA FALTA O ESTÁ ROTO**

**Drug Facts** **KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION**

**Active ingredient (in each caplet)**  
Bismuth subsalicylate 525 mg ..... Upset stomach reliever/anti-diarrheal

**Purpose**  
relieves ■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including:  
■ indigestion ■ fullness ■ gas ■ nausea ■ heartburn ■ belching

**Warnings**  
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
Allergy alert: Contains salicylate. Do not take if you are allergic to salicylates (including aspirin).  
■ taking other salicylate products

**Do not use if you have**  
■ an ulcer ■ a bleeding problem ■ bloody or black stool

**Ask a doctor before use if you have**  
■ fever ■ mucus in the stool

**Ask a doctor or pharmacist before use if you are taking any drug for**  
■ gout ■ diabetes ■ arthritis ■ anticoagulation (thinning the blood)

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.

**Stop use and ask a doctor if**  
■ symptoms get worse ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days

**If pregnant or breast-feeding, ask a health professional before use.** Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
■ do not take more than directed ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Drug Facts (continued)**

■ swallow with water; do not chew  
■ adults and children 12 years and over: 1 caplet every 1/2 to 1 hour as needed. Do not exceed 8 caplets in 24 hours.  
■ do not use for more than 2 days unless directed by a doctor  
■ use until diarrhea stops, but not more than 2 days  
■ children under 12 years: ask a doctor

**Other information**  
■ each caplet contains: calcium 45 mg, salicylate 206 mg, sodium 3 mg  
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)  
■ avoid excessive heat  
■ see end flap for expiration date and lot number

**Inactive ingredients** calcium carbonate, corn starch, D&C red #27 aluminum lake, D&C red #30 aluminum lake, magnesium stearate, mannitol, microcrystalline cellulose, povidone, silicon dioxide, sodium starch glycolate, stearic acid

**Questions or comments?**  
Call 1-(833) 604-1084 (English and Spanish)  
8:00 AM-5:00 PM CST, Monday-Friday

Manufactured for: Laboratorios Sanfer S.A. de C.V.  
Blvd. Adolfo López Mateos No. 314  
Col. Itzacopac, CDMX, 01049, México

This product is produced in compliance with all United States regulatory and quality requirements for pharmaceutical product.

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

  
8 50036 90493 9 50844  
ORG032174908

B-1090-749-08  
ORG032174908

NO PRINTING • NO VARNISH  
LOT & EXP

**Treda Ultra 44-749**

<b>TREDA ULTRA</b>			
bismuth subsalicylate tablet			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:83393-749
<b>Route of Administration</b>	ORAL		
<b>Active Ingredient/Active Moiety</b>			
<b>Ingredient Name</b>		<b>Basis of Strength</b>	<b>Strength</b>
BISMUTH SUBSALICYLATE (INN: G2TEX51BB) (SALICYLIC ACID, INN: Q11AB74UBZ)		BISMUTH	

<b>BISMUTH SUBSALICYLATE</b> (UNII: 021EY51RRI) (SALICYLIC ACID - UNII:0414PZ4LPZ, BISMUTH CATION - UNII:ZS9CD1I8YE)	<b>BISMUTH SUBSALICYLATE</b>	525 mg
--	------------------------------	--------

### Inactive Ingredients

Ingredient Name	Strength
<b>CALCIUM CARBONATE</b> (UNII: H0G9379FGK)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>D&amp;C RED NO. 27 ALUMINUM LAKE</b> (UNII: ZK64F7XSTX)	
<b>D&amp;C RED NO. 30</b> (UNII: 2S42T2808B)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6130)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	

### Product Characteristics

<b>Color</b>	pink	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	44;749
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83393-749-08	1 in 1 CARTON	02/06/2024	
1		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	02/06/2024	

**Labeler** - Laboratorios Sanfer, S.A. de C.V. (810007732)

Revised: 2/2026

Laboratorios Sanfer, S.A. de C.V.