

LEADER INFANTS PAIN AND FEVER- acetaminophen suspension
Cardinal Health 110, LLC. dba Leader

Cardinal Health Infants' Pain & Fever Drug Facts

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
- the common cold
- flu
- headache
- sore throat
- toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has

liver disease

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

When using this product

do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed (see overdose warning)**
- **shake well before using**
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise, use age.
- push air out of syringe. Insert syringe tip into bottle opening.
- flip bottle upside down. Pull yellow part of syringe to the first dose line and then push product back into bottle.
- pull yellow part of syringe until it reaches and stays at the correct dose
- dispense liquid slowly into child’s mouth, toward inner cheek
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- replace cap tightly to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor

24-35	2-3 years	5 mL
-------	-----------	------

* or as directed by a doctor

Attention: use only enclosed syringe specifically designed for use with this product. Do not use any other dosing device.

Other information

- store at 20-25°C (68-77°F)
- **do not use if printed neckband is broken or missing**

Inactive ingredients

anhydrous citric acid, calcium sulfate, carrageenan, flavor, glycerin, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, tribasic sodium phosphate

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

LEADER™

DYE-FREE

Infants' Pain & Fever

Acetaminophen 160 mg per 5 mL

Oral Suspension

Pain Reliever / Fever Reducer

Grape Flavor

Ages 2 to 3 Years

Alcohol-Free

Aspirin-Free

Ibuprofen-Free

No Parabens

COMPARE TO INFANTS' TYLENOL® ORAL SUSPENSION

active ingredient

100% Money Back Guarantee

2 FL OZ (59 mL)

USE ONLY WITH ENCLOSED SYRINGE.
 SEE SIDE PANEL FOR MORE INFORMATION.
 PEDIATRICIAN PREFERRED DOSING SYSTEM

Drug Facts

Active ingredient (in each 5 mL) **Purpose**
 Acetaminophen 160 mg Pain reliever/fever reducer

Uses temporarily: ■ reduces fever
 ■ relieves minor aches and pains due to: ■ the common cold
 ■ flu ■ headache ■ sore throat ■ toothache

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes
 ■ more than 5 doses in 24 hours, which is the maximum daily amount
 ■ with other drugs containing acetaminophen
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if your child has ever had an allergic reaction to this product or any of its ingredients
Ask a doctor before use if your child has liver disease
Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin
When using this product do not exceed recommended dose (see overdose warning)
Stop use and ask a doctor if ■ new symptoms occur
 ■ pain gets worse or lasts more than 5 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present. These could be signs of a serious condition.
Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions
 ■ this product does not contain directions or complete warnings for adult use
 ■ do not give more than directed (see overdose warning)
 ■ shake well before using ■ mL = milliliter
 ■ find right dose on chart. If possible, use weight to dose; otherwise, use age.
 ■ push air out of syringe. Insert syringe tip into bottle opening.
 ■ flip bottle upside down. Pull yellow part of syringe to the first dose line and then push product back into bottle. ▶

Drug Facts (continued)

■ pull yellow part of syringe until it reaches and stays at the correct dose
 ■ dispense liquid slowly into child's mouth, toward inner cheek
 ■ repeat dose every 4 hours while symptoms last
 ■ do not give more than 5 times in 24 hours
 ■ replace cap tightly to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL

* or as directed by a doctor
Attention: use only enclosed syringe specifically designed for use with this product. Do not use any other dosing device.

Other information
 ■ store at 20-25°C (68-77°F)
 ■ do not use if printed neckband is broken or missing

Inactive ingredients anhydrous citric acid, calcium sulfate, carrageenan, flavor, glycerin, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, tribasic sodium phosphate

Questions or comments?
 1-800-719-9260

**This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of infants' Tylenol® Oral Suspension.

©2024 Cardinal Health. All Rights Reserved. CARDINAL HEALTH, the Cardinal Health Logo, ESSENTIAL TO CARE, LEADER, and the LEADER Logo are trademarks or registered trademarks of Cardinal Health. All other marks are the property of their respective owners.

✓ All LEADER™ Brand Products Have A 100% Money Back Guarantee
 Return to place of purchase if not satisfied.

LEADER²

Infants' Pain & Fever
 Acetaminophen 160 mg per 5 mL
 Oral Suspension
 Pain Reliever / Fever Reducer

NDC 70000-0674-1

DYE-FREE

Infants' Pain & Fever
 Acetaminophen 160 mg per 5 mL
 Oral Suspension
 Pain Reliever / Fever Reducer

Grape Flavor

Ages 2 to 3 Years

Alcohol-Free
 Aspirin-Free
 Ibuprofen-Free
 No Parabens

COMPARE TO INFANTS' TYLENOL® ORAL SUSPENSION active ingredient**

100% Money Back Guarantee

2 FL OZ (59 mL)

USE ONLY WITH ENCLOSED SYRINGE. SEE SIDE PANEL FOR MORE INFORMATION. PEDIATRICIAN PREFERRED DOSING SYSTEM

CIN 5899166 REV. 03/24

0 96295 14339 3

76616 E9 C 1

IMPORTANT:
 Please see dosing chart or ask a doctor for your child's correct dose.

STEP 1. See *Directions* for the correct dose for your child.

STEP 2. Shake well before using.

STEP 3. Push air out of syringe. Insert syringe tip into bottle opening.

STEP 4. Flip bottle upside down. Pull yellow part of syringe to the first dose line and then push product back into bottle.

STEP 5. Pull yellow part of syringe until it reaches and stays at the correct dose.

STEP 6. Dispense liquid slowly into child's mouth.

DISTRIBUTED BY CARDINAL HEALTH
 DUBLIN, OHIO 43017
 www.myleader.com
 1-800-200-6315
 Essential to Care™ since 1979

LEADER INFANTS PAIN AND FEVER

acetaminophen suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70000-0674
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CALCIUM SULFATE, UNSPECIFIED FORM (UNII: WAT0DDB505)	
CARRAGEENAN (UNII: 5C69YCD2YJ)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
SODIUM PHOSPHATE, TRIBASIC (UNII: A752Q30A6X)	

Product Characteristics

Color	WHITE	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70000-0674-1	1 in 1 CARTON	05/06/2024	
1		59 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/06/2024	

Labeler - Cardinal Health 110, LLC. dba Leader (063997360)

