

CHILDRENS GILTUSS D- childrens giltuss d syrup
Gil Pharmaceutical Corp

Children's Giltuss-D
ALLERGY & CONGESTION
ANTIHISTAMINIC AND NASAL DECONGESTANT

Drug Facts

Active ingredients (in each 5 mL) Purpose

Chlorpheniramine maleate 1.75 mg Antihistamine

Phenylephrine HCl 5 mg Nasal decongestant

Uses

Temporarily relieves these Symptoms due to hay fever (allergic rhinitis) or other upper-respiratory allergies:

- runny nose
- Sneezing
- Itchy, watery eyes
- nasal congestion
- itching of the nose or throat
- sinus congestion and pressure

Warnings

Do not use

- to sedate a child or to make a child sleepy
- if the child is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug.

If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if the child has

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- glaucoma
- a breathing problem or persistent or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if the child is

- taking sedatives or tranquilizers

When using this product

- **do not use more than directed**
- may cause drowsiness
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- if the child gets nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **not take more than 6 doses in any 24-hour period**
- **use enclosed dosing cup**
- **keep dosage cup for use with this product**
- **mL = milliliter**

Children under 6 years of age	Consult a Doctor
children 6 to under 12 years of age	5 mL every 4 hours
Adults and children 12 years and over	10 mL every 4 hours

Other information

- store at room temperature 15°-30°C (59°-86°F)
- close cap tightly
- **SAVE CARTON FOR COMPLETE DRUG FACTS**

Inactive ingredients

Citric Acid, Flavor, Glycerin, Methylparaben, Propylene Glycol, Propylparaben, Purified Water, Sodium Carboxymethylcellulose, Sucralose.

Questions or comments?

Call 1-787-848-9114, Monday-Friday 9AM-5PM EST. Call your doctor for medical advice in the event of side effects.

Label revised: 04/24

**MANUFACTURED FOR
GIL PHARMACEUTICAL CORP.
PONCE, PUERTO RICO 00717-1565**

PRINCIPAL DISPLAY PANEL

Children's Giltuss-D ALLERGY & CONGESTION - NDC 58552-148-04 - Bottle Label

<p>Drug Facts</p>	<p>Active ingredients (in each 5 mL)</p> <p>Chlorpheniramine maleate, 1.75 mg..... Antihistamine Phenylephrine HCl, 5 mg..... Nasal decongestant</p>	<p>Purposes</p>		<p>When using this product • do not use more than directed • may cause drowsiness • avoid alcoholic beverages • alcohol, sedatives, and tranquilizers may increase drowsiness • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • excitability may occur, especially in children</p>							
<p>Uses Temporarily relieves these symptoms due to hay fever (allergic rhinitis) or other upper-respiratory allergies:</p> <ul style="list-style-type: none"> runny nose • Sneezing • Itchy, watery eyes • nasal congestion itching of the nose or throat • sinus congestion and pressure 	<p>Warnings</p> <p>Do not use • to sedate a child or to make a child sleepy • if the child is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p>	<p>Ask a doctor before use if the child has:</p> <ul style="list-style-type: none"> heart disease • high blood pressure • thyroid disease • diabetes trouble urinating due to an enlarged prostate gland • glaucoma a breathing problem or persistent or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema. 	<p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>	<p>Stop use and ask a doctor if • if the child gets nervous, dizzy, or sleepless • symptoms do not get better within 7 days or are accompanied by fever</p>	<p>MANUFACTURED FOR GIL PHARMACEUTICAL CORP. PONCE, PUERTO RICO 00717-1565</p>						
<p>Ask a doctor or pharmacist before use if the child is taking sedatives or tranquilizers</p>	<p>Ask a doctor or pharmacist before use if the child is taking sedatives or tranquilizers</p>	<p>Ask a doctor or pharmacist before use if the child is taking sedatives or tranquilizers</p>	<p>Directions • do not take more than 6 doses in any 24-hour period • use enclosed dosing cup • keep dosage cup for use with this product • mL = milliliter</p>	<table border="1"> <tr> <td>children under 6 years</td> <td>Consult a Doctor</td> </tr> <tr> <td>children 6 to under 12 years</td> <td>5 mL every 4 hours</td> </tr> <tr> <td>Adults and children 12 years and over</td> <td>10 mL every 4 hours</td> </tr> </table>	children under 6 years	Consult a Doctor	children 6 to under 12 years	5 mL every 4 hours	Adults and children 12 years and over	10 mL every 4 hours	<p>For most recent product information, visit www.giltuss.com Label revised: 02/23</p>
children under 6 years	Consult a Doctor										
children 6 to under 12 years	5 mL every 4 hours										
Adults and children 12 years and over	10 mL every 4 hours										
<p>Ask a doctor or pharmacist before use if the child is taking sedatives or tranquilizers</p>	<p>Ask a doctor or pharmacist before use if the child is taking sedatives or tranquilizers</p>	<p>Ask a doctor or pharmacist before use if the child is taking sedatives or tranquilizers</p>	<p>Other information • store at room temperature 15°-30°C (59°-86°F) • close cap tightly • SAVE CARTON FOR COMPLETE DRUG FACTS</p>	<p>Inactive ingredients Citric Acid, Flavor, Glycerin, Methylparaben, Propylene Glycol, Propylparaben, Purified Water, Sodium Carboxymethylcellulose, Sucralose.</p>	<p>Questions or comments? Call 1-787-348-9114, Monday-Friday 9AM-5PM EST. Call your doctor for medical advice in the event of side effects.</p>						

Children's Giltuss-D ALLERGY & CONGESTION - NDC 58552-148-04 - Carton Label

DO NOT USE IF SAFETY SEAL IS BROKEN OR MISSING. VERIFICAR SI LA SELLA DE SEGURIDAD ESTÁ INTACTA.

ANTIHISTAMINE AND NASAL DECONGESTANT
ANTIHIISTAMÍNICO Y DESCONGESTIVANTE NASAL

NO PSEUDOEPHEDRINE Children Niños 6-12

Children's Giltuss-D ALLERGY & CONGESTION ALERGIJA Y CONGESTIÓN

FAST ANTIHISTAMINE EFFICACY TO RELIEVE SNEEZING, RUNNY NOSE, AND ITCHY & WATERY EYES, ITCHY THROAT OR NOSE
RÁPIDA EFICACIA ANTIHIISTAMÍNICA PARA ALIVIAR LOS ESTORNIDOS, CORRIENTE NASAL, Y OJOS LLOROSOS, PICOR DE GARGANTA O NASAL

DECONGESTANT EFFICACY TO RELIEVE SINUS PRESSURE & STUFFY NOSE
EFICACIA DESCONGESTIVANTE PARA EL ALIVIO DE PRESIÓN POR SINUSITIS Y NASAL TAPADA

Drug Facts

Active ingredients (in each 5 mL)

Chlorpheniramine maleate, 1.75 mg..... Antihistamine
Phenylephrine HCl, 5 mg..... Nasal decongestant

Purposes

Temporarily relieves these symptoms due to hay fever (allergic rhinitis) or other upper-respiratory allergies:

- sneezing
- runny nose
- itching of the nose or throat
- sinus congestion and pressure

Warnings

Do not use • to sedate a child or to make a child sleepy • if the child is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if the child has:

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- glaucoma
- a breathing problem or persistent or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema.

Ask a doctor or pharmacist before use if the child is taking sedatives or tranquilizers

Directions

Do not take more than 6 doses in any 24-hour period • use enclosed dosing cup • keep dosage cup for use with this product • mL = milliliter

children under 6 years of age Consult a Doctor
children 6 to under 12 years 5 mL every 4 hours
Adults and children 12 years and over 10 mL every 4 hours

Other information

Store at room temperature 15°-30°C (59°-86°F) • close cap tightly • SAVE CARTON FOR COMPLETE DRUG FACTS

Inactive ingredients
Citric Acid, Flavor, Glycerin, Methylparaben, Propylene Glycol, Propylparaben, Purified Water, Sodium Carboxymethylcellulose, Sucralose.

Questions or comments?
Call 1-787-348-9114, Monday-Friday 9AM-5PM EST. Call your doctor for medical advice in the event of side effects.

YEAST FREE, GLUTEN FREE SIN VEGANURA, SIN GLUTEN

Dye Free, Preservative Free, Sodium Free, Sugar Free
Sin colorantes, Sin Conservantes, Sin sodio, Sin azúcar

Tutti Frutti Flavor Sabor a Tutti Frutti

4 FL OZ (118 mL)

3 58552 14804 7

MANUFACTURED FOR
GIL PHARMACEUTICAL CORP.
PONCE, PUERTO RICO 00717-1565
giltuss.com

CHILDRENS GILTUSS D
childrens giltuss d syrup

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58552-148
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	1.75 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE	5 mg

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID ACETATE (UNII: DSO12WL7AU)	
GLYCERIN (UNII: PDC6A3C0OX)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	TUTTI FRUTTI	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58552-148-04	1 in 1 CARTON	04/11/2024	
1		1 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/11/2024	

Labeler - Gil Pharmaceutical Corp (176826592)

