

PAIN RELIEF- acetaminophen tablet, coated
CHAIN DRUG CONSORTIUM

1092-PRV-2020-0826

Drug Facts

Active ingredient (in each tablet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or

pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none">▪ take 2 tablets every 6 hours while symptoms last▪ swallow whole – do not crush, chew, or dissolve▪ do not take more than 6 tablets in 24 hours, unless directed by a doctor▪ do not use for more than 10 days unless directed by a doctor
children under 12 years	<ul style="list-style-type: none">▪ ask a doctor

Other information

- store between 20-25°C (68-77°F)
- retain carton for complete product information

Inactive ingredients

acesulfame potassium, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

1-844-705-4384

PRINCIPAL DISPLAY PANEL

Premier Value®

COMPARE TO THE ACTIVE INGREDIENT IN TYLENOL® EXTRA STRENGTH EZ TABS†
EXTRA STRENGTH

Pain Relief

For Adults

ACETAMINOPHEN • EASY TABS

PAIN RELIEVER/FEVER REDUCER

Easy to Swallow, Sweet Coated

50 TABLETS, 500 MG EACH

INK AND COATING FREE
FOR LOT AND
EXPIRATION STAMPING



DO NOT USE IF IMPRINTED SEAL
UNDER CAP IS BROKEN OR MISSING

This product is not manufactured or distributed by Micelle II Consumer Healthcare, distributor of Tylenol® Extra Strength EZ Tabs. F11092039REV_B3

Drug Facts

Active ingredient (in each tablet)
Acetaminophen 500 mg — Pain reliever/fever reducer

Purpose

- Uses**
- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - minor pain of arthritis
 - muscular aches
 - toothache
 - menstrual and menstrual cramps
 - temporarily reduces fever

- Warnings**
- Liver warnings:** This product contains acetaminophen. Severe liver damage may occur if you take
 - more than 4,000 mg of acetaminophen in 24 hours
 - with other drugs containing acetaminophen
 - 3 or more alcoholic drinks every day while using this product.

- Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:
 - skin redness
 - hives
 - rash
 If a skin reaction occurs, stop use and seek medical help right away.

- Do not use**
- with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 - if you are allergic to acetaminophen or any of the inactive ingredients in this product.
- Ask a doctor before use if you have liver disease. ▶

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

- Stop use and ask a doctor if**
- pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.


- Directions** — do not take more than directed (see overdose warning)
- adults and children 12 years and over
 - take 2 tablets every 6 hours while symptoms last
 - swallow whole — do not crush, chew, or dissolve
 - do not take more than 6 tablets in 24 hours, unless directed by a doctor
 - do not use for more than 10 days unless directed by a doctor
 - children under 12 years
 - ask a doctor

- Other Information**
- store between 20-25°C (68-77°F)
 - retain carton for complete product information

Drug Facts (continued)

Inactive Ingredients acetaminophen potassium, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, polyvinylpyrrolidone K90, polyethylene glycol, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?
1-844-705-4384



NC

NC

NC

PAIN RELIEF acetaminophen tablet, coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68016-502
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	red	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	A92
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68016-502-50	1 in 1 CARTON	01/01/2008	05/31/2026
1		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:68016-502-00	1 in 1 CARTON	01/01/2008	06/30/2026
2		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:68016-502-24	1 in 1 CARTON	06/22/2018	05/31/2024
3		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/01/2008	06/30/2026

Labeler - CHAIN DRUG CONSORTIUM (101668460)

Revised: 10/2024

CHAIN DRUG CONSORTIUM