

PREP-SAN ULTRA- lactic acid, linear dodecylbenzene sulfonic acid, sulfonated oleic acid solution
Ecolab Inc.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Danger

Causes severe skin burns and eye damage.

Wash skin thoroughly after handling. Wear protective gloves/ protective clothing/ eye protection/ face protection. Do not mix with bleach or other chlorinated products - will cause chlorine gas.

IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. IF ON SKIN (or hair): Remove/ Take off immediately all contaminated clothing. Rinse skin with water/ shower.

IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER or doctor/ physician. Wash contaminated clothing before reuse.

DIRECTIONS:

Consult your Ecolab representative for specific use instructions and recommended dispensing equipment.

PREP-SAN ULTRA MUST BE DILUTED BEFORE USE

DO NOT MIX WITH ANYTHING BUT COOL, POTABLE WATER

FOR USE ONLY IN A DISPENSING SYSTEM RECOMMENDED BY ECOLAB

Dilution Parameter: A dilute solution of a 0.5% lactic acid and 1.0% linear alkylate sulfonates teat dip will consist of 1 part Prep-San Ultra and 9 parts potable water by volume.

IMPORTANT: Do not mix Prep-San Ultra or diluted Prep-San Ultra with any other products unless specified in the product use instructions. If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for Prep-San Ultra. If product in dip cup becomes visibly dirty, discard contents and replenish with diluted, ready-to-use product. Do not reuse or return any unused product to the original container.

Before Milking: Wash teats thoroughly just prior to next milking with appropriate udder wash solution or pre-milking teat dip to avoid contamination of milk. Teats should then be dried with single-service towels. Use proper procedures for udder washing or pre-dipping.

Application of Diluted Prep-San Ultra

Directions for Teat Dipping

Pre-Milk Dipping: Fill teat dip cup with diluted Prep-San Ultra. Before each cow is

milked, dip each teat fulllength into the teat dip cup containing diluted Prep-San Ultra. Wipe teats dry after dipping using single-service towels to avoid contamination of milk.

Post-Milk Dipping: Immediately after each milking, dip each teat full-length into the teat dip cup containing diluted Prep-San Ultra. Allow to air dry. Do not wipe. Always use fresh, diluted Prep-San Ultra. Do not turn cows out in freezing weather until teat dip is completely dry.

Directions for Teat Spraying

Pre-Milk Spraying: Before each cow is milked, spray entire teat with diluted Prep-San Ultra. Wipe teats after spraying using single-service towels to avoid contamination of milk.

Post-Milk Spraying: Immediately after each milking, spray entire teat with diluted Prep-San Ultra. Allow to air dry. Do not wipe. Always use fresh, diluted Prep-San Ultra. Do not turn cows out in freezing weather until diluted Prep-San Ultra is completely dry.

Expanded Usage: When freshening cows, begin dipping teats twice daily for about 10 days before calving.

PRECAUTION: Prep-San Ultra is not intended to cure or help the healing of chapped or irritated teats. As with any germicide, irritation or sensitization may occur in sensitive animals. In case of teat irritation or chapping, have the condition examined and, if necessary, treated by a veterinarian.

NOTE: Use blended Prep-San Ultra within six (6) months of blended date.

READ SAFETY DATA SHEET (SDS) BEFORE USING THIS PRODUCT

EMERGENCY HEALTH INFORMATION: 1 800 328 0026. If located outside the United States and Canada, call collect 1 651 222 5352 (number is in the US).

A & L Laboratories

Prep-San™ Ultra

Teat Dip Concentrate

An aid in reducing the spread of organisms which may cause mastitis

ACTIVE INGREDIENTS:

Lactic acid.....4.2%

Linear alkylate sulfonates.....8.2%

(includes 5.0% linear dodecyl benzene sulfonic acid and 3.2% sulfonated oleic acid)

INERT INGREDIENTS:.....47.6%

SKIN CONDITIONING SYSTEM:

(contains sorbitol)..... 40.0%

TOTAL:.....100.0%

Manufactured by

Ecolab · 1 Ecolab Place · St. Paul, MN 55102 USA · tel: 1 800 392 3392

© 2017 Ecolab USA Inc · All rights reserved

Made in U.S.A. | Hecho en EE.UU.

762092/5301/0517

56.8 L (15 US GAL) 6301638



An aid in reducing the spread of organisms which may cause mastitis

ACTIVE INGREDIENTS:	
Lactic acid	4.2%
Linear alkylate sulfonates	3.2%
(Includes 5.0% linear dodecyl benzene sulfonic acid and 3.2% sulfonated oleic acid)	
INERT INGREDIENTS:	
	47.6%
SKIN CONDITIONING SYSTEM:	
(contains sorbitol)	40.0%
TOTAL:	100.0%

FOR INDUSTRIAL USE ONLY

DIRECTIONS:

Consult your Ecolab representative for specific use instructions and recommended dispensing equipment.
PREP-SAN ULTRA MUST BE DILUTED BEFORE USE
DO NOT MIX WITH ANYTHING BUT COOL, POTABLE WATER
FOR USE ONLY IN A DISPENSING SYSTEM RECOMMENDED BY ECOLAB
Dilution Parameter: A dilute solution of a 0.5% lactic acid and 1.0% linear alkylate sulfonates teat dip will consist of 1 part Prep-San Ultra and 9 parts potable water by volume.
IMPORTANT: Do not mix Prep-San Ultra or diluted Prep-San Ultra with any other products unless specified in the product use instructions. If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for Prep-San Ultra. If product in dip cup becomes visibly dirty, discard contents and replenish with diluted, ready-to-use product. Do not reuse or return any unused product to the original container.
Before Milking: Wash teats thoroughly just prior to next milking with appropriate udder wash solution or pre-milking teat dip to avoid contamination of milk. Teats should then be dried with single-service towels. Use proper procedures for udder washing or pre-dipping.
Application of Diluted Prep-San Ultra
Directions for Teat Dipping
Pre-Milk Dipping: Fill teat dip cup with diluted Prep-San Ultra. Before each cow is milked, dip each teat full-length into the teat dip cup containing diluted Prep-San Ultra. Wipe teats dry after dipping using single-service towels to avoid contamination of milk.
Post-Milk Dipping: Immediately after each milking, dip each teat full-length into the teat dip cup containing diluted Prep-San Ultra. Allow to air dry. Do not wipe. Always use fresh, diluted Prep-San Ultra. Do not turn cows out in freezing weather until teat dip is completely dry.
Directions for Teat Spraying
Pre-Milk Spraying: Before each cow is milked, spray entire teat with diluted Prep-San Ultra. Wipe teats after spraying using single-service towels to avoid contamination of milk.
Post-Milk Spraying: Immediately after each milking, spray entire teat with diluted Prep-San Ultra. Allow to air dry. Do not wipe. Always use fresh, diluted Prep-San Ultra. Do not turn cows out in freezing weather until diluted Prep-San Ultra is completely dry.
Expanded Usage: When freshening cows, begin dipping teats twice daily for about 10 days before calving.
PRECAUTION: Prep-San Ultra is not intended to cure or help the healing of chapped or irritated teats. As with any germicide, irritation or sensitization may occur in sensitive animals. In case of teat irritation or chapping, have the condition examined and, if necessary, treated by a veterinarian.
NOTE: Use blended Prep-San Ultra within six (6) months of blended date.

KEEP OUT OF REACH OF CHILDREN

NOT FOR HUMAN USE

Danger

Causes severe skin burns and eye damage.

Wash skin thoroughly after handling. Wear protective gloves/protective clothing/eye protection/face protection. Do not mix with bleach or other chlorinated products - will cause chlorine gas.

IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. IF ON SKIN (or hair): Remove/ Take off immediately all contaminated clothing. Rinse skin with water/ shower. IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER or doctor/physician. Wash contaminated clothing before reuse.

READ SAFETY DATA SHEET (SDS) BEFORE USING THIS PRODUCT

EMERGENCY HEALTH INFORMATION: 1 800 328 0426. If located outside the United States and Canada, call collect 1 651 222 5352 (number is in the US).

Manufactured by
 Ecolab · 1 Ecolab Place · St. Paul, MN 55102 USA · tel: 1 800 392 3392
 © 2017 Ecolab USA Inc · All rights reserved
 Made in U.S.A. | Hecho en EE.UU.
 762092/5301/0517

Ayuda a evitar la proliferación de organismos que pueden causar mastitis

PARA USO INDUSTRIAL SOLAMENTE

INSTRUCCIONES:

Consulte con su Representante de Ecolab para obtener instrucciones específicas sobre el uso y el equipo dispensador recomendado.
PREP-SAN ULTRA DILUIR ANTES DE USAR
NO MEZCLAR CON NADA QUE NO SEA AGUA POTABLE FRÍA
USAR ÚNICAMENTE EN UN SISTEMA DE DOSIFICACIÓN RECOMENDADO POR ECOLAB
Parámetro de dilución: Una solución diluida de baño antiséptico para tetillas de 0.5% de ácido láctico y 1% de sulfonatos alquilo lineales estará compuesta por 1 parte de concentrado Prep-San Ultra y 9 partes de agua potable por volumen.
IMPORTANTE: No mezcle Prep-San Ultra o Prep-San Ultra diluido con otros productos a menos que se indique en las instrucciones de uso del producto. Si el producto fuese transferido desde este envase a cualquier otro, asegúrese de que ese otro envase haya sido íntegramente prelavado y que contenga el etiquetamiento del envase de Prep-San Ultra.
 Si el producto en el recipiente para realizar el baño está visiblemente sucio, deseché el contenido y rellene el recipiente con producto diluido y listo para usar. No vuelva a utilizar el mismo producto, ni vuelva a colocar el producto sin usar dentro del envase original.
Antes de ordeñar: Lave los pezones completamente justo antes de ordeñar con una solución de lavado apropiada para ubres o baño antiséptico para pezones previo al ordeño, o rocíe el pezón para evitar la contaminación de la leche. Posteriormente, los pezones se deben secar con toallas de servicio único o descartables. Utilice el procedimiento adecuado para el lavado o el prelavado antiséptico de las ubres.
Aplicación de dilución Prep-San Ultra
Instrucciones para realizar el sellado del pezón
Sellado del pezón previo al ordeño: Llene el recipiente para el baño de pezones con Prep-San Ultra diluido. Antes de ordeñar cada vaca, bañe toda la superficie del pezón en el recipiente que contiene Prep-San Ultra diluido. Seque los pezones después de realizar el sellado, utilizando toallas desechables para evitar la contaminación de la leche.
Sellado del pezón posterior al ordeño: Inmediatamente después de ordeñar cada vaca, bañe toda la superficie del pezón en el recipiente que contiene Prep-San Ultra diluido. Deje secar al aire. No utilice paños. Siempre utilice Prep-San Ultra fresco y diluido. Hasta que Prep-San Ultra no se haya secado completamente, no saque las vacas al exterior en climas fríos.
Instrucciones para el Rociado de los Pezones
Rociado previo al ordeño: Antes de ordeñar cada vaca, rocíe toda la superficie del pezón con Prep-San Ultra diluido. Seque los pezones después de rociarlos utilizando toallas de un solo servicio para evitar la contaminación con la leche.
Rociado posterior al ordeño: Inmediatamente después de ordeñar cada vaca, rocíe toda la superficie del pezón con Prep-San Ultra diluido. Deje secar al aire. No utilice paños. Siempre utilice Prep-San Ultra fresco y diluido. No deje las vacas en un clima helado antes de que Prep-San Ultra diluido se haya secado por completo.
Uso Expandido: Al refrescar a las vacas, comience con un baño antiséptico de sus pezones dos veces por día por aproximadamente 10 días antes de parir.

PRECAUCIÓN: Prep-San Ultra no está diseñado para curar o ayudar con la cicatrización de los pezones agrietados o irritados. Como cualquier otro germicida, puede provocar irritación o sensibilización en animales sensibles. En caso de irritación o agrietamiento de los pezones, haga examinar dichas afecciones y, de ser necesario, hágalas tratar con un veterinario.
AVISO: Utilice Prep-San Ultra mezclado dentro de los seis (6) meses a partir de la fecha de mezclado.

MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS

NO ES PARA USA HUMANO

Peligro

Provoca graves quemaduras en la piel y lesiones oculares.

Lave la piel minuciosamente después del manejo. Usar guantes/ropa de protección/equipo de protección para los ojos/la cara. No se mezcle con el blanqueador u otros productos tratados con cloro - causará el gas de cloro.

EN CASO DE INGESTIÓN: Enjuagar la boca. NO provocar el vómito. EN CASO DE CONTACTO CON LA PIEL (o el pelo): Quitar inmediatamente la ropa contaminada. Enjuagar la piel con agua/ ducharse. EN CASO DE INHALACIÓN: Transportar la víctima al aire libre y mantenerla en reposo en una posición que le facilite la respiración. EN CASO DE CONTACTO CON LOS OJOS: Enjuagar con agua cuidadosamente durante varios minutos. Quitar las lentes de contacto cuando estén presentes y pueda hacerse con facilidad. Proseguir con el lavado. Llamar inmediatamente a un CENTRO DE TOXICOLOGÍA o a un médico. Lave la ropa contaminada antes de volverla a usar.

LEA LA FICHA DE DATOS DE SEGURIDAD ANTES DE USAR EL PRODUCTO
INFORMACIÓN DE EMERGENCIA SOBRE SALUD: 1 800 328 0426. Si está fuera de los Estados Unidos y Canadá, llame a cobro revertido al 1 651 222 5352 (número en los EE.UU.).

56.8 L (15 US GAL)

6301638



PREP-SAN ULTRA

lactic acid, linear dodecylbenzene sulfonic acid, sulfonated oleic acid solution

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:47593-547
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LACTIC ACID (UNII: 33X04XA5AT) (LACTIC ACID - UNII:33X04XA5AT)	LACTIC ACID	42 mg in 1 mL

DODECYLBENZENESULFONIC ACID (UNII: 60NSK897G9) (DODECYLBENZENESULFONIC ACID - UNII:60NSK897G9)	DODECYLBENZENESULFONIC ACID	50 mg in 1 mL
SULFONATED OLEIC ACID (UNII: DDS68Y21OL) (SULFONATED OLEIC ACID - UNII:DDS68Y21OL)	SULFONATED OLEIC ACID	32 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47593-547-17	56800 mL in 1 DRUM		
2	NDC:47593-547-21	208000 mL in 1 DRUM		
3	NDC:47593-547-44	1150000 mL in 1 CONTAINER		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		06/11/2015	

Labeler - Ecolab Inc. (006154611)

Revised: 10/2025

Ecolab Inc.