

LOW DOSE ASPIRIN- aspirin tablet, coated

Akron Phama

Aspirin

Low Dose

Drug Facts

Active Ingredient (in each tablet)

Aspirin 81 mg (NSAID)*

*** nonsteroidal anti-inflammatory drug**

***Purpose* Pain reliever**

Uses

- temporarily relief of minor aches and pains or as recommended by your doctor.
Because of its delayed release action, this product will not provide fast relieve of headache or symptoms needing immediate relief.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because the symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- shock
- facial swelling
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. This chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription of nonprescription NSAIDS [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin, or any other pain reliever/fever reducer

Ask a doctor before use if

- you have asthma
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn

- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
- under a doctor's care for any serious condition
- taking any other drug

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- have bloody or black stools
- vomit blood
- have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- ringing in the ears or loss of hearing occurs
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- redness or swelling present in the painful area
- new symptoms occur.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- drink a full glass of water with each dose
- **adults and children 12 years and over:**
- take 4 to 8 tablets every 4 hours while symptoms last
- do not take more than 48 tablets in 24 hours unless directed by a doctor
- **children under 12 years:** ask a doctor

Other information

- store at room temperature 15 °-30 °C(59 °-86 °F)
- read all product information before using

Inactive ingredients

anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, FD&C yellow#10 lake, FD&C yellow#6 lake, iron oxide ochre, methacrylic acid copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, talc, titanium dioxide, triethyl citrate

Questions or comments? Call toll-free 1-877-225-6999

Monday to Friday 9AM to 5PM

Akrion Pharma

NDC 71399-8627-2

Compare to Bayer® Low Dose Aspirin active ingredient*

Low Dose

Aspirin

81 mg

Pain Reliever (NSAID)

Safety Coated Aspirin Regimen

120 Tablets

Drug Facts

Active ingredient (in each tablet)	Purpose
Aspirin 81 mg (NSAID)*	Pain reliever
* nonsteroidal anti-inflammatory drug	

Uses

■ temporarily relief of minor aches and pains or as recommended by your doctor. **Because of its delayed release action, this product will not provide fast relief of headache or symptoms needing immediate relief.**

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ shock ■ facial swelling ■ asthma

(wheezing). **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed
Do not use ■ if you are allergic to aspirin, or any other pain reliever/fever reducer

DRUG FACTS CONTINUED on back of the label 

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL
UNDER CAP IS BROKEN OR MISSING**

***This product is not manufactured or distributed by Bayer HealthCare LLC.,
owner of the registered trademark Bayer® Low Dose Aspirin.**

PEEL HERE 

Akron Pharma

Manufactured for:
Akron Pharma, Inc.
Fairfield, NJ-07004
www.akronpharma.com
Made in USA 
Rev.:12/20



Drug Facts (Continued)

Ask a doctor before use if

- you have asthma
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
- under a doctor's care for any serious condition
- taking any other drug

Stop use and ask a doctor if

- you experience any of the following signs of stomach

bleeding: ■ feel faint ■ have bloody or black stools ■ vomit blood ■ have stomach pain that does not get better

■ an allergic reaction occurs.

Seek medical help right away.

■ ringing in the ears or loss of hearing occurs

■ pain gets worse or lasts for more than 10 days

■ fever gets worse or lasts for more than 3 days

■ redness or swelling present in the painful area

■ new symptoms occur.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ drink a full glass of water with each dose

■ **adults and children 12 years and over:**

■ take 4 to 8 tablets every 4 hours while symptoms last

■ do not take more than 48 tablets in 24 hours unless directed by a doctor

■ **children under 12 years:** ask a doctor.

Other information

■ store at room temperature 15°-30°C (59°-86°F)

■ read all product information before using

Inactive ingredients

anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, FD&C yellow#10 al lake, FD&C yellow#6 al lake, iron oxide ochre, methacrylic acid copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, talc, titanium dioxide, triethyl citrate

Questions or comments? Call toll-free 1-877-225-6999

Monday to Friday 9AM to 5PM

LOW DOSE ASPIRIN

aspirin tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:71399-8627
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
METHACRYLIC ACID - ETHYL ACRYLATE COPOLYMER (4500 MPA.S) (UNII: T967IEU43C)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	A1
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71399-8627-1	1000 in 1 BOTTLE; Type 0: Not a Combination Product	05/12/2021	
2	NDC:71399-8627-2	120 in 1 BOTTLE; Type 0: Not a Combination Product	05/12/2021	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/12/2021	

Labeler - Akron Phama (067878881)

Revised: 7/2025

Akron Phama