## BACIGUENT - bacitracin ointment Woodward Pharma Services LLC

-----

### **BACIGUENT (bacitracin) Ophthalmic Ointment USP**

# STERILE Rx Only

#### **DESCRIPTION:**

Each gram of ointment contains 500 units of bacitracin in a low melting special base containing White Petrolatum and Mineral Oil.

#### **CLINICAL PHARMACOLOGY:**

The antibiotic, bacitracin, exerts a profound action against many gram-positive pathogens, including the common Streptococci and Staphlococci. It is also destructive for certain gram-negative organisms. It is ineffective against fungi.

### **INDICATIONS AND USAGE:**

For the treatment of superficial ocular infections involving the conjunctiva and/or cornea caused by bacitracin susceptible organisms.

#### **CONTRAINDICATIONS:**

This product should not be used in patients with a history of hypersensitivity to bacitracin.

#### **PRECAUTIONS:**

Baciguent<sup>®</sup> ophthalmic ointment should not be used in deep-seated ocular infections or in those that are likely to become systemic. The prolonged use of antibiotic containing preparations may result in overgrowth of nonsusceptible organisms particularly fungi. If new infections develop during treatment appropriate antibiotic or chemotherapy should be instituted.

#### **ADVERSE REACTIONS:**

Bacitracin has such a low incidence of allergenicity that for all practical purposes side reactions are practically non-existent. However, if such reaction should occur, therapy should be discontinued.

To report SUSPECTED ADVERSE REACTIONS, contact Woodward Pharma Services LLC at 1-844-200-7910 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

### **DOSAGE AND ADMINISTRATION:**

The ointment should be applied directly into the conjunctival sac 1 to 3 times daily. In blepharitis all scales and crusts should be carefully removed and the ointment then spread uniformly over the lid margins. Patients should be instructed to take appropriate measures to avoid gross contamination of the ointment when applying the ointment directly to the infected eye.

#### **HOW SUPPLIED:**

NDC 69794-880-013-1 g sterile tamper evident tubes with ophthalmic tip.

NDC 69784-880-35 3.5 g (1/8 oz.) sterile tamper evident tubes with ophthalmic tip.

Store at 20°-25°C (68°-77°F) [see USP Controlled Room Temperature].

Manufactured For Woodward Pharma Services LLC Wixom, MI 48393

Rev 11-19 5G000 CJ J1

# PACKAGE/LABEL PRINCIPAL DISPLAY PANEL - 1 g Label

Rx Only

NDC 69784-880-35

 $Baciguent^{\hbox{\scriptsize $\mathbb{R}$}}$ 

(Bacitracin Ophthalmic Ointment, USP)

**STERILE** 

NET WT 3.5 g (1/8 oz)

USUAL DOSAGE: Apply 1 to 3 times daily. See package insert for full prescribing information.

Each gram contains: 500 units of bacitracin in a base of white petrolatum and mineral oil.

KEEP TIGHTLY CLOSED

Store at 20° to 25°C (68°-77°F) [see USP Controlled Room Temperature].

KEEP OUT OF REACH OF CHILDREN



## **BACIGUENT**

bacitracin ointment

Prod	net	Info	иma	tion
PIOU			ппп	

Product Type HUMAN PRESCRIPTION DRUG Item Code (Source) NDC:69784-880

Route of Administration OPHTHALMIC

## **Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
BACITRACIN (UNII: 58 H6 RWO 52I) (BACITRACIN - UNII: 58 H6 RWO 52I)	BACITRACIN	500 [USP'U] in 1 g

## **Inactive Ingredients**

Ingredient Name	Strength
PETROLATUM (UNII: 4T6H12BN9U)	
MINERAL OIL (UNII: T5L8T28FGP)	

P	Packaging				
#	Item Code	Package Description	<b>Marketing Start Date</b>	Marketing End Date	
1	NDC:69784-880-01	3 in 1 CARTON	04/15/2020		
1		1 g in 1 TUBE; Type 0: Not a Combination Product			
2	NDC:69784-880-35	1 in 1 CARTON	04/15/2020		
2		3.5 g in 1 TUBE; Type 0: Not a Combination Product			

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA061212	04/15/2020		

Labeler - Woodward Pharma Services LLC (026749066)

Registrant - Woodward Pharma Services LLC (026749066)

Revised: 4/2020 Woodward Pharma Services LLC