HEALTH MART CHILDRENS PAIN AND FEVER- acetaminophen suspension Strategic Sourcing Services LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

McKesson Children's Pain & Fever Drug Facts

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
- the common cold
- flu
- headache
- sore throat
- toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

• with any other drug containing acetaminophen (prescription or nonprescription). If you are not

- sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has

liver disease

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

When using this product

do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

*or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each 5 mL contains: sodium 3 mg
- store at 20-25°C (68-77°F)
- do not use if printed neckband is broken or missing

Inactive ingredients

anhydrous citric acid, calcium sulfate, carrageenan, flavor, glycerin, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, propylparaben, purified water, sodium benzoate, sorbitol solution, sucralose, tribasic sodium phosphate

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Compare to Children's Tylenol® Oral Suspension active ingredient

SUSPENSION LIQUID

Children's Pain & Fever

Acetaminophen 160 mg per 5 mL

Pain Reliever/Fever Reducer

For ages 2 to 11 years

No high fructose corn syrup

Aspirin Free

Alcohol Free


Ibuprofen Free

Gluten Free

Dye Free

GRAPE FLAVOR

4 Fl oz (118 mL)



SUSPENSION LIQUID Children's Pain & Fever

Health Mart.

Acetaminophen 160 mg per 5 mL

Pain Reliever/Fever Reducer

Drug Facts

Active ingredient (in each 5 mL)

Purpose

.Pain reliever/fever reduce cetaminophen 160 mg

- Uses temporarily:

 reduces fever
 relieves minor aches and pains due to:
 the common cold
 flu headache
 sore throat
 toothache

Warnings

- warnings:
 Liver warning: This product contains acetaminophen.
 Severe liver damage may occur if your child takes
 more than 5 doses in 24 hours, which is the maximu daily amount
- dairy amount

 with other drugs containing acetaminophen

 Allergy alert: Acetaminophen may cause severe skin
 reactions. Symptoms may include:

 skin reddening

 ti a skin reaction occurs, stop use and seek medical help

right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

- with any other drug containing acetaminophen (prescription or nonprescription), If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

 if your child has ever had an allergic reaction to this
- product or any of its ingredients

Ask a doctor before use if your child has liver disease Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
 fever gets worse or lasts more than 3 days
 new symptoms occur
- redness or swelling is present
 These could be signs of a serious condition

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical ven if you do not notice any signs or symptoms.

Drug Facts (continued)

- Directions

 this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose

- warning)

 shake well before using

 mL = milliliter

 find right dose on chart below. If possible, use weight
- to dose; otherwise, use age.

 remove the child protective cap and squeeze your child's dose into the dosing cup

 repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

*or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each 5 mL contains: sodium 3 mg
 store at 20-25°C (68-77°F)
 do not use if printed neckband is broken or

Inactive ingredients

anhydrous citric acid, calcium sulfate, carrageenan, flavor, glycerin, hydroxyethyl cellulose, microcrystal cellulose and carboxymethylcellulose sodium, propylene glycol, propylparaben, purified water, sodium benzoate, sorbitol solution, sucralose, tribasic

Questions or comments? 1-800-719-9260

**This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Children's Tylenol® Oral Suspension.

MSKESSON Another Quality Product
Distributed by Molikeson
655 Stata Highway 161, Iving, TX 75039
Money Back Guarantee
healthmant.com



Compare to Children's Tylenol® Oral Suspension active ingredient**

SUSPENSION LIQUID

Children's Pain & Fever

Acetaminophen 160 mg per 5 mL

Pain Reliever/Fever Reducer

For ages 2 to 11 years No high fructose corn syrup

Aspirin Free • Alcohol Free Ibuprofen Free • Gluten Free

Dye Free



4 Fl oz (118 mL)



SUSPENSION LIQUID

Children's Pain & Fever

DIRECTIONS: STEP 1. Shake well before using



STEP 2. Remove child protective cap



STEP 3. Squeeze your child's dose into the dosing cup



IMPORTANT:

Please see dosing chart or ask a doctor for your child's correct dose.





HEALTH MART CHILDRENS PAIN AND FEVER

acetaminophen suspension

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:62011-0183

Route of Administration

ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINO PHEN (UNII: 36209 ITL9D) (ACETAMINO PHEN - UNII: 36209 ITL9D)	ACETAMINOPHEN	160 mg in 5 mL	

Inactive Ingredients		
Ingredient Name	Strength	
ANHYDRO US CITRIC ACID (UNII: XF417D3PSL)		
CALCIUM SULFATE, UNSPECIFIED FORM (UNII: WAT0 DDB505)		
CARRAGEENAN (UNII: 5C69 YCD2YJ)		
GLYCERIN (UNII: PDC6 A3C0 OX)		
MICRO CRYSTALLINE CELLULO SE (UNII: OP1R32D61U)		
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
PROPYLPARABEN (UNII: Z8IX2SC1OH)		
WATER (UNII: 059QF0KO0R)		
SODIUM BENZOATE (UNII: OJ245FE5EU)		
SORBITOL (UNII: 506T60A25R)		
SUCRALOSE (UNII: 96K6UQ3ZD4)		
SODIUM PHO SPHATE, TRIBASIC (UNII: A752Q30 A6 X)		

Product Characteristics			
Color	WHITE (viscous)	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

	Packaging			
1	# Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC:62011-0183-1	1 in 1 CARTON	0 1/17/20 13	
	1	118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	0 1/17/20 13	

Labeler - Strategic Sourcing Services LLC (116956644)