MAXIMUM STRENGTH MUCINEX SINUS-MAX PRESSURE, PAIN AND COUGHacetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, film coated RB Health (US) LLC

Maximum Strength Mucinex® Sinus-Max® Pressure, Pain & Cough Drug Facts

Active ingredients (in each caplet)	Purposes	
Acetaminophen 325 mg	Pain reliever	
Dextromethorphan HBr 10 mg	Cough suppressant	
Guaifenesin 200 mg	Expectorant	
Phenylephrine HCl 5 mg	Nasal decongestant	

Uses

- temporarily relieves:
 - nasal congestion
 - headache
 - minor aches and pains
 - cough
 - sinus congestion and pressure
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information

store between 20-25°C (68-77°F)

Inactive ingredients

croscarmellose sodium, crospovidone, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, talc, titanium dioxide

Questions?

1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.

Dist. by: RB Health (US) Parsippany, NJ 07054-0224

PRINCIPAL DISPLAY PANEL - 20 Caplet Blister Pack Carton

MAXIMUM STRENGTH NDC 63824-242-20

Mucinex® SINUS-MAX®

Actual Size

PRESSURE, PAIN & COUGH

FOR AGES 12+

Acetaminophen – Pain Reliever • Dextromethorphan HBr – Cough Suppressant Guaifenesin – Expectorant • Phenylephrine HCl – Nasal Decongestant

- Relieves Sinus Pressure & Congestion
- Relieves Headache
- Controls Cough
- Thins & Loosens Mucus

20 CAPLETS

Do not use a skin reaction occurs, stop use and seek medica

high blood pressure thyroid

inestions:

-866-MUCINEX (1-866-682-4639)

may also report side effects to this phone

EXP.:

麗

TOT

98659TE MADE IN: smoking, asthma, chronic bronchills, or emphysem

OCCURS WITH

Ask a doctor before use if you have doctor or pharmacist before taking this produc your prescription drug contains an MAOI, ask a stopping the MAOI drug. If you do not know if heartdisease Idiabetes

If you are now taking a prescription monoamine with any other drug containing acetaminophen ask a doctor or pharmacist oxidase inhibitor (MAOI) (certain drugs for (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen,

depression, psychiatric, or emotional conditions or Parkinson's disease), or for 2 weeks after

Other information store at 20-25°C (68-77°F)

children under 12 years of age: do not use

over: take 2 caplets every 4 hours

cellulose, polyethylene glycol, polyvinyl alcohol povidone, talc, titanium dioxide no. 40 aluminum lake, FD&C yellow no. 6 aluminun croscarmellose sodium, crospovidone, FD&C red ake, magnesium steerate, microcrystalline nactive ingredients amings

keep out of reach of children. professional before use.

f pregnant or breast-feeding, ask a health

Serious condition.

3 or more alcoholic drinks daily while using Severe liver damage may occur if you take: more than 12 caplets in 24 hours, which is the Liver warning: This product contains acetaminoph with other drugs containing ac etaminophen maximum daily amount

skin reactions. Symptoms may include: lergy alert: Acetaminophen may cause severe ■ blisters rash

damage. In case of overdose, get medical help o Overdose warning: Taking more than file recommended dose (overdose) may cause do not take more fran directed (see children even if you do not notice any signs or nedical attention is critical for adults as well as for ontact a Poison Control Center right away. Quick do not take more than 12 caplets in any 24-hour period adults and children 12 years of age and Overdose warning

Maximum Strength per 4-hour dose

Tamper evident: Do not use if carton is damaged or if printed seal on blister is broken or missing.

pain, nasal congestion, or cough gets worse or

asts more than 7 days

nervousness, dizziness, or sleeplessness occur

temporarily relieves

i helps loosen phlegm (mucus) and thin bronchia

bothersome mucus and make coughs secretions to rid the bronchial passageways of temporarily promotes nasal and/or sinus drainage

 sinus congestion and pre nasal congestion

minor aches and pains

■ cough

new symptoms occur

cough comes back, or occurs with fever, rash,

neadache that lasts. These could be signs of a

redness or swelling is present

fever gets worse or lasts more than 3 days

Drug Facts

in each caplet, active ingredients

Purposes

cough that occurs with too much philegm (mucus

| Facts

(continued

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more

NDC 63824-242-20

omethorphan HBr 10 mg

Keep carton for complete product information.

PRESSURE, PAIN & COUGH

Mucinex

SINUS-MAX®

Actual Size

PRESSURE, PAIN & COUGH

Acetaminophen - Pain Reliever • Dextromethorphan HBr - Cough Suppressant Guaifenesin - Expectorant • Phenylephrine HCI - Nasal Decongestant

Relieves Sinus Pressure & Congestion

Relieves Headache

Controls Cough

Thins & Loosens Mucus

CAPLETS



FOR AGES 12+

Dist. by: RB Health (US) Parsippany, NJ 07054-0224 2020 RB Health

www.mucinex.com Patents: www.rb.com/patents 092520 3165986



MAXIMUM STRENGTH



PARENTS: www.StopMedicineAbuse.org

MAXIMUM STRENGTH

MAXIMUM STRENGTH MUCINEX SINUS-MAX PRESSURE, PAIN AND

COUGH

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, film coated

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63824-242	
Route of Administration	ORAL			

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg	
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS 297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg	

Inactive Ingredients	
Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
CROSPOVIDONE (UNII: 2S7830E561)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics				
Color	red	Score	no score	
Shape	OVAL	Size	20mm	
Flavor		Imprint Code	VVV;MSC	
Contains				

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:63824- 242-20	2 in 1 CARTON	05/01/2018		
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC Monograph Drug	M012	05/01/2018			

Labeler - RB Health (US) LLC (081049410)

Revised: 12/2024 RB Health (US) LLC