

LIDOCAINE HYDROCHLORIDE - lidocaine injection

Agri Laboratories, Ltd.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

LIDOCAINE HYDROCHLORIDE INJECTION 2% 20 mg/mL

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

FOR ANIMAL USE ONLY

KEEP OUT OF REACH OF CHILDREN

INDICATIONS

Lidocaine is a potent local anesthetic for producing epidural, nerve conduction and infiltration anesthesia.

CONTRAINDICATIONS

Lidocaine is contraindicated in animals with a known hypersensitivity to the drug.

PRECAUTIONS

Lidocaine is usually well tolerated. Nevertheless, as with all local anesthetics, untoward effects may occur due to hypersensitivity, faulty technique, overdosage and inadvertent intravascular or subarachnoid injection. In case of respiratory arrest, immediate resuscitation with oxygen is indicated.

DOSAGE AND ADMINISTRATION

Epidural

Cattle and Horses- 5 to 15 mL

Dogs and Cats- 1 mL per 10 pounds of body weight.

Nerve Block

Cattle and Horses- 5 to 20 mL

Infiltration

Dilute to 0.5% concentration (1 mL of 2% solution diluted with 3 mL of sterile water = 4 mL of a 0.5% solution)

COMPOSITION

Each mL of sterile aqueous solution contains:

Lidocaine Hydrochloride.....2.0%

Propylene Glycol.....5.2%

Sodium Chloride.....0.5%

Sodium Lactate.....0.5%

with Methylparaben 0.15%, Sodium Metabisulfite 0.10%,

Propylparaben 0.03%, and Disodium Edetate 0.001% as preservatives.

Store at controlled room temperature between 15° and 30°C (59° - 86°F)

TAKE TIME OBSERVE LABEL DIRECTIONS

INDICATIONS: Lidocaine is a potent local anesthetic for producing epidural, nerve conduction and infiltration anesthesia.

CONTRAINDICATIONS: Lidocaine is contraindicated in animals with a known hypersensitivity to the drug.

PRECAUTIONS: Lidocaine is usually well tolerated. Nevertheless, as with all local anesthetics, untoward effects may occur due to hypersensitivity, faulty technique, overdosage and inadvertent intravascular or subarachnoid injection. In case of respiratory arrest, immediate resuscitation with oxygen is indicated.

DOSAGE AND ADMINISTRATION:

Epidural

Cattle and Horses - 5 to 15 mL

Dogs and Cats - 1 mL per 10 pounds of body weight.

Nerve Block

Cattle and Horses - 5 to 20 mL

Infiltration

Dilute to 0.5% concentration (1mL of 2% solution diluted with 3 mL of sterile water = 4 mL of a 0.5% solution).

L-2000-04

Rev. 05-08

Lot

Exp

NDC 57561-221-04

**LIDOCAINE
HYDROCHLORIDE
INJECTION 2%
20 mg/mL**

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**FOR ANIMAL USE ONLY
KEEP OUT OF REACH OF CHILDREN**

NET CONTENTS: 100 mL



COMPOSITION: Each mL of sterile aqueous solution contains:

Lidocaine Hydrochloride 2.0%
Propylene Glycol 5.2%
Sodium Chloride 0.5%
Sodium Lactate 0.5%

with Methylparaben 0.15%,
Sodium Metabisulfite 0.10%,
Propylparaben 0.03% and
Disodium Edetate 0.001% as
preservatives.

**Store at controlled room
temperature between 15°
and 30°C (59° - 86°F).**

TAKE TIME  OBSERVE LABEL DIRECTIONS

**Manufactured for
Agri Laboratories, Ltd.
St. Joseph, MO 64503**

© Registered trademark of Agri
Laboratories, Ltd.



LIDOCAINE HYDROCHLORIDE

lidocaine injection

Product Information

Product Type	PREScription ANIMAL DRUG	Item Code (Source)	NDC:57561-221
Route of Administration	EPIDURAL, INFILTRATION		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE HYDROCHLORIDE ANHYDROUS	0.02 g in 1 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57561-221-04	100 mL in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		08/21/2008	

Labeler - Agri Laboratories, Ltd. (155594450)

Revised: 6/2013

Agri Laboratories, Ltd.