

CMPK2- cmpk2 liquid
Radix Laboratories, Inc.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

CMPK2

• **CONTENTS (grams/500ml):**

- Calcium10.0g
- (Calcium Borogluconate)
- Phosphorus.....6.03g
- (Sodium Hypophosphite)
- Magnesium.....2.76g
- (Magnesium Chloride Hexahydrate)
- Potassium.....0.525g
- (Potassium Chloride)
- Dextrose Monohydrate.....50.0g

Indications:

- For use as a supplemental nutritive source of calcium, phosphorus, magnesium, potassium and dextrose in cattle.

Storage:

- Store at controlled room temperature between 59°F to 86°F (15°C -30°C).

Warnings:

- Avoid storage at cold temperatures which can cause product to become cloudy.
- To clarify place product in warm water bath (100°F-120°F) for about 30 minutes.

Dosage and administration:

- Administer orally as a drench or by use of a stomach tube.
- The usual recommended dose in adult cattle is 16.9 fl. oz (500mL) to 33.8 fl.oz (1000mL) depending on size and condition.
- Keep away from children.

Current CMPK2 Label

CONTENTS (grams/500ml):

Calcium (Calcium borogluconate)	10.0g
Phosphorus (Sodium hypophosphite)	6.03g
Magnesium (Magnesium chloride hexahydrate)	2.76g
Potassium (Potassium chloride)	0.525g
Dextrose monohydrate	50.0g
Purified water	q.s.
Inactives: Methylparaben, Propylparaben, Propylene Glycol	

MANUFACTURED BY:

Radix Laboratories, Inc.
1334 International Drive
Eau Claire, WI 54701
1-800-927-2349



NDC 50480-052-50



**CMPK
2**

TAKE TIME



READ LABEL INSTRUCTIONS

**Keep away from children
FOR ANIMAL USE ONLY**

**NET CONTENTS
16.9 fl. oz. (500ml)**

WARNING:

Avoid prolonged storage at cold temperatures, which can cause preservatives to precipitate. To clarify, place product in warm water bath (100°F to 120°F) for about 30 minutes.

INDICATIONS:

For use as a supplemental nutritive source of calcium, magnesium, phosphorus, potassium and dextrose in cattle.

DOSAGE AND ADMINISTRATION:

Administer orally as a drench or by use of a stomach tube. The usual recommended dose in adult cattle is 16.9 fl. oz. (500ml) to 33.8 fl. oz. (1000ml) depending on size and condition.

STORAGE:

Store at room temperature between 59°F and 86°F (15°C to 30°C)

Product Image



CMPK2

cmpk2 liquid

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:50480-052
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CALCIUM BORO GLUCONATE (UNII: 6915G2PQ10) (CALCIUM CATION - UNII:2M83C4R6ZB)	CALCIUM CATION	2 g in 100 mL
SODIUM HYPOPHOSPHITE (UNII: 8TU1537O43) (HYPOPHOSPHITE ION - UNII:238U65NZ04)	PHOSPHATE ION	1.2 g in 100 mL

MAGNESIUM CHLORIDE (UNII: 02F3473H9O) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM CATION	0.552 g in 100 mL
POTASSIUM CHLORIDE (UNII: 660YQ98I10) (POTASSIUM CATION - UNII:295O53K152)	POTASSIUM CATION	0.105 g in 100 mL
DEXTROSE MONOHYDRATE (UNII: LX22YL083G) (ANHYDROUS DEXTROSE - UNII:5SL0G7R0OK)	DEXTROSE MONOHYDRATE	10 g in 100 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50480-052-50	12 in 1 CARTON		
1		500 mL in 1 BOTTLE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/27/2013	

Labeler - Radix Laboratories, Inc. (108500091)

Revised: 3/2014

Radix Laboratories, Inc.