

**WALGREENS GENTLE LAXATIVE- bisacodyl suppository suppository  
Walgreens**

-----  
**Walgreens® Gentle Laxative 8ct**

**Active Ingredient (in each Suppository)**

Active Ingredient (in each Suppository)	Purpose
Bisacodyl 10 mg.....	Stimulant Laxative

**Uses**

- for relief of occasional constipation
- this product generally produces bowel movement in 1/4 to 1 hour

**Warnings**

For rectal use only.

**Do not use**

- Do not use
- when abdominal pain, nausea, or vomiting are present
  - for a period of longer than 1 week

**Ask a doctor**

Ask a doctor before use if you have noticed a sudden change in bowel habits that lasts over a period of 2 weeks

**When using this product**

When using this product it may cause abdominal discomfort, faintness, rectal burning and mild cramps

**Stop use and ask a doctor if**

- Stop use and ask a doctor if
- you have rectal bleeding
  - you fail to have a bowel movement after using this product.

This may indicate a serious condition

### **If pregnant or breast-feeding**

If pregnant or breast-feeding, ask a health professional before use.

### **Keep out of reach of children**

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

### **Directions**

- detach one suppository from the strip
- remove wrapper before inserting into the rectum

adults and children 12 years of age and older	1 suppository once daily
children 6 to under 12 years	1/2 suppository once daily
children under 6 years	do not use

### **Other information**

Other information

- Store at room temperature. Do not exceed 30 °C (86 °F)
- Lot No. & Exp. Date: see wrapper or box

### **Inactive ingredient**

Inactive ingredient hydrogenated vegetable oil

### **Directions**

- detach one suppository from the strip
- remove wrapper before inserting into the rectum

adults and children 12 years of age and older	1 suppository once daily
children 6 to under 12 years	1/2 suppository once daily
children under 6 years	do not use

### **Label**

Walgreens

Relieves constipation

ORF0418-F2

ACTUAL SIZE



# Gentle Laxative Suppositories

Walgreens

Compare to Dulcolax® active ingredient<sup>††</sup>

NDC 0363-0441-08

BISACODYL USP 10 mg

- Prompt, predictable relief of constipation

8 SUPPOSITORIES

DISTRIBUTED BY: **WALGREEN CO.**  
 200 WILMOT RD., DEERFIELD, IL 60015  
**100% SATISFACTION GUARANTEED**  
 walgreens.com  
 ©2018 Walgreen Co.

<sup>†</sup>Walgreens Pharmacist Survey  
<sup>††</sup>This product is not manufactured or distributed by Chatterm, Inc., owner of the registered trademark Dulcolax®.



Health expertise you rely on.<sup>™</sup>

**Drug Facts** (continued)

**Directions**

- detach one suppository from the strip
- remove wrapper before inserting into the rectum

**Other information**

adults and children	1 suppository
12 and over	once daily
children 6 to 11 years	½ suppository
children 5 years and under	once daily
children under 6 years	do not use

**Warnings**

- this product generally produces bowel movement in ½ to 1 hour
- for relief of occasional constipation
- do not use
  - when abdominal pain, nausea or vomiting are present
  - for a period longer than 1 week
- ask a doctor before use if you have noticed a sudden change in bowel habits that lasts over a period of 2 weeks

**Do not use**

- for rectal use only

**When using this product** it may cause abdominal discomfort, flatness, rectal bleeding and mild cramps

**Stop use and ask a doctor if**

- you have rectal bleeding
- you feel to have a bowel movement after using this product

**Ask a doctor before use if you have** noticed a serious condition

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Drug Facts** (continued)

**Active ingredient** (in each suppository)  
 Bisacodyl 10 mg

**Purpose**  
 Stimulant Laxative

**Uses**  
 Relief of occasional constipation

**Warnings**  
 This product generally produces bowel movement in ½ to 1 hour

**Do not use**  
 when abdominal pain, nausea or vomiting are present  
 for a period longer than 1 week  
 ask a doctor before use if you have noticed a sudden change in bowel habits that lasts over a period of 2 weeks

**When using this product** it may cause abdominal discomfort, flatness, rectal bleeding and mild cramps

**Stop use and ask a doctor if**  
 you have rectal bleeding  
 you feel to have a bowel movement after using this product

**Ask a doctor before use if you have** noticed a serious condition

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

ITEM 324579 W00000-0000-0



3 119171 6262 1

bisacodyl suppository suppository

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-0466
<b>Route of Administration</b>	RECTAL		

### Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	10 mg

### Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
PALM KERNEL OIL (UNII: B0S90M0233)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	
<b>Shape</b>	BULLET	<b>Size</b>	32mm
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0466-08	2 in 1 PACKAGE	09/18/2014	
1		4 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M007	09/18/2014	

**Labeler** - Walgreens (008965063)

**Registrant** - Unipack LLC (116015769)

### Establishment

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
Unipack LLC		009248480	manufacture(0363-0466)

